

Final Report

Title: Planetary Geophysics and Tectonics

PI: E. Marc Parmentier

Period of Award: 11/1/98-10/31/02

Recipient's Institution:

**Brown University
Box 1929
Providence, RI 02912**

Grant Number: NAG5-3659

Models for the thermal and chemical evolution of the Moon

We have carried out several studies that explore explanations for the role of chemical density variations in Moon's evolution. Meaningful models for the evolution of the Moon must explain a number of important magmatic characteristics. Volcanic activity subsequent to the formation of its anorthositic crust was dominated by the eruption of mare basalt. 1) The main phase of mare volcanism began ~500 Myr after the crystallization of the anorthositic crust and continued for ~1Gyr. 2) The picitic glasses, considered to be representative of mare basalt least affected by low pressure, near-surface fractionation, were generated by melting, at 400-600 km depth, of a source containing components that, on the basis of the magma ocean hypothesis, should have crystallized at much shallower depth during fractionation of the anorthositic crust. 3) Mare basalts occur primarily in one region of the Moon. Recent topographic data demonstrate that the earlier idea that mare basalt flooded areas of low elevation is not correct. Large areas of very low elevation do not contain mare basalt. The hemispheric asymmetry of mare basalt distribution on the lunar surface must be explained in some other way (Zhong, et al. 2000). 4) A region of the surface roughly correlating with that containing mare basalts also is thought to contain high subsurface concentrations of KREEP which was excavated during the formation of large impact basins. This so-called Procellarum KREEP Terrane (PKT) is responsible for the Imbrium basin-centered thorium anomaly mapped by Lunar Prospector.

A model in which dense ilmenite and pyroxene-rich cumulates (IC), that crystallize during the solidification of the last 5-10% of a magma ocean, sink into the underlying mantle may explain important aspects of lunar evolution. The IC may carry with it high concentrations of incompatible elements, including heat producing U, Th, and K, that would have become progressively enriched in the residual liquid as the magma ocean crystallized. We suggested that dense IC cumulates may have differentiated to form a core or a layer surrounding an existing metallic core. It is interesting to note that a recent inversion of lunar free oscillation data indicates a lunar core with a density appropriate for IC, rather than metal. Subsequent thermal expansion due to radiogenic heating would make this material buoyant, giving rise to upwelling and melting (Zhong, et al. 2000). This model can thus potentially explain the timing of mare volcanism and, if this upwelling developed at spherical harmonic degree-one, why mare volcanism is concentrated in one large area of the surface.

KREEP-rich materials excavated during the formation of the Imbrium basin must have pre-existed mare basalts that fill the basin. Both the concentration of KREEP-rich material beneath the PKT and later mare volcanism in the same region due to the upwelling and melting of initially deep IC cumulates are not a natural consequence of a model of buoyant upwelling of a core or deep IC layer unless some preconditioning of the mantle beneath this region determined the location of later upwelling.

It is thus important to explore mechanisms which may regionally concentrate a KREEP layer early in lunar evolution and which ~400-500 Myr later will give rise to mare basalt generation beneath the *same* region. It has been suggested that melting due to radioactive heating of a thickened KREEP layer beneath the PKT may give rise to mare volcanism in that region. We explored (Parmentier, et al., 2002) whether gravitational differentiation of a layer containing IC cumulates is a possible mechanism to explain the concentration of KREEP beneath the PKT. Thickening of the KREEP layer beneath the PKT during differentiation of the IC cumulates and the later partial melting of these cumulates due to radiogenic heating could have generated the mare basalts that were localized in the region of the PKT. In this case, differentiated IC cumulates may never have reached depths greater than that of the deepest mare basalt source. However, for this explanation to be valid, differentiation of IC cumulates in a relatively thin layer beneath the anorthositic crust must have occurred at a long wavelength (spherical harmonic degree-one); and the rate of differentiation and the rate radiogenic heating must be of an appropriate magnitude to explain the timing and depth of origin of mare basalts. We also showed (Hess and Parmentier, 2001) that a KREEP layer formed over the whole Moon concentrated at shallow depth beneath the PKT would have thermal and magmatic consequences that are not observed.

To understand the conditions for which this is possible, we examined the Rayleigh-Taylor instability of a dense layer near the surface of a sphere, representing the lunar mantle, in which the viscosity varies with depth (Parmentier, et al, 2002). Our results show that spherical harmonic degree-one is the fastest growing wavelength of instability if the viscosity of the dense layer is sufficiently low relative to that of the deeper mantle or if the viscosity of the mantle increases with depth. However, the viscosity increase cannot be distributed over a depth that is too large. We showed that with a temperature dependent viscosity, that the depth over which radioactivity in the IC will heat the mantle, thus reducing the viscosity, is consistent with the timing of mare basalt eruption.

Convective cooling of a compositionally stratified viscous fluid

Convective cooling of a stably stratified viscous fluid with strongly temperature dependent viscosity is important in understanding the thermal and chemical evolution of planets. Several different mechanisms may lead to an initially stably stratified mantle. The overturn of initially gravitationally unstable lunar magma ocean cumulates could lead to a stably, compositionally stratified mantle (Hess and Parmentier, 1995). This could result in a continuous Mg/Fe and density stratification over a depth comparable to that of the original magma ocean and/or to a more sharply defined layer of very dense, highly radiogenic late stage ilmenite-rich cumulates. Alley and Parmentier (1998) describe numerical experiments on the heating from below of an initially stable compositional stratification that provides a basis to test the role of compositional stratification in the evolution of the Moon. Zhong, et al. (2000) examined a layer surrounding a metallic core or a central core of dense, heating producing element-rich silicate material as discussed above.

The formation of crust by decompression melting in upwelling mantle would also result in a stably stratified mantle. The amount of melting in an upwelling parcel of mantle increases the height over which it decompresses, and increasing depletion of melt decreases the density of the residual mantle. Therefore, the progressive accumulation of buoyant mantle, depleted as a

consequence of crustal genesis by partial melting, should result in stable stratification of mantle beneath the lithosphere. We have previously discussed the possible role of such a buoyant depleted mantle layer in Venus (Parmentier and Hess, 1992). Recent studies have pointed out that the viscosity increase due to depleting small amounts of intergranular water along with melt may be larger than originally thought, perhaps as large as a factor of 300.

For Mars, SNC meteorites record an early magmatic differentiation. Even though Mars is a larger planet than the Moon and therefore expected to have a longer, more active convective evolution, isotopic heterogeneity recorded in SNCs is as large as those that have been found on Moon. Rb-Sr and Sm-Nd isotopes preserve a record of a major depletion event in the mantle, indicating mantle has resisted homogenization. Lu-Hf and rare earth element distributions reflect small degrees of melting of a source with residual garnet, corresponding to a depth in Mars of at least 250 km. If Martian crust, making up approximately 4% of total planetary volume, were all generated by adiabatic decompression melting, a significant compositional stratification of the top of the mantle would result. This amount of crust would require about 20% melting of the upper 200 km of Mars. More likely lower degrees of melt distributed over greater depths. How much modal cpx and garnet are represented in a chondritic material? 20% melting of garnet or spinel bearing mantle would create a density change equivalent to a temperature change of almost 500°C. Could inhibit convective heat transfer in the mantle. As discussed below, a low mantle heat flux may help explain relatively low temperatures at the base of the crust required to explain large elastic thicknesses and the long-term preservation of crustal thickness variations (Parmentier and Zuber, 2001; 2002). It is thus plausible that planets are initially stably stratified over substantial depths and that the density variations are large.

We have carried out numerical experiments examining the instability and subsequent mixing in a stably stratified fluid layer cooled from above (Zaranek and Parmentier, 2003). The objective is to use numerical experiments to derive scaling relationships that can be applied to planetary evolution – not detailed models of a particular planet or evolutionary scenario. We consider a 2D cartesian domain containing a Boussinesq fluid with infinite Prandtl number. On the top, no-slip boundary of this domain, temperature is set at zero. The free-slip bottom boundary is adiabatic and the solutions are horizontally periodic. With no volumetric heating or heating from below, cooling of the fluid, initially at a uniform temperature T_i , creates convective instability that is opposed by a stable compositional density stratification with a linear gradient

$$\gamma = \frac{1}{\rho} \frac{d\rho}{dz}$$

Composition advects but does not diffuse. For thermally activated creep with a large activation energy Q , viscosity increases sharply with decreasing temperature thus significantly reducing the thermal buoyancy in the boundary layer at the top of the fluid available to drive convective motions. We consider an exponentially temperature dependent viscosity

$$\mu(T) = \mu_i \exp(\varepsilon(T_i - T)/T_i)$$

With $\varepsilon = Q/RT_i$, where Q is the activation energy, this relationship provides a good approximation to the Arrhenius law with if the temperature range is not too large. Since viscosity variations as large as indicated by creep activation energies are difficult or impossible to treat with high numerical resolution, the scaling laws that we derive for realizable viscosity variations can be extrapolated to stronger temperature dependence.

Convective instability begins with the initial penetration of cold plumes to a depth determined by the compositional stratification. Subsequently formed plumes penetrate to greater depth resulting in a mixed layer that cools and thickens with time. The cooling and thickening of mixed layer makes it less buoyant than fluid at the top of the underlying, still hot and stratified

fluid. Continuity of density across the boundary between the bottom of the mixed layer and the top of the underlying fluid gives

$$(z_{mixed} - z_L) = \frac{2\alpha(T_i - T_m)}{\gamma}$$

where z_{mixed} and z_L are the mixed layer and conductive lid thicknesses respectively, T_m is the mean temperature in the mixed layer, and α is the thermal expansion of the fluid. This relationship is satisfied well by numerical experiments over a range of conditions, indicating that the mixed layer thickens mainly by buoyant, rather than purely viscous, entrainment.

Cooling of the convecting mixed layer is determined by heat conduction through the lid and by the heatflux delivered convectively to its base

$$f = ck \left(\frac{\rho\alpha g}{\mu(T_m)\kappa} \right)^{1/3} \Delta T_c^{4/3}$$

This relationship follows from earlier work (e.g. Grasset and Parmentier, 1998) where $\Delta T_c = 2.23T_i/\varepsilon$ is the temperature over which the viscosity increases by a factor of ten. Cooling of the mixed layer, assuming that it thickens fast enough that no heat is conducted into it from the still stratified region beneath, is given by the energy balance

$$\frac{d}{dt}(\rho c_p (T_i - T_m) z_{mixed}) = f.$$

Combining these equations to eliminate f and T_m gives the change in mixed layer thickness with time

$$\frac{d(z_{mixed} - z_L)^2}{dt} = \kappa_M = ck \left(\frac{\rho\alpha\Delta T_c g}{\mu(T_m)\kappa} \right)^{1/3} \left(\frac{\alpha\Delta T_c}{\gamma} \right).$$

The mixed layer thickens approximately as the squareroot of time so that the expression on the right side of the above equation can be considered an effective diffusivity for mixing κ_M . Numerical experiments agree well with this predicted scaling which accounts very well for the dependence of mixed layer thickness on γ and ε . We find values of c in the range of 0.130 to 0.159 consistent with that for laboratory and numerical experiments with no compositional stratification. Some of our numerical experiments also indicate that the mixed layer stops thickening, possibly due to the development of a separately convecting layer beneath it. We are continuing to study the mechanisms and scaling of the behavior.

Mars thermal evolution

Variations in the thickness of a uniform density crust cause lateral pressure gradients that generate flow in a ductile lower crust from areas of larger to smaller crustal thickness, thus evening out the thickness of the crust over time. Analysis of gravity and topography data for Mars provide estimates of crustal thickness showing long wavelength lateral variations present even beneath features of great age. The longest wavelength present is the north-south crustal thickness variation, but it is not clear that some of the apparent crustal thickness variation (derived assuming a constant density crust) does not reflect variations in crustal density. Alternatively, the Tharsis Rise appears to be a large center of basaltic volcanism, that began forming early in the evolution of Mars, with topography supported largely by thickened crust, presumably of relatively uniform composition. Other features such as large basins, like Hellas, while apparently younger than Tharsis, preserve shorter wavelength crustal thickness variations.

In this study we are exploring the implications that the survival of crustal thickness variations would have for the thermal evolution of Mars.

Previous studies have noted that the preservation of crustal thickness variations places constraints on either the thickness of the crust through the temperature dependence of crustal viscosity and previous studies have investigated constraints on crustal thickness using steady-state thermal evolution models which assume a heatflux at the base of the crust equal to the radiogenic heat production in the mantle. With a dry diabase flowlaw, crustal thickness as large as 100 km will allow the preservation of crustal thickness variations.

Since the rheology of the lower crust is likely to very strongly temperature dependent, this places important constraints on thermal evolution. We have formulated thermal models that include both radiogenic heating and secular cooling. These models treat the strong temperature dependence of flow by thermally activated creep and the presence of conductive lid above a convecting interior using the results of Grasset and Parmentier (1998). Our results (Parmentier and Zuber, 2001; 2002) indicate that the contribution of the heat flux from secular cooling of the core and mantle to temperatures at the base of crust is very significant. For initial temperatures for subsolidus cooling in the range of 1500 – 2000°C, and with a crustal thickness as thin as 50 km and with laboratory flow laws representing a range of plausible crustal compositions, both wet and dry, we find that crustal thickness variations even with a lateral scale as large as Tharsis cannot be preserved for times on the order of 4 Gyr. We suggest two mechanisms that can reduce the temperatures of the lower crust thus preserving crustal thickness variations without requiring a very thin crust: compositional stratification of the mantle (Zaranek and Parmentier, 2003) and hydrothermal cooling of the crust.

If the early crust formed by adiabatic decompression melting, for example in mantle plumes, the extraction of melt to form crust would leave behind a compositionally buoyant, depleted mantle, as discussed above. The results reported above show that stable compositional stratification due to crustal genesis may significantly inhibit convective instability at the top of the mantle. An estimate of the relaxation rate of crustal thickness variations assuming only the conduction of heat from a non-convecting mantle is shown in Figure 4a. Relaxation rates are calculated for a dry diabase rheology following the approach outlined by Parmentier and Zuber (2001, 2002). Only low values of the fraction of chondritic heat sources fractionated into the crust would give relaxation rates smaller than 10^{-17} /sec as required to preserve crustal thickness variations over the last ≈ 4 Gyr. This appears possible only if the crustal heating fraction is less than about 20%. Noachian elastic lithosphere thicknesses as large as 50 km (McGovern, et al. 2001) would also require a relatively low surface heatflux of about 40-50 mW/m².

As an alternative to low mantle heat flux, a porous, permeable regolith, if water saturated, may allow hydrothermal convective cooling of upper crust. On the Moon, low seismic velocity in the upper 25 km of the crust is thought to reflect fractured, porous rock with a porosity of about 1%. An impact-fractured and brecciated layer should also have been present on Mars perhaps with a similar porosity to that on the Moon. The bottom of such a porous layer is determined by the close of fractures under hydrostatic stress, then the layer thickness on Mars should be about 1/3 that on the Moon. Hydrothermal cooling is determined a Rayleigh number that depends on the product of the permeable and layer thickness and the physical properties of water. Assuming that fractures occur a planar orthogonal network of cracks, the permeability is determined by the product of the porosity and the square of the crack spacing. For fractures spaced 1 cm apart, perhaps typical of the clast sizes in lunar breccia samples, and using the appropriate physical properties of water, values of Nu (Nusselt number), the ratio hydrothermal convective heat transfer to purely conductive heat flux, would about a factor of ten. For Nu=10 and an 8 km layer thickness, the relaxation time for a mantle heatflux expected early in Martian history (60-80 mW/m²) is small enough to preserve crustal thickness variations.

Hydrothermal alteration may also explain the difference in the intensity of magnetic anomalies between the southern highlands and northern plains. If crustal mineralogy is

responsible for remnant magnetization and if hydrothermal alteration extends to the same depth in both the highlands and the plains, then thicker crust in the highlands would result in a thicker, unaltered, still magnetized layer. Hydrothermally altered minerals may also fill cracks reducing the permeability. Hydrothermal cooling may thus be most effective when impacts are frequent enough to continually generate new fractures.

Published papers:

- O. Grasset and E.M. Parmentier, Thermal convection in a volumetrically heated, infinite Prandtl number fluid with strongly temperature-dependent viscosity: implications for planetary thermal evolution, *J. Geophys. Res.*, 103, 18171-18181, 1998.
- K.M. Alley and E.M. Parmentier, Thermal convection in an initially stratified fluid heated from below: application the early evolution of planets and particularly the Moon, *Phys. Earth Planet. Int.*, 108, 15-32, 1998.
- S. Zhong, E.M. Parmentier, M.T. Zuber, A dynamic origin for the global asymmetry of Lunar mare basalts, *Earth Planet. Sci. Lett.* 177, 131-140, 2000.
- P.C. Hess and E.M. Parmentier, Thermal Evolution of a Thicker KREEP Liquid Layer, *J. Geophys. Res.*, 106, 28023-28032, 2001.
- E.M. Parmentier, S. Zhong, and M.T. Zuber, Gravitational differentiation due to initial chemical stratification: origin of lunar asymmetry by the creep of dense KREEP, *Earth Planet. Sci. Lett.* 201, 473-480, 2002.
- S. Zarnek and E.M. Parmentier, Convective cooling of an initially stably stratified fluid with temperature-dependent viscosity including applications to planetary evolution, submitted to *J. Geophys. Res.* (2003).
- L.T. Elkins Tanton, E.M. Parmentier, and P.C. Hess, Magma ocean fractional crystallization and cumulate overturn in terrestrial planets: implications for Mars, submitted to *Meteoritics and Planetary Science* (2003).

Abstracts of recent and ongoing work:

- E.M. Parmentier and P.C. Hess, On the chemical differentiation and subsequent evolution of the Moon, *Lunar Planet. Sci.* 30, 1289, 1999.
- P.C. Hess and E.M. Parmentier, Asymmetry and timing of mare volcanism, *Lunar Planet. Sci.* 30, 1360, 1999.
- S. Zhong, E.M. Parmentier, and M.T. Zuber, Early Lunar evolution and origin of asymmetric distribution of mare basalts, *Lunar Planet. Sci.* 30, 1789, 1999.
- G. Choblet, O. Grasset, E.M. Parmentier, and C. Sotin, Mars thermal evolution revisited, *Lunar Planet. Sci.* 30, 1556, 1999.
- Parmentier, E.M., S. Zhong, and P.C. Hess, Asymmetric evolution of the Moon: a possible consequence of chemical differentiation, Workshop on New Views of the Moon, September, 1999.
- Choblet, G., O. Grasset, E.M. Parmentier, and C. Sotin, Mars thermal evolution revisited, *Lunar Planet. Sci.* 30, 1556, 1999.
- Parmentier, E.M., S. Zhong, and M.T. Zuber, On the relationship of chemical differentiation and the origin of lunar asymmetries, *Lunar Planet. Sci.* 31, 1614, 2000.
- Hess, P.C., and E.M. Parmentier, Evolution of the Moon with a thickened KREEP layer, *Lunar Planet. Sci.* 31, 1659, 2000.
- Spaun, N., E.M. Parmentier and J.W. Head, Modelling Europa's crustal evolution: Implications for an ocean, *Lunar Planet. Sci.* 31, 1039, 2000.
- Hess, P.C. and E.M. Parmentier, Implications for cumulate overturn on Mars, *Lunar Planet. Sci.* 32, 1319, 2001
- Parmentier, E.M., S. Zhong, and M.T. Zuber, Gravitational differentiation of an initially unstable chemical stratification: origin of lunar asymmetries, *Lunar Planet. Sci.* 32, 1329, 2001.
- Parmentier, E.M. and M.T. Zuber, Relaxation of crustal thickness variations on Mars: implications for thermal evolution, *Lunar Planet. Sci.* 32, 1357, 2001.
- Sotin, C., J.W. Head, and E.M. Parmentier, Convective pattern and type of volcanism on Mars, *Eos Trans. AGU* 82(47) *Fall Meeting Supplement*, abstract P31B-02, 2001.
- Choblet, G., E.M. Parmentier, and C. Sotin, Thermal boundary layer instabilities for a fluid heated from below and from within, *Eos Trans. AGU* 82(47) *Fall Meeting Supplement*, abstract T21A-0866, 2001.
- Parmentier, E.M. and M.T. Zuber, Preservation of crustal thickness variations on Mars: mantle compositional stratification or hydrothermal crustal cooling?, *Lunar Planet. Sci.* 33, 1737, 2002.
- Zarnek, S., and E.M. Parmentier, Convective cooling of an initially stably stratified fluid: implications for Martian planetary evolution, *Lunar Planet. Sci.* 33, 1580, 2002.
- Sotin, C., J.W. Head, and E.M. Parmentier, Models for Mars early evolution: formation of Tharsis and cessation of magnetic field, *Lunar Planet. Sci.* 33, 1922, 2002.