

# Evaluation and Validation of Case 2 Algorithms in Chesapeake Bay

Lawrence W. Harding, Jr.  
Andrea Magnuson

Horn Point Laboratory  
University of Maryland  
Center for Environmental Science  
P.O. Box 775  
Cambridge, MD 21613

## INTRODUCTION

The high temporal and spatial resolution of satellite ocean color observations will prove invaluable for monitoring the health of coastal ecosystems where physical and biological variability demands sampling scales beyond that possible by ship. However, ocean color remote sensing of Case 2 waters is a challenging undertaking due to the optical complexity of the water. The focus of this SIMBIOS support has been to provide *in situ* optical measurements from Chesapeake Bay (CB) and adjacent mid-Atlantic bight (MAB) waters for use in algorithm development and validation efforts to improve the satellite retrieval of chlorophyll (chl *a*) in Case 2 waters. CB provides a valuable site for validation of data from ocean color sensors for a number of reasons. First, the physical dimensions of the Bay ( $> 6,500 \text{ km}^2$ ) make retrievals from satellites with a spatial resolution of  $\sim 1 \text{ km}$  (*i.e.*, SeaWiFS) or less (*i.e.*, MODIS) reasonable for most of the ecosystem. Second, CB is highly influenced by freshwater flow from major rivers, making it a classic Case 2 water body with significant concentrations of chlorophyll, particulates and chromophoric dissolved organic matter (CDOM) that highly impact the shape of reflectance spectra. Finally, past and ongoing research efforts provided an extensive data set of optical observations that support the goal of this project.

Our SIMBIOS contribution has the following objectives:

- To provide to SeaBASS an on-going stream of *in-situ* optical measurements from CB/MAB for use by the ocean color community in the development and validation of atmospheric correction and ocean bio-optical models;
- To evaluate the performance of SeaWiFS bio-optical algorithms in CB/MAB;
- To evaluate the performance of SeaWiFS  $L_{WN}$  and chl *a* products in CB/MAB on a wide range of time and space scales;
- To utilize our bio-optical database to develop a regionally-tuned algorithm for CB/MAB;
- To produce accurate satellite chl *a* fields that can be used in existing primary productivity models (Harding *et al.*, 2002) to improve the spatial and temporal resolution of primary productivity estimates for CB/MAB.

## RESEARCH ACTIVITIES

### *Field Program*

Throughout our SIMBIOS support we have maintained a seasonal (*i.e.*, spring, summer, fall) sampling schedule in CB proper that started in 1995 with the NSF Land-Margin Ecosystem

Research (LMER) program Trophic Interactions in Estuarine Systems (TIES) cruises (1995-2000) and continued with the NSF Biocomplexity cruises (2001-2003). In addition, five separately funded Office of Naval Research (ONR) cruises were conducted in adjacent offshore waters including coverage to the Gulf Stream. All cruises collected the same suite of measurements. Vertical profiles of downwelling irradiance ( $E_d$ ) and upwelling radiance ( $L_u$ ) were performed with a Biospherical Instruments MER 2040/2041. Discrete samples of particulate and dissolved fractions were collected for determination of absorption spectra of phytoplankton,  $a_{ph}(\lambda)$ , non-pigmented particulates,  $a_d(\lambda)$ , and colored dissolved material,  $a_{CDOM}(\lambda)$ . Discrete particulate samples were collected for pigment analyses of chl *a* by fluorometry and a suite of pigments by HPLC. Radiometric profiles were used to calculate attenuation coefficients for downwelling irradiance ( $K_d$ ) and upwelling radiance ( $K_u$ ), and surface values of  $E_d$  and  $L_u$  ( $E_d\theta$  and  $L_u\theta$ , respectively) by simple linear regression. Instrument self-shading corrections were applied to each  $L_u\theta$  according to Gordon and Ding (1992).

In 2001 we expanded our sampling protocol to include two new radiometric instruments, the Satlantic MicroPro free-fall radiometer and the Satlantic Hyperspectral Tethered Spectral Radiometer Buoy (Hyper-TSRB). The MicroPro is designed for work in more turbid waters with a small sensor cross-section, and is sufficiently compact to allow deployment from small boats. The Hyper-TSRB measures surface  $E_d$  and  $L_u$  with a 120-channel detector from 400 to 800 nm. On each cruise in CB, sequential radiometric profiles were made with the MER (typically one cast) and MicroPro (2 to 5 casts). All casts were completed within ~5 to 8 minutes, depending on the number of MicroPro casts that were made. The TSRB was deployed prior to the radiometric profiles and continuously operated while on station. Since 2001 we also have been using a MicroTops sun photometer from the SIMBIOS instrument pool to measure aerosol optical thickness.

In 2002 and 2003 we participated in six EPA/NASA Atlantic Coast Estuaries Indicators Consortium (ACE INC) cruises that sampled two tributaries of CB, the Patuxent and Choptank Rivers. ACE INC cruises collected the same suite of measurements as the LMER-TIES and Biocomplexity cruises, with the exception that radiometric profiles of  $E_d$  and  $L_u$  were limited to the MicroPro because of vessel limitations.

We have submitted over 3000 *in-situ* measurements to SeaBASS, significantly expanding bio-optical data for Case 2 waters (Table 1). Data from summer and fall cruises of 2003 will be submitted prior to the end of this contract. We also participated in the CHORS/Horn Point Laboratory HPLC Round Robin. We worked together with Jason Perl to identify and minimize methodological and instrument differences between laboratories such that an internally consistent pigment database could be created.

#### *Algorithm Evaluation/Development*

We have evaluated the performance of the empirical algorithm OC4v.4 (O'Reilly *et al.*, 1998) and the semi-analytical Garver-Siegel-Maritorena (GSM01) model (Maritorena *et al.*, 2002) for the CB/MAB, using bio-optical data collected during from SIMBIOS. This entailed using regional bio-optical data as input parameters to GSM01, specifically the chl *a*-normalized absorption,  $a_{ph}^*(\lambda)$ , and spectral slope of the combined absorption of dissolved and non-pigmented particulate matter,  $S_{cdm}$ , to produce a regionally tuned version of the model that we termed GSM01-CB. We subsequently modified the native GSM01 parameters in the SeaWiFS Data Analysis System (SeaDAS) code to evaluate the performance of the GSM01-CB using satellite-derived radiances as input.

### SeaWiFS Image Analysis

We have continued acquisition and analysis of SeaWiFS ocean color imagery from the Goddard Distributed Active Archive Center (DAAC) for CB/MAB. Approximately 100 clear images are available each year from 1998-present. We have used *in-situ* data to validate SeaWiFS performance in match-up analyses and comparisons with continuous underway measurements. A three-year times series (1998-2000) has been used to determine the extent to which SeaWiFS resolves seasonal and interannual variability of chl *a* evident in the *in-situ* data.

## RESEARCH RESULTS

### Radiometric Comparisons

Expansion of our radiometric measurement suite allowed us to evaluate performance of each profiling instrument in nearly simultaneous (*i.e.*, within minutes) deployments. The MicroPro is more suitable for work in Case 2 waters than the MER due to: (1) smaller diameter (MicroPro 4.8 cm relative to the MER 21 cm) that reduces instrument self-shading; (2) higher sampling frequency that gives greater depth resolution in waters where  $K_d$  is high and slight changes in the depth of the sensor have a larger effect on the measurements. The MicroPro exhibited clear advantages in waters with higher  $K_d$ . The improved depth resolution of the MicroPro (~25 measurements per meter) compared to the MER (~10 measurements per meter) compensated for the shallow optical depths in CB where the depth range available for the extrapolation interval was limited to the upper 4 m (or less). Measurements of  $E_d\sigma^+$  from the MicroPro were much easier to reconcile with surface irradiance measurements ( $E_s$ ) than were those from the MER, which exhibited an increasing bias as  $K_d$  increased. Measurements of  $L_u\sigma$  from the MER and MicroPro also showed increased disagreement as  $K_d$  increased. Based on advantages of the MicroPro over the MER in comparisons of  $E_d\sigma^+$  and  $E_s$ , we feel confident that the MicroPro measurements of  $L_u\sigma$  are more accurate than the MER for the Case 2 waters in which we work. Lastly, the increased depth resolution of the MicroPro reduced the effects of small changes in the selection of the depth interval on the calculated  $K_d$ ,  $K_u$ ,  $E_d\sigma$ , and  $L_u\sigma$ . The MicroPro also allowed us to expand our radiometric sampling to more turbid waters of the tributaries, and has improved measurement capabilities in the upper reaches of CB. In addition, these comparisons of the MER and MicroPro indicated strong agreement in the clearer waters of the mesohaline and polyhaline CB, confirming confidence in our multi-year time series of radiometric measurements with the MER in the CB/MAB.

### Characterization of Bio-optical Properties in Chesapeake Bay

Distributions of inherent optical properties in CB/MAB showed the strong influence of freshwater flow that complicates retrievals of chl *a* from satellite ocean color measurements.  $a_{ph}(440)$ ,  $a_{CDOM}(440)$ ,  $a_d(440)$  ranged from 0.1 to 1.0  $m^{-1}$  in CB, and were an order of magnitude lower in the MAB. Whereas  $a_d(440)$  dominated in upper CB, the magnitudes of  $a_{ph}(440)$ ,  $a_{CDOM}(440)$ ,  $a_d(440)$  were comparable in the mid- and lower CB and inshore MAB.  $a_{CDOM}(440)$  dominated water column absorption in the offshore MAB. The distribution  $a_{CDOM}(440)$  was conservative in CB proper, due to its predominant terrestrial origin.  $a_{ph}(440)$  and  $a_d(440)$  were subject to much higher spatial variability. The distribution of dissolved and particulate materials regulated the magnitude and spectral shape of apparent optical properties such as the attenuation coefficient for downwelling irradiance,  $K_d$ , and the remote-sensing reflectance,  $R_{RS}$ .

#### OC4v.4 Algorithm Evaluation

The distribution of inherent optical properties had a pronounced effect on the performance of OC4v.4 in CB/MAB (Fig. 1a). OC4v.4 returned chl *a* estimates for CB that had a small positive bias and high uncertainty, reflecting that  $a_{ph}$ ,  $a_{CDOM}$ ,  $a_d$  did not co-vary. A larger positive bias of chl *a* occurred in the MAB where too much of the total absorption was attributed to phytoplankton. Match-ups of SeaWiFS chl *a* with *in-situ* data showed a larger positive bias in CB than observed for OC4v.4 applied to *in-situ* radiometric data, suggesting SeaWiFS-derived radiances introduced additional errors (Fig. 1b). Whereas the fourth SeaWiFS reprocessing lessened the frequency of negative  $L_{WN}$ , comparisons of *in-situ* and satellite-derived  $L_{WN}$  as part of this work and others have shown that low and negative  $L_{WN}$  from SeaWiFS remain a problem in coastal and estuarine waters (Harding *et al.*, 2003; O'Reilly and Yoder, 2003). SeaWiFS  $L_{WN}$  at 'blue' wavebands are low in CB, causing the return of higher-than-observed chl *a* values by OC4v.4. The lack of increased bias in chl *a* for match-ups from the MAB suggested errors in SeaWiFS chl *a* could be attributed primarily to OC4v.4 rather than errors in atmospheric correction.

#### Resolution of Interannual, Seasonal and Event-scale Variability by SeaWiFS

High uncertainties we detected in match-up analyses of *in-situ* and SeaWiFS chl *a* restricted the usefulness of SeaWiFS data for CB/MAB, stimulating additional analyses to lessen time-space aliasing we believed was responsible. High frequency and small-scale variability of biomass typical of estuaries virtually assures time-space aliasing of satellite observations and *in-situ* measurements. To minimize aliasing, we compared continuous shipboard measurements of chl *a* were to SeaWiFS chl *a* along ship tracks to evaluate the extent to which SeaWiFS resolved variability on small spatial scales. We found SeaWiFS accurately captured small-scale (tens-of-kilometers) variability in the mesohaline and polyhaline regions of CB (Figs. 2 and 3) (see Harding *et al.*, 2003). Increased noise in chl *a* with distance north corresponded to sharp gradients of  $R_{RS}$ . We suspect that the magnitude of the  $R_{RS}$  signal often approaches the limits of sensitivity for the SeaWiFS sensor in turbid waters with high  $K_d$  typical of mesohaline and oligohaline CB, thereby limiting the accuracy of chl *a* retrievals. This gradient of  $R_{RS}$  was persistent in the imagery and was suggested in field radiometric measurements. Comparisons of several transects suggested axial movement of this gradient of  $R_{RS}$  could determine the areal extent of accurate chl *a* retrievals.

Our evaluation of the 1998 – 2000 SeaWiFS chl *a* time series demonstrated that SeaWiFS accurately represented seasonal and interannual variability of chl *a* revealed by *in-situ* data (Fig. 4). Previous research has shown that the timing, position, and magnitude of the spring bloom are regulated by freshwater flow from the Susquehanna River, supplying nutrients and suspended matter to CB (Malone *et al.*, 1988; Harding, 1994). We found that positive deviations of flow in 1998 resulted in a large spring bloom that extended to the polyhaline CB, shown in both satellite imagery and *in-situ* data. Measurements of chl *a* in 1999, a year of lower-than-average freshwater flow, showed a smaller bloom that was confined primarily to the mesohaline CB. Lastly, average freshwater flow in 2000 supported a bloom of intermediate magnitude that extended throughout the mesohaline and part of the polyhaline CB. As observed in transect data, the agreement of SeaWiFS and *in-situ* chl *a* improved with distance down the Bay. These comparisons with *in-situ* data confirm that SeaWiFS chl *a* retrievals are reasonable and can significantly augment the temporal resolution of shipboard measurements.

The benefit of increased temporal resolution afforded by satellite observations was demonstrated in the three-year time series of chl *a* (Fig. 5). Mean, daily chl *a* for a section of polyhaline CB indicated several doublings over a period of days in summer that were not captured by *in-situ* observations. We hypothesize that these doublings could be associated with short-lived,

but dense blooms of dinoflagellates that have been observed before during stratified conditions in summer. The sedimentation of organic matter from the spring bloom fuels at least 50% of the primary productivity in surface waters (Kemp and Boynton, 1984). Transient destratification of the summer pycnocline provides the nutrients to support these short-lived summer blooms (Malone *et al.*, 1986, 1988). The higher frequency of these events in 1998 and 2000, contrasted with 1999, may indicate a connection between these transient blooms in summer and the magnitude of the spring bloom. Further study is needed to test this hypothesis.

### Regional Parameterization of GSM01

Semi-analytical models such as GSM01 have the potential to distinguish multiple constituents of the water column that absorb and scatter light. This attribute is crucial for accurate retrievals of chl *a* in optically-complex waters. The original parameterization of GSM01 was based on *in-situ* observations from Case 1 waters and optimized for performance in the global ocean (Maritorena *et al.*, 2002). We hypothesized that regional parameterization was necessary to improve the accuracy of chl *a* retrievals in CB/MAB.

We evaluated our extensive data set of *in-situ* observations to derive regionally and seasonally (when appropriate) values for  $a_{ph}^*(\lambda)$  and  $S_{cdm}$  to produce a regional version of the model, GSM01-CB (Table 2) (see Magnuson *et al.*, 2003). Regional differences of  $a_{ph}^*(\lambda)$  were significant across the study area from the upper, oligohaline CB to the offshore MAB.  $a_{ph}^*(\lambda)$  increased with distance offshore. This trend of increasing  $a_{ph}^*(\lambda)$  can be explained by two factors: decreases of pigment packaging and increases in concentrations of accessory pigments (Yentsch and Phinney, 1989; Bricaud *et al.*, 1995). We also found significant seasonal differences in mean  $a_{ph}^*(\lambda)$  in CB that we attributed primarily to changes in floral composition and associated changes in pigment packaging. Highest  $a_{ph}^*(\lambda)$  occurred in spring when phytoplankton in CB consist primarily of large cells (*i.e.*, centric diatoms) (Malone *et al.*, 1988; Marshall and Nesiuis, 1996). Lowest  $a_{ph}^*(\lambda)$  occurred in summer when phytoplankton consist of a more diverse flora of smaller, flagellated cells (Malone *et al.*, 1991; Marshall and Nesiuis, 1996).  $S_{cdm}$  exhibited an increasing trend with distance offshore, reflecting the decrease in the influence of terrigenous CDOM (Blough *et al.*, 1993; Nelson and Guarda, 1995; Vodecek *et al.*, 1997). Small, but significant differences in  $S_{cdm}$  along the axis of CB reflected changes in the relative contributions of  $a_{CDOM}$  and  $a_d$  to  $a_{cdm}$ .

Evaluation of chl *a* from OC4v.4, GSM01, and GSM01-CB for CB, inshore MAB, and offshore MAB showed that GSM01-CB improved upon OC4v.4 in all regions (Table 3). The local parameterization was essential to improve performance in CB. Both GSM01 and GSM01-CB showed improvement over OC4v.4 in the MAB where they specifically accounted for high absorption due to CDOM. Analysis of SeaWiFS products using each model supported results of evaluations using *in-situ*  $R_{RS}$  as inputs, specifically that: (1) one or both of the semi-analytical models performed as well or better than OC4v.4; (2) both semi-analytical models clearly improved upon OC4v.4 in the offshore MAB (Fig. 6). Disagreement between the *in-situ* observations and model estimates for the inshore MAB likely reflect a sampling bias as *in-situ* observations were concentrated near the Chesapeake and Delaware Bay mouths where chl *a* are generally high, whereas mean chl *a* from SeaWiFS represented the entire inshore region. Disregarding *in-situ* values, both semi-analytical models returned much lower mean chl *a* for the inshore MAB than did OC4v.4, and GSM01-CB chl *a* were lower than GSM01 chl *a*, consistent with results obtained using *in-situ* data as input to the models (Table 3). The semi-analytical models return estimates of  $a_{cdm}(443)$  and  $b_{bp}(443)$  in addition to chl *a*. Estimates of  $a_{cdm}(443)$  were closer to *in-situ* values than were estimates of chl *a* (Table 3). Meaningful evaluation of the  $b_{bp}(443)$  product was not possible due to the lack of direct measurements of scattering. Relatively high uncertainties remained for all models, reflecting the difficulty of working in bio-optically complex waters. It is notable that the

performance of GSM01 was surprisingly good, considering the parameterizations differed significantly from bio-optical properties of CB. The relatively strong performance of GSM01 in CB likely reflects the robust tuning procedures used by Maritorena *et al.* (2002) to derive global parameters for GSM01.

## CONCLUSIONS

It is useful and necessary to assess the performance of satellite products at various stages to recognize limitations, as well as to demonstrate strengths and identify the types of applications for which current products are suitable. We have made use of an extensive database of bio-optical measurements to provide several different evaluations of SeaWiFS performance. We have identified several factors that inhibit the accuracy of the standard chl *a* product, including: (1) use of an empirical algorithm that does not account for all of the absorbing and scattering components of the water column; (2) atmospheric correction errors in coastal waters resulting in low satellite radiances; (3)  $R_{RS}$  signals approaching the lower sensitivity limits of the sensor. Nonetheless, the importance of these factors lessened, and SeaWiFS recoveries improved, with increasing water clarity. We found that in mesohaline and polyhaline CB, SeaWiFS accurately and reliably represented seasonal and interannual variability of biomass, resolved event-scale phenomena that were missed in our field sampling, and resolved small-scale (*i.e.*, tens-of-kilometers) regional variability. We are encouraged by our initial attempts to improve chl *a* estimates by tuning a semi-analytical model for regional conditions. We are exploring modifications to the parameterization that could lessen the use of abrupt transitions in parameter values at seasonal and regional boundaries. Improvements in SeaWiFS radiances also will be necessary for maximal performance of semi-analytical models in Case 2 waters.

## FIGURE LEGENDS

Figure 1. Comparisons of *in-situ* chl *a* to: (a) mean chl *a* estimated from OC4v.4 applied *in-situ*  $R_{RS}$ ; (b) SeaWiFS retrievals of chl *a*. SeaWiFS values represent the mean of a 3x3 pixel box centered on the station location for *in-situ* sampling. A chl *a* match-up was considered valid if the satellite observation and *in-situ* measurement occurred on the same day, and half (5 out of 9) pixels in a box centered on the latitude and longitude of the sampling station returned valid estimates. The original match-up dataset contained over 1400 chl *a* observations. Mean ratio = SeaWiFS chl *a* / *in-situ* chl *a*.

Figure 2.  $R_{RS555}$  and chl *a* from SeaWiFS for a CB scene, 15 October 1999. The white lines superimposed on these images give the ship track for concurrent measurements that included bio-optical measurements. Vertical line at ~ 150 km marks the time of the satellite pass.

Figure 3. (a) Comparison of *in-situ* and SeaWiFS chl *a*; (b) SeaWiFS  $R_{RS510}$  and  $R_{RS555}$ ; (c) ratio of SeaWiFS  $R_{RS510/555}$  along a ship track in the mesohaline to polyhaline CB.

Figure 4. Monthly mean chl *a* for the polyhaline (I, II, south of 37.8°N) and mesohaline (III, IV, 37.8 – 38.8°N) regions of CB from *in situ* (closed symbols) and SeaWiFS (open symbols) data for 1998-2000.

Figure 5. Time-series of mean, daily SeaWiFS (open symbols) and *in-situ* (closed symbols) chl *a* for a section of CB from 37.3° to 37.8°N.

Figure 6. Histograms of *in-situ* and SeaWiFS chl *a* observations for CB, inshore and offshore MAB for image from 4 May 2000: (a-c) *in-situ*, (d-f) SeaWiFS OC4v.4, (g-i) SeaWiFS GSM01, and

(j-l) SeaWiFS GSM01-CB. *In-situ* data from the MAB were not available for this image. The inshore and offshore MAB histograms (b-c) reflect all field data available for those regions during May-September 1996-1998.

## REFERENCES

- Blough, N.V., O.C. Zafirou, and J. Bonilla, 1993: Optical absorption spectra of waters from the Orinoco River outflow: terrestrial input of colored organic matter to the Caribbean. *Journal of Geophysical Research* **98**, 2,271-2,278.
- Bricaud, A., M. Babin, A. Morel, & H. Claustre, 1995: Variability in the chlorophyll-specific absorption coefficients of natural phytoplankton: analysis and parameterization, *Journal of Geophysical Research* **100**, 13,321-13,332.
- Gordon, H.R. and K. Ding. 1992. Self-shading of in-water optical instruments. *Limnology and Oceanography*, **37**, 491-500.
- Kemp, W.M., & W.R. Boynton, 1984: Spatial and temporal coupling of nutrient inputs to estuarine primary production: the role of particulate transport and decomposition, *Bulletin of Marine Science* **35**, 242-247.
- Harding, L.W., Jr., E.C. Itsweire, and W.E. Esaias, 1994: Estimates of phytoplankton biomass in the Chesapeake Bay from aircraft remote sensing of *chl-a*, 1989-1992, *Remote Sensing of Environment* **49**, 41-56.
- Harding, L.W., Jr., A. Magnuson, and M.E. Mallonee, 2003: Bio-Optical and Remote Sensing Observations in Chesapeake Bay, *Estuarine, Coastal, and Shelf Science* (in review).
- Magnuson, A., L.W. Harding, Jr., M.E. Mallonee, and J.E. Adolf, 2003: Bio-optical modeling in the Middle Atlantic Bight, *Estuarine, Coastal, and Shelf Science* (to be submitted).
- Malone, T.C., W.M. Kemp, H.W. Ducklow, W.R. Boynton, J.H. Tuttle, and R.B. Jonas, 1986: Lateral variation in the production and fate of phytoplankton in a partially stratified estuary, *Marine Ecology Progress Series* **32**, 149-160.
- Malone, T.C., L.H. Crocker, S.E. Pike, and B.W. Wendler, 1988: Influences of river flow on the dynamics of phytoplankton production in a partially stratified estuary, *Marine Ecology Progress Series* **48**, 235-249.
- Malone, T.C., H.W. Ducklow, E.R. Peele, and S.E. Pike, 1991: Picoplankton carbon flux in Chesapeake Bay, *Marine Ecology Progress Series* **78**, 11-22.
- Maritorena, S., D.A. Siegel, and A.R. Peterson, 2002: Optimization of a semianalytical ocean color model for global-scale applications, *Applied Optics* **41**, 2,705-2,714.
- Marshall, H.G. and K.K. Nesius, 1996: Phytoplankton composition in relation to primary production in Chesapeake Bay, *Marine Biology* **125**, 611-617.
- Nelson, J.R. and S. Guarda, S. 1995: Particulate and dissolved spectral absorption on the continental shelf of the southeastern United States, *Journal of Geophysical Research* **100**, 8,715-8,732.

O'Reilly, J.E., S. Maritorena, B.G. Mitchell, D.A. Siegel, K.L. Carder, S.A. Garver, M.Kahru, & C.R. McClain, 1998: Ocean color algorithms for SeaWiFS, *Journal of Geophysical Research* **103**, 24,937-24,953.

O'Reilly, J.E. and J.A. Yoder, 2003: A comparison of SeaWiFS LAC Products from the Third and Fourth Reprocessing: Northeast US Ecosystem, in *Algorithm Updates for the Fourth SeaWiFS Data Reprocessing* (S.B. Hooker & E.R. Firestone, eds). NASA Tech. Memo. 2003-206892, NASA Goddard Space Flight Center, Greenbelt, MD, pp. 60-67.

Vodacek, A., N.V. Blough, M.D. DeGrandpre, E.T. Peltzer, and R.K. Nelson, 1997: Seasonal variation of CDOM and DOC in the Middle Atlantic Bight: Terrestrial inputs and photooxidation, *Limnology and Oceanography* **42**, 674-686.

Yentsch, C.S. and D.A. Phinney, 1989: A bridge between ocean optics and microbial ecology, *Limnology and Oceanography* **34**, 1,694-1,705.

#### PEER REVIEWED PUBLICATIONS

Harding, L.W., W. . Miller, R.N. Swift, and C.W. Wright, 2001: Aircraft remote sensing, In: Steele, J., S. Thorpe, and K. Turekian, (eds.), *Encyclopedia of Ocean Sciences*, Academic Press, London, UK, 113-122.

Harding, L.W., Jr., M.E. Mallonee, and E.S. Perry, 2002: Toward a predictive understanding of primary productivity in a temperate, partially stratified estuary, *Estuarine, Coastal and Shelf Science* **55**, 437-463.

Harding, L.W., Jr., A. Magnuson, and M.E. Mallonee, 2003: Bio-Optical and remote sensing observations in Chesapeake Bay, *Estuarine, Coastal, and Shelf Science* (in review).

Magnuson, A., L.W. Harding, Jr., M.E. Mallonee, and J.E. Adolf, 2003: Bio-optical modeling in the middle Atlantic bight, *Estuarine, Coastal, and Shelf Science* (to be submitted).



#### PRESENTATIONS

Harding, L.W., Jr., Seasonal and inter-annual variability of primary productivity in Chesapeake Bay from remotely sensed aircraft observations, at Phytoplankton Productivity - An Appreciation of 50 Years of the Study of Production in Oceans and Lakes, 19 March 2002, Bangor, Wales, United Kingdom.


Magnuson, A., L.W. Harding, Jr., and M.E. Mallonee, Progress toward a case II bio-optical algorithm optimized for Chesapeake Bay and the adjacent waters of the mid-Atlantic bight. Ocean Optics XVI, 18-22 November 2002, Sante Fe, NM.

Magnuson, A., L.W. Harding, Jr., M.E. Mallonee, and W.D. Miller, Comparison of *In Situ* radiometric measurements in turbid waters, poster presented at the 5th SIMBIOS Science Team Meeting, 15-17 January 2002, Baltimore, MD.







Harding and Magnuson, Figure 1




Harding and Magnuson, Figure 2




Harding and Magnuson, Figure 3



Harding and Magnuson, Figure 4



Harding and Magnuson, Figure 5



Harding and Magnuson, Figure 6

Table 1. Status of the Chesapeake Bay data set submitted to SeaBASS. Numbers for 2003 are estimated based on cruises yet to be completed.

Direct Measurements	Derived Products	1996	1997	1998	1999	2000	2001	2002	2003
<b>MER*</b>									
$E_d(z, \lambda)$	$R_{RS}(0^+, \lambda)$	82	93	83	44	40	25	39	25
$L_u(z, \lambda)$	$K_d(m, \lambda)$								
$E_s(0^+, \lambda)$	$K_u(m, \lambda)$								
<b>MicroPro**</b>									
$E_d(z, \lambda)$	$R_{RS}(0^+, \lambda)$	---	---	---	---	---	49	130	90
$L_u(z, \lambda)$	$K_d(m, \lambda)$								
$E_s(0^+, \lambda)$	$K_u(m, \lambda)$								
$a_p(0, \lambda), a_d(\lambda)$	$a_{ph}(\lambda)$	79	32	130	66	46	27	32	25
$a_{CDOM}(0, \lambda)$		79	32	130	66	46	27	32	25
Chl <i>a</i> (z)		323	709	591	514	330	189	182	60
Pigments (z)		143	95	104	63	56	49	32	20
$T(z), S(z)$		---	30	55	49	43	26	31	20
MicroTops		---	---	---	---	---	---	24	15

\* MER wavebands 412, 443, 455, 490, 510, 532, 550, 650, 589, 625, 671, 683, 700

\*\*MicroPro wavebands 400, 412, 443, 455, 490, 510, 532, 555, 565, 590, 625, 670, 684, 700.

Direct Measurements	
$E_d(z, \lambda)$	Downwelling irradiance depth profiles
$L_u(z, \lambda)$	Upwelling radiance depth profiles
$E_s(0^+, \lambda)$	Incident irradiance
$a_p(z, \lambda)$	Total particulate absorption (300-900 nm) of discrete samples by QFT
$a_d(z, \lambda)$	Non-pigmented particulate absorption (300-900 nm) of discrete samples by QFT with MeOH extraction
$a_{cdom}(z, \lambda)$	Colored dissolved absorption (190-820 nm) of discrete samples
Chl <i>a</i> (z)	Determination of chl <i>a</i> for discrete samples by Turner fluorometry
pigments (z)	HPLC analysis for a suite of pigments
$T(z), S(z)$	SeaBird temperature and conductivity profiles
Derived Products	
$R_{RS}(0^+, \lambda)$	Remote sensing reflectance from calculated from radiance/irradiance profiles
$K_d(m, \lambda), K_u(m, \lambda)$	Attenuation coefficients for $E_d(z, \lambda)$ and $L_u(z, \lambda)$ for the surface mixed layer, calculated from radiance/irradiance profiles
$a_{ph}(z, \lambda)$	Pigmented particulate absorption (300-900 nm), calculated as $a_p(0, \lambda) - a_d(0, \lambda)$

Table 2. Parameter values used in GSM01 and GSM01-CB.

Region	Season	$a_{ph}^*(412)$	$a_{ph}^*(443)$	$a_{ph}^*(490)$	$a_{ph}^*(510)$	$a_{ph}^*(555)$	$a_{ph}^*(671)$	$S_{cdm}$	$\eta$
GSM01		0.00665	0.05582	0.02055	0.01910	0.01015	0.01424	0.02061	1.03373
GSM01-CB									
Upper Bay	Spring	0.02119	0.02509	0.01282	0.00910	0.00427	0.02087	0.01218	0
	Summer	0.02653	0.02979	0.01655	0.01208	0.00470	0.02122	0.01218	0
	Fall	0.02259	0.02573	0.01372	0.01019	0.00376	0.02066	0.01218	0
Mid-Bay	Spring	0.02001	0.02212	0.01279	0.00974	0.00449	0.01588	0.01385	0
	Summer	0.03345	0.03900	0.02318	0.01664	0.00609	0.02285	0.01385	0
	Fall	0.02758	0.03080	0.01826	0.01438	0.00691	0.02112	0.01385	0
Lower Bay	Spring	0.02001	0.02212	0.01279	0.00974	0.00449	0.01588	0.01330	0
	Summer	0.03345	0.03900	0.02318	0.01664	0.00609	0.02285	0.01330	0
	Fall	0.02758	0.03080	0.01826	0.01438	0.00691	0.02112	0.01330	0
Inshore		0.07123	0.08843	0.06024	0.04072	0.01693	0.03815	0.01236	0
Offshore		0.11331	0.14678	0.09832	0.06048	0.01920	0.04349	0.01646	1



Table 3. Comparison of the mean ratio ( $\pm$ SD) of modeled and measured chl *a* and  $a_{cdm}(443)$ , and estimated  $b_{bp}(443)$  for Chesapeake Bay, inshore MAB, and offshore MAB. Mean ratio = modeled/measured.

	Chesapeake Bay chl <i>a</i>	Inshore MAB chl <i>a</i>	Offshore MAB chl <i>a</i>	Chesapeake Bay $a_{cdm}(443)$	Inshore MAB $a_{cdm}(443)$	Offshore MAB $a_{cdm}(443)$	Chesapeake Bay $b_{bp}(443)$	Inshore MAB $b_{bp}(443)$	Offshore MAB $b_{bp}(443)$
GSM01	1.99 $\pm$ 2.21	1.84 $\pm$ 0.92	1.31 $\pm$ 0.68	1.24 $\pm$ 0.58	1.01 $\pm$ 0.54	0.87 $\pm$ 0.61	0.94 $\pm$ 0.65	2.28 $\pm$ 1.35	1.43 $\pm$ 0.88
GSM01- CBAY	1.10 $\pm$ 1.08	0.59 $\pm$ 0.57	0.51 $\pm$ 0.29	0.92 $\pm$ 0.40	1.12 $\pm$ 0.42	0.95 $\pm$ 0.67	1.35 $\pm$ 0.91	2.88 $\pm$ 1.64	1.34 $\pm$ 1.26
OC4v.4	1.20 $\pm$ 0.71	3.12 $\pm$ 2.05	2.40 $\pm$ 1.03	NA	NA	NA	NA	NA	NA

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> 10-14-2003	<b>3. REPORT TYPE AND DATES COVERED</b> Final 12-1-2001 - 11-30-2003	
<b>4. TITLE AND SUBTITLE</b> Evaluation and Validation of Case 2 Algorithms in Chesapeake Bay			<b>5. FUNDING NUMBERS</b> NAS5-00195	
<b>6. AUTHOR(S)</b> Harding, Lawrence W., Jr. Magnuson, Adrea				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> University of Maryland Center for Environmental Science (Horn Point Laboratory) PO Box 775 Cambridge, Maryland 21613			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> NASA/Goddard Space Flight Center Greenbelt Road, Code 210.5 Greenbelt, Maryland 20771			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Unclassified Unlimited			<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 Words)</b> The SIMBOIS Project "evaluation and validation of Case 2 algorithms in Chesapeake Bay" focused on obtaining regional in situ observation of bio-optical properties in support of the SeaWiFS and SIMBIOS programs. Our study sites included Chesapeake Bay and the middle Atlantic bight where optical complexity imposed by high concentrations of photosynthetic pigments, chromophoric dissolved organic matter, and suspended particulate material makes retrievals of chlorophyll (chl-a) problematic. We have provided data from several field programs to the SeaBASS data archive at NASA, participated in algorithm development in SIMBIOS, generated time-series data from satellite and aircraft instruments (SeaWiFS, SAS III) and developed models of primary productivity and chl-a for application to remotely sensed data.				
<b>14. SUBJECT TERMS</b> SEAWiFS, SIMBIOS, ocean optics, ocean color sensor validation			<b>15. NUMBER OF PAGES</b> 18	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> unclassified	<b>20. LIMITATION OF ABSTRACT</b> Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)

Prescribed by ANSI Std. Z39-18

298-102