

28th Annual International Gravitational Physiology Meeting
8-13 April, 2007
Menger Hotel, San Antonio, TX, USA
Abstract Form

Email Abstract (in Word or PDF) to:
bjcoon@ucdavis.edu

Your entire abstract, including Title, Author names and Institution must fit entirely in the outlined box below (7" x 4.5" or 17.8 cm x 11.4 cm)

Preferred Presentation Method:

- Oral Poster

Audio/Visual Requirements:

- Slide Projector
 Over-Head Projector
 LCD Projector

Postal address of the corresponding author:

Andre' Diedrich, M.D., Ph.D.
 Autonomic Dysfunction Center, Vanderbilt
 University
 1161 21st Avenue South, Suite AA3228
 MCN
 Nashville, TN 37232-2195

Corresponding author Email address,
 Phone & Fax:

ph (615) 343-6499
 fax (615) 343-8649
 e-mail andre.diedrich@vanderbilt.edu

**EFFECTS OF DAILY CENTRIFUGATION ON SEGMENTAL
 FLUID DISTRIBUTION IN BED-RESTED SUBJECTS**


A DIEDRICH¹, ST MOORE², M STENGER³, T. M ARYA³, N

NEWBY³, JM TUCKER³, L MILSTEAD³, K ACOCK³, C

KNAPP⁴, J EVANS⁴, W PALOSKI³

¹Vanderbilt University Medical School, Nashville, TN, ²Mount Sinai
 Hospital, New York, ³Johnson Space Center UTMB Galveston T.,
⁴Biomedical Engineering, University of Kentucky

The effect of daily centrifugation on segmental fluid distribution have been studied during 21 days of 6° head down bedrest. One group (N=7) underwent no countermeasure while the other (N=8) received a daily, one hour, dose (2.5 gz at the foot, decreasing to 1.0 gz at the heart) of artificial gravity (AG) training on the Johnson Space Center short radius centrifuge. Fluid shifts of thoracic (VTO), abdominal (VAB), thigh (VTH), and calf (VCA) regions were measured by the tetrapolar segmental body impedance technique. *Results:* Untrained subjects reduced their total volume from 18.9±0.5L to 17.9±0.9L (MN±SE, P<0.05) while trained subjects maintained their total volume. In untrained, control, subjects after bed rest, there was a trend toward reduced volume in all segments, with significant reductions in thigh and calf (fig, P<0.05). Trained subjects maintained volume in all segments. Our data indicate that artificial gravity treatment counteracts bed rest-induced hypovolemia. *Supported by the NASA AG-Bed Rest Pilot Project. We thank the Cardio and Centrifuge Teams for their support.*



TO BE RECEIVED NO LATER THAN 1 FEBRUARY, 2007