

Microbial extremophiles in aspect of limits of life.

Elena V. Pikuta¹, Richard B. Hoover², and Jane Tang³

^{1,2}-National Space Sciences and Technology Center/NASA, VP-62, 320 Sparkman Dr., Astrobiology Laboratory, Huntsville, AL 35805, USA.

³- Noblis, 3150 Fairview Park Drive South, Falls Church, VA 22042, USA.

Introduction.

During Earth's evolution accompanied by geophysical and climatic changes a number of ecosystems have been formed. These ecosystems differ by the broad variety of physicochemical and biological factors composing our environment. Traditionally, pH and salinity are considered as geochemical extremes, as opposed to the temperature, pressure and radiation that are referred to as physical extremes (Van den Burg, 2003). Life inhabits all possible places on Earth interacting with the environment and within itself (cross species relations). In nature it is very rare when an ecotope is inhabited by a single species. As a rule, most ecosystems contain the functionally related and evolutionarily adjusted communities (consortia and populations). In contrast to the multicellular structure of eukaryotes (tissues, organs, systems of organs, whole organism), the highest organized form of prokaryotic life in nature is the benthic colonization in biofilms and microbial mats. In these complex structures all microbial cells of different species are distributed in space and time according to their functions and to physicochemical gradients that allow more effective system support, self-protection, and energy distribution. *In vitro*, of course, the most primitive organized structure for bacterial and archaeal cultures is the colony, the size, shape, color, consistency, and other characteristics of which could vary specifics on species or subspecies levels. In table 1 all known types of microbial communities are shown (Pikuta et al., 2005).

In deep underground (lithospheric) and deep-sea ecosystems an additional factor – pressure, and irradiation - could also be included in the list of microbial communities. Currently the best-studied ecosystems are: human body (due to the medical importance), and fresh water and marine ecosystems (due to the reason of an environmental safety). For a long time, extremophiles were *terra incognita*, since the environments with aggressive parameters (compared to the human body temperature, pH, mineralization, and pressure) were considered *a priori* as a dead zone. It took time to find out that the environments with extreme physico-chemical and climatic parameters are inhabited by a wide spectrum of different microorganisms. In chronological consequences the discoveries of extremophiles are presented as follows:

Slight acidophilic property (pH 4-6) was known in many fungi a long time ago, but the first obligately acidophilic bacterium described was *Acidithiobacillus ferrooxydans* (formally *Thiobacillus ferrooxidans*), later thermophilic lithotrophic acidophiles were found, and the hyper-acidophilic species of the genus *Picrophilus* (capable of grow at negative pH values) were described in 1996 (Schleper et al., 1996).

The epoch of study of thermophilic microorganisms starts with the discovery of *Thermus aquaticus* (Brock and Freeze, 1969), and now the maximum temperature for growth at 113 °C was found for *Pyrolobus fumarii* (Blöchl et al., 1997). Another hyperthermophilic microorganism, strain 121(not validly published) that was isolated by Kashefi and Lovley (2003) survives for short periods of time at 130 °C (Cowan, 2004).

Table 1. Known types of microbial communities.

Types of communities:	pH	NaCl, %(w/v)	Temperature, °C
1. Freshwater psychrophilic 2. Freshwater, meso-thermal 3. Freshwater moderately thermophilic 4. Freshwater thermophilic	5-7	0-1	< 10 15-40 50-60 70-110
5. Marine psychrophilic 6. Marine, meso-thermal 7. Marine moderately thermophilic 8. Marine thermophilic	8	3-4	<10 15-40 50-60 70-120
9. Alkaliphilic psychrophilic 10. Alkaliphilic mesophilic 11. Alkaliphilic moderately thermophilic 12. Alkaliphilic thermophilic	9-11	0-1	<10 15-40 50-60 70-110
13. Haloalkaliphilic psychrophilic 14. Haloalkaliphilic mesophilic 15. Haloalkaliphilic moderately thermophi	9-10	3-25	<10 15-40 50-60
16. Halophilic psychrophilic 17. Halophilic mesophilic 18. Halophilic moderately thermophilic	8.0	3-30	<10 15-40 50-60
19. Acidophilic mesophilic* 20. Acidophilic moderately thermophilic 21. Acidophilic thermophilic	0-4	0-2	15-40 50-60 70-120

The first mention of the term “psychrophile” was made by Schmidt-Nielsen in 1902 for the description of bacteria capable of growth at 0 °C (Morita, 1975), but arctic diatoms already were studied more than one hundred years ago without using this term (Van Heurck, 1909). Now with the development of a special technique of cooled instruments and well equipped cold rooms it has become possible to study truly psychrophilic microorganisms. Firstly in our Astrobiology Laboratory at NASA/MSFC/NSSTC the bacterial growth (in pure culture) was determined at – 5 °C on a liquid media (Hoover et al., 2002; Pikuta and Hoover, 2003; Pikuta et al., 2003b), and our Russian colleges successfully cultivated aerobic bacterial cultures on solid agar media also at – 5 °C (Gilichinsky et al., 2005). The observations of living microorganisms *in situ* at – 20 °C in highly mineralized media were reported previously (Staley and Gosink, 1999).

* One exception – *Thiobacillus halophilus* – it has optimal NaCl concentration at 5-6 % (w/v) and maximum concentration at 24 % (w/v).

The first mentioned alkaliphile was the bacterium *Streptococcus fecalis* (Downie and Cruickshank, 1928), but several years earlier an article about alkalitolerant nitrifying bacteria was published (Meek and Lipman, 1922). Extreme alkaliphiles belonging to genera *Clostridium* and *Bacillus* were isolated from soils (Horikoshi and Akiba, 1982), but truly alkaliphilic microorganisms belonging to separate genera such as *Natronobacterium* and *Natronococcus* were described much later (Tindall, et al., 1984).

Study of halophilic microorganisms has started from work with saline soils and lakes, and now the record of good growth for *Haloferax mediterranei* has been demonstrated at 30 % NaCl.

Study of barophiles became possible after great engineering of submersible crafts and achievements in Black Smokers study. It was shown that there are microorganisms that require high pressure in addition to the high temperature, and the highest known limit of life was detected at 100 MPa (Yayanos *et al.*, 1979).

The first radioresistant bacterium *Deinococcus radiodurans* was found during the process of food conservation and storage. This bacterium could survive ionizing irradiation and other DNA-damaging assaults at doses that are lethal to all other organisms (Raj *et al.*, 1960). Among archaea the hyperthermophilic sulfur-reducing *Thermococcus gammatolerans* that is capable of resisting 30 kGy of γ -irradiation was described comparatively recently (Edmond *et al.*, 2003).

Anaerobiosis, as an alternative to the aerobic life, was introduced after Pasteur's discovery of fermentation (Pasteur, 1861), but an anaerobic technique for the cultivation of obligately anaerobic microorganisms was developed much later (Hungate, 1969). Among the bacteria and archaea there are many anaerobic species, and most of them are not extremophiles. For this reason an anaerobiosis is not considered as imperative to extremophilic life. However, it should be remembered that the first life forms on early Earth were anaerobic extremophiles, and therefore, this capacity is a very important issue for the logical discussion about the limits of life.




From the systematics point of view life in extreme ecosystems is represented by both eukaryotic and prokaryotic organisms, and the comparison of all taxa in complex biocoenosis of each ecosystem will provide answers to the fundamental questions of the origin, distribution, and evolution of life (direction of changes in the phenotype).

1. Acidophilic microorganisms.


It is well known that proteins denature at low pH, and from this point of view extreme acidophilic organisms represent great potential for biotechnology and significance for the molecular and genetic research. In nature fish and cyanobacteria have not been found at pH below 4, plants and insects below pH 2-3 (Rothschild & Mancinelli, 2001). It is exceptionally interesting that the first representative of extreme acidophiles growing near pH 0 were found to be not prokaryotic but eukaryotic microorganisms. *Cyanidium caldarium* is a red algae belonging to rhodophytes. It contains chlorophyll "a" and C-phycocyanin in its chloroplasts, and it colonizes hot, acid "soils" and waters all over the world except for Hawaii, which are geologically relatively young islands. Three other extreme acidophilic eukaryota are fungi: *Aconitium cylatium*, *Cephalosporium* sp., and *Trichosporon cerebriae* (Schleper *et al.*, 1995). The green alga *Dunaliella acidophila* can survive pH 0, but maximum of growth for this organism occurs at pH 1 (Pick, 1999), but discovered archaea *Picrophilus oshimae* and *P. torridus* broke all records and demonstrated that life could exist at a minimum pH of -0.2. The cells of these archaea grow within a moderately thermal regime and were

found near a hydrothermal spring with solfataric gases in Japan (Schleper et al., 1996). Another extremely acidophilic microorganism *Ferroplasma acidarmanus* is eubacterium growing at pH 0 in acid mine drainage in Iron Mountain, California. This species does not have a cell wall, and the cell membrane is the only barrier between the cytoplasm and concentrated sulfuric acid with high concentrations of copper, arsenic, cadmium and zinc in the surrounding medium.

Bacteria growing at pH less than 3.0 could have chemolithotrophic, chemolithomixotrophic, or chemoorganoheterotrophic metabolism. Among acidophiles there are hyperthermophilic, moderately thermophilic and mesophilic species. They could be spore-forming and non spore-forming, aerobic, micro-aerophilic, or obligately anaerobic. Cell membrane could have a positive or negative reaction to the gram-stain, and sometimes the cell wall could be reduced to a single membrane. They could receive energy from hydrogen, iron, sulfur, or organic molecules. Usually participating in the process of transformation of sulfur and iron from minerals and rocks (bioleaching), acidophiles form acidic media (equilibria 1-3):



But they are also capable of the consumption of a proton that can lead to alkalinity in some mine waters (equilibrium 4):



Some acidophilic microorganisms have the ability to maintain a neutral pH inside of cytoplasm, and intracellular enzymes from these organisms do not need to be adapted to extremely low pH. The internal pH of acidophiles has been measured at between 5 and 7 (Norris and Johnson, 1998). However the extra cellular proteins usually have adaptation to high acidity and their optimum of function occur at low pH.

We have already discussed how these microorganisms could have developed an adaptation to the several extremes at the same time. High temperature, low pH, and high concentration of certain metals (the term *metalophile* was used in some literature, but has not received wide acceptance) are examples of multiple effects of environment. In acidic environments the solubility of metals is much greater, and in many cases can reach the g l^{-1} concentrations. In cells of acidophiles the metals may accumulate above the normal physiological concentrations by the action of unspecific, constitutively expressed transport systems, whereby they become toxic. Intracellular metals can exert a toxic effect by forming coordinate bonds with anions blocking functional groups of enzymes, inhibiting transport systems, displacing essential metals from their native binding sites and disrupting cellular membrane integrity. There are five basic mechanisms (Dopson *et al.*, 2003) that convey an increased level of cellular resistance to metals: (1) efflux of the toxic metal out of the cell; (2) enzymatic conversion; (3) intra- or extracellular sequestration; (4) exclusion by a permeability barrier; and (5) reduction in sensitivity of cellular targets. Metal resistance mechanisms of acidophiles were extensively studied for arsenic, copper, zinc, cadmium, nickel, mercury, silver, ferric iron, molybdenum, chromium, and uranium. The examples of acidophilic bacteria and archaea resistant to the toxic concentrations of As(III) are *Acidithiobacillus caldus* KU and *Sulfolobus metallicus* BC respectively. *At. ferrooxidans* strains adapted to high levels of Cu(II) could be tolerant to 800 mM (Dew *et al.*, 1999). The toxicity of Zn(II) to *At. ferrooxidans* depends upon the growth substrate. One strain resistant to 153 mM (10 g l^{-1}) Zn(II) while growing on Fe(II) is sensitive to 92 μM (0.6 mg l^{-1}) when growing on thiosulfate (Trevors *et al.*, 1985). Acidophiles resistant to Cd(II) include a

number of *Acidiphilum* spp. And an *Acidocella* strain resistant to 700 mM Cd(II), while the maximum resistance in archaea was 10 mM in *Sulfolobus acidocaldarius* and *S. solfataricus* (Dopson *et al.*, 2003). Adapted strains of *At. ferrooxidans* can tolerate up to 1 M Ni(II) (Dew *et al.*, 1999). Not much data is available for mercury resistance among prokaryotes. Huber and colleagues showed that the archaeon *Metallosphaera sedula* is sensitive to mercury at concentration 5 μM (Huber *et al.*, 1998). Guay *et al.* (1989) found that Ag(I) is the most toxic of the metals and ion studied: silver, cadmium, cobalt, copper, zinc, uranyl, and As(III). According to De *et al.* (1996) silver is toxic to *At. ferrooxidans* Fe(II) oxidation at a concentration of 0.93 μM (0.10 mg l^{-1}), however after previous silver exposure this species was tolerant to 1 mM (108 mg l^{-1}) Ag(I) (Ehrlich, 1984). *Leptospirillum ferrooxidans* and *Acidimicrobium cryptum* have been shown to be resistant to 500 and 300 mM Fe(III), respectively (Johnson *et al.*, 1992). Uranium, molybdenum, and chromium also were found in acidic environments, and a number of different acidophiles were isolated from the sites containing these metals. Wong *et al.* (1982) showed inhibition of growth of *At. ferrooxidans* by > 15 mM Cr(III). Bioleaching processes are now widely applied in bioremediation of polluted environments and for the extraction of certain metals from low-grade ores that would not be economically viable by any other method. The calculation of the contribution of bioleaching processes for acid mine drainages showed that the ferrous iron oxidation occurs with minor participation of extremely acidophilic prokaryotes. But moderately acidophilic iron and sulfur oxidizing bacteria appear to be more significant, particularly in waters of pH.3 (Hallberg and Johnson, 2003).

It should be noted that in systematics and taxonomy of acidophilic microorganisms hyperthermophilic and moderately thermophilic species could have absolutely distant and isolated phylogenetic positions on the levels of separate genera and families, and alternatively, among psychophilic acidophiles the specialized taxa are absent. The cases of identification of acidophiles in permafrost ecosystems were reported (Schleper *et al.*, 1995; Johnson, 1998), but it was shown only for mesophilic species *Acidithiobacillus* and only as a surviving process for psychrotolerant strains without detailed characteristics of metabolism.

Enzymes from acidophiles growing at $\text{pH} < 2-3$, such as amylases and glucoamylases are applied in the starch processing; proteases and cellulases are used in feed component; and oxidases are applied in the desulfurization of coal.

Observing the wide spectrum of phylogenetic diversity of microorganisms inhabiting low pH environments, different mechanisms of resistance to metals and low $[\text{H}^+]$, and various cell structure, it is possible to conclude that acidophiles as a major group of extremophiles, did not have a monophilic ancestral root, but instead the survival in aggressive acidity could possibly have resulted from widespread adaptive processes during the earliest stages of biological evolution on Earth. Furthermore, the existence of specialized genera and families for the hyperthermophilic and moderately thermophilic acidophiles representing separate distant taxa in systematics of prokaryotes and the absence of obligately psychophilic acidophilic microorganisms perhaps support our point of view on the origin, development, and distribution of the ancient original biota precisely in the acidic environment as a primary ecosystem on early Earth.

2. Thermophilic microorganisms.

Diversity of thermophiles. For most known species of *Eucarya* the temperatures around 100° C usually denature proteins and nucleic acids, increase the fluidity of membranes to lethal levels, and degrade chlorophyll (above 75° C), making photosynthesis impossible. Some members of two other domains *Bacteria* and *Archaea* could exist at much higher temperatures. Traditionally all thermophilic microorganisms are divided into three groups: moderately thermophilic (growth optimum at 50-60° C), thermophilic (optimum higher than 70° C), and hyperthermophilic (optimum higher than 80° C). *Archaea* consist of four phyla: *Crenarchaeota* (Sulfolobales-Thermoproteales branch), *Euryarchaeota* (extreme halophiles-Methanogens branch), "*Korarchaeota*" and "*Nanoarchaeota*". The "*Korarchaeota*" phylum (Barns *et al.*, 1994; Burggraf *et al.*, 1997) has been identified from analysis of DNA obtained from the natural environment without pure cultivation of organisms, and for this reason, its establishment remains not validated (the name written in quotations).

Another phylum of archaea, "*Nanoarchaeota*" (Huber *et al.*, 2003) is represented by a nanosized hyperthermophilic symbiont "*Nanoarchaeum equitans*" that does not even have a strain number, and lives obligately dependent upon *Igniococcus* strain KIN4/1, cells of which possess an outer membrane that it is a notable exception among the *Archaea* (Rachel *et al.*, 2002). This coculture was isolated from a sample of a marine hydrothermal system near Iceland, and grows under strictly anaerobic conditions between 70 and 98° C. Minimal doubling times (70 min for *Igniococcus* strain KIN4/1, and 45 min for "*N. equitans*") was observed at 90° C, pH 6, and 2 % NaCl concentration in the medium. By changing the gas mixture flow ($H_2:CO_2 = 80:20$) to 30 l min⁻¹ to allow removal of H₂S and increase the supply of H₂, the cell density of "*N. equitans*" was raised about 10-fold, while the cell density of *Igniococcus* remained unchanged. This means that "*N. equitans*" is sensitive to high concentrations of H₂S. It is interesting that during the late exponential growth phase of the coculture, about 80 % of the "*N. equitans*" cells detach from the surface of *Igniococcus* cells and occur freely in suspension. Separated cells "*N. equitans*" could not grow on cell homogenates of *Igniococcus* strain KIN4/1, and they require a physiologically active *Igniococcus* culture. Even experiments with separation of the co-culture cells by a semipermeable membrane have failed, resulting in the conclusion that direct cell contact with the "host" microorganism was required for growth of "*N. equitans*". Furthermore the parasitic lifestyle of "*N. equitans*" relatively to *Igniococcus* strain KIN4/1 is also not excluded. This co-culture demonstrated that symbiosis or parasitism occur not only between members of different domains, but even within the *Archaea*. In addition to the 16S RNA sequence of "*N. equitans*", three others sequences were received from the environmental high temperature biotopes (LPC33, OP9, and CU1). The sequence for LPC33 from the East Pacific Rise was identical to the sequence of "*N. equitans*", but two others showed difference. It was shown that genome size of "*N. equitans*" (490 kb) is the smallest of all known *Archaea*. Continuing study will show if it is due to the elimination of unused genes or if it is the original shortest primitive genome. The genetics study of "*N. equitans*" showed that this species has a large number of split genes (11 protein-coding and 4 tRNA genes). The splitting of tRNA genes has not been described previously, and the splitting occurs at the position, in which other archaeal tRNA contain an intron, and joining of tRNA halves is thought to be by way of a trans-splicing reaction. Furthermore, "*N. equitans*" encodes five of the nine proteins that are conserved in all *Euryarchaea* but they are not found in any of the *Crenarchaea* (Makarova and Koonin, 2005).

Subsequent discoveries of the members of “*Korarchaeota*” and “*Nanoarchaeota*” with a validly published new species could also validate the status of these phyla within the *Archaea*.

All other phyla of the domains Bacteria and Archaea have been validated, and have well-established phylogenetic positions. The phylum Crenarchaeota contains a single class, Thermoprotei, represented by the orders Thermoproteales, Desulfurococcales and Sulfolobales. All these orders are exclusively hyperthermophilic or thermophilic (Itoh, 2003). The order Thermoproteales is represented by two families, Thermoproteaceae (includes 5 genera: *Thermoproteus*, *Pyrobaculum*, *Thermocladium*, *Caldivirga*, and *Vulcanisaeta*) and Thermofilaceae (includes genus *Thermofilum*). The order Desulfurococcales comprise two families: Desulfurococcaceae (with genera: *Desulfurococcus*, *Staphylothermus*, *Aeropyrum*, *Sulphobococcus*, *Thermosphaera*, *Ignicoccus*, and *Thermodiscus*) and Pyrodictiaceae (include three genera: *Pyrodictium*, *Hyperthermus*, and *Pyrolobus*). The order Sulfolobales contains a single family, Sulfolobaceae, composed of six genera. Strictly aerobic genera: *Sulfolobus*, *Metallosphaera*, and *Sulfurococcus*; the facultatively aerobic genera *Acidianus* and *Sulfurisphaera*; and the strictly anaerobic genus *Stygiolobus*.

The phylum “*Euryarchaeota*” consists of eight classes and nine orders: Methanobacteria (including the order Methanobacteriales), Methanococci (with orders Methanococcales, Methanomicrobiales and Methanosarcinales), Halobacteria (order Halobacteriales), Thermoplasmata (order Thermoplasmatales), Archaeoglobi (order Archaeoglobales), and Methanopyri (order Methanopyrales). The hyperthermophiles of the phylum “*Euryarchaeota*” include both methanogenic and nonmethanogenic orders. Among the methanogenic orders there are both hyperthermophilic and mesophilic species. The order Thermoplasmatales contains three families comprising a single genus each: Thermoplasmataceae (genus *Thermoplasma*), Picrophilaceae (genus *Picrophilus*), and Ferroplasma (genus *Ferroplasma*). The order Thermococcales includes one family Thermococcaceae with three genera: *Thermococcus*, *Pyrococcus*, and *Paleococcus*. The order Archaeoglobales includes one family with three genera *Archaeoglobus*, *Ferroglobus*, and *Geoglobus*.

Thermophilic species of methanogenic archaea are distributed among orders Methanobacteriales (one genus *Methanothermobacter*), Methanococcales with genus *Metanococcus* (*M. thermolithotrophicus*, *M. jannaschii*, *M. igneus*), and Methanopyrales (genus *Methanopyrus*). Among the Bacteria two deep-branched and short lineages represented by the members of orders Thermotogales (includes the genera *Thermotoga*, *Thermosiphon*, and *Fervidobacterium*) and Aquificales (includes one genus *Aquifex*); they are all hyperthermophiles. Moderately thermophilic bacteria have enormous diversity in phylogeny, probably due to horizontal genes transfer during the latest stages of life evolution.

Physiology of thermophiles. The physiology of thermophilic microorganisms has a wide diversity, and they are usually well adjusted to their biotopes (Huber and Stetter, 1998). They could be primary producers of organic matter, as well as primary and secondary decomposers within the community. Chemolithoautotrophs and organotrophs with mixotrophic and heterotrophic anabolism occur among them. Some of them have respiratory catabolism and are dependent upon external electron acceptors (sulfur, iron, sulfate, nitrate, oxygen), or they could have fermentative metabolism. In *P. furiosus* the Embden-Meyerhof pathway was found to be modified and includes a number of novel enzymes – ADP-dependent glucokinase, ADP-dependent phosphofruktokinase, glyceraldehydes-3-phosphate ferredoxin oxidoreductase, phosphoenolpyruvate synthase, pyruvate-ferredoxin oxidoreductase, and ADP-forming acetyl-CoA synthetase (Sakuraba and Ohshima, 2002).

In particular, enzymes from hyperthermophiles, besides having high-temperature stability, are usually highly stable against detergents, organic solvents, and some other chemical reagents. Relative to pH, salinity and redox potential all thermophiles could be divided on several physiological groups: extreme acidophiles, moderate acidophiles, neutrophiles, alkaliphiles, etc.). They could be aerobic and anaerobic. But it is important to note that for deep-sea and deep underground ecosystems the biological standard state refers to neutral pH, which is not pH 7. Only at 25 °C and 1 bar pressure does the pH 7 represent neutrality, but at the pressure of water saturation and 100 °C, the neutral pH is 6.13, and at 200 °C, it drops to 5.64 (Amend & Teske, 2005).

Moderately thermophilic species of the genus *Picrophilus* still keep the record as the most acidophilic among extreme acidophiles (see section 1). Members of the genus *Sulfolobus* are strict aerobes growing autotrophically by oxidation of sulfur, hydrogen sulfide, or hydrogen, forming sulfuric acid or water as an end product. Members of the genus *Acidiamus* are able to grow by oxidation of hydrogen, organic matter, sulfur, sulfides, and even as *A. brierleyi* by leaching of sulfidic ores.

Among moderately acidophilic and neutrophilic thermophiles there are endemic species, for example species of the genus *Methanothermus* were found only in hot springs of the Kerlingarfjöll mountains in Iceland (Stetter *et al.*, 1981).

Thermococcus alkaliphilus is able to grow up to pH 10.5. But in nature alkaline hot springs are less widely distributed than acidic (at least at the bottom of the ocean).

Some hyperthermophiles appear to be highly adapted to toxic metal ions or metalloids (Holden and Adams, 2003). Arsenate and selenate are generally toxic to microorganisms; however, the crenarchaeon *Pyrobaculum arsenaticum* isolated from a hot freshwater spring near Naples, Italy, grows on hydrogen and carbon dioxide as its energy and carbon sources using 10 mM arsenate or 50 mM selenate as the electron acceptor. *P. islandicum* is unable to reduce arsenic or selenium, but it is capable of oxidizing hydrogen or components of peptone or yeast extract and reducing various metals such as Fe(III), U(VI), Tc(VII), Cr(VI), Co(III), and Mn(IV) at 100 °C. The iron reduction in *P. islandicum* occurs by a mechanism that differs from the NADH- and cytochrome *c*-dependent process that occurs in mesophilic iron reducers such as species of *Shewanella* and *Geobacter* (Childers and Lovley, 2001). *P. islandicum* does not contain *c*-type cytochromes and its NADPH is better than NADH as an electron donor for Fe(III) reductases activity ($K_m = 0.04$ and 3.33 mM).

Thermostable enzymes are classified into three groups (Yano and Poulos, 2003) defined by their range of temperature stability: moderately thermostable (45-65 °C), thermostable (65-85 °C), and extremely thermostable (>85 °C). Hyperthermophilic enzymes can have higher temperature optimum than original organism; for example, activity of amylopullulanase could have activity up to 142 °C (Schuliger *et al.*, 1993). In some literature the enzymes from thermophiles called thermozymes (Li *et al.*, 2005). Thermostable proteins generally show higher α -helical and β -sheet contents than mesophilic proteins (Nagi and Regan, 1997), and they show a very slow unfolding rate (Vieille & Zeikus, 2001), indicating that retaining the native structure in denatured conditions is anticipated. Also it was shown that thermostable proteins contain many charged and hydrophobic residues (Haney *et al.*, 1999). Extracellular enzymes like amylases, proteases, xylanases or pullulanases, and intracellular enzymes like dehydrogenases, oxidoreductases, and DNA polymerases were isolated from *Thermotoga maritima*, *Pyrococcus furiosus*, *Thermococcus* spp. and other hyperthermophiles. Three principle methods (*i.e.*, large-scale studies utilizing databases, the directed evolution, and the comparative structural analyses and protein engineering) have identified several factors that increase

thermal stability. These include a decrease in loop length and a concomitant increase in secondary structure; a decrease in labile residues (such as cysteines, asparagines and glutamines); an increase in aromatic stacking, increased hydrophobic interactions; increased metal-binding capacity; and increased oligomerization (Yano and Poulos, 2003).

Chromosomes of hyperthermophiles appear to be densely packed with genes, most of which are required for essential functions. This suggests that the earliest life forms may have had small genomes (Fujiwara 2002).

3. Barophilic microorganisms.

Pressure is a key physical parameter, which has influenced the evolution and distribution of both microorganisms and macroorganisms (Bartlett, 2002). Hominids evolved at an atmospheric pressure of 0.101 MPa (= 1 atmosphere = 1.013 bar), although our aquatic ancestors originated under hydrostatic pressure (Rothschild and Mancinelli, 2001). The oceans have an average depth of 3800 m and thus an average pressure of 38 MPa as well as a maximal depth of approximately 11000 m (~110 MPa). Hydrostatic pressure increases at a rate of 10.5 kPa per meter depth, compared with 22.6 kPa per meter for lithostatic pressure. Pressure decreases with altitude, so that by 10 km above sea level, atmospheric pressure is almost a quarter of that at sea level. The boiling point of water increases with pressure, so water at the bottom of the ocean remains liquid at 400 °C. Since liquid water at the Earth's surface does not normally occur above 100 °C, increased pressure can increase the optimal temperature for microbial growth, but usually by only a few degrees (Pledger *et al.*, 1994). High-pressure environments include deep lakes and the deep subsurface regions. It includes Lake Baikal in Siberia, Russia and Lake Vostok in Antarctica. Subsurface communities of microorganisms were found as deep as 3500 m below the surface (Szewzyk *et al.*, 1994). It is predicted that the largest number of prokaryotes in the biosphere occurs under ground (Whitman *et al.*, 1998).

The pressure changes the volume and compresses packaging of lipids resulting in decreased membrane fluidity. Many organisms on Earth have adapted to very high pressure, and a sudden change of pressure could be lethal. The Mariana trench is the world's deepest sea floor at 10,898 m, yet it harbours organisms that can grow at standard temperature and pressure. It is also inhabited by an obligately piezophilic species that can grow at 70 to 80 MPa, but not below 50 MPa (Kato *et al.*, 1998).

Barophilic bacteria are defined as those displaying optimal growth at pressures more than 40 MPa, whereas barotolerant bacteria display optimal growth at pressure less than 40 MPa and can grow well at a pressure of one atmosphere. Biodiversity of deep-sea ecosystems usually divides on hyperthermophiles and psychrophiles.

The bottom of the deep sea is a world exposed to extremely high pressure and low temperature (+ 2 °C) but in areas around hydrothermal vents the temperature can rise to 400 °C. The first barophilic bacterium was isolated from a depth of 10,500 m in 1979 (Yayanos *et al.*, 1979). This isolate grew at a pressure more than 100 MPa at 2 °C and more than 40 MPa at temperatures exceeding 100 °C. Takami *et al.* (1997) isolated thousands of microorganisms from the mud samples collected from the Mariana Trench. The agar plates in this investigation were incubated at 4-75 °C at atmospheric pressure or at 100 MPa for 1-4 weeks. The biodiversity was found to be composed of: actinomycetes, fungi, non-extremophilic bacteria and various extremophiles such as alkaliphiles,

thermophiles, three barophiles and psychrophiles. Kato with coauthors (Kato *et al.*, 1995; Kato *et al.*, 1996) have reported several high pressure adapted bacteria isolated from samples of deep-sea sediment from a depth of 2500-6500 m. Most of these isolates had barophilic and psychrophilic physiology. At atmospheric pressure strain DB6705 grew only at 4 °C, but not above 10 °C. At conditions of high pressure (>50 MPa) this strain was able to grow better at high temperature than at the lower temperature. Another extremely barophilic bacteria from the Mariana Trench could grow at pressure no less than 50 MPa. The isolates obtained by Yayanos became more barophilic at higher temperatures (Yayanos, 1986).

Hyperthermophilic barophiles *Thermococcus profundus* and *Pyrococcus horikoshii* were isolated from bacterial mat samples and hot fluid samples of hydrothermal vents in the Mid-Okinawa Trough at a depth of 1395 m. Another archaeon *T. peptonophilus* OG1 and SM-2 were isolated from hot fluid samples of hydrothermal vents in the Izu-Bonin forearc at a depth of 1380 m and in the South Mariana Trench at a depth of 1484 m respectively (Gonzales *et al.*, 1995). Maximal growth rates of *T. peptonophilus* were observed at 30-45 MPa whereas growth at 60 MPa was slower; the optimal growth temperature shifted from 85 °C at 30 MPa to 90-95 °C at 45 MPa. The maximum growth rate of *P. horikoshii* was observed at 95 °C under pressure of 0.1-15 MPa, and at 100 °C at a pressure of 30 MPa, with a doubling time of 35 min.

Some calculations indicate that all oceanic water passes through the basement rocks every 30 million years. Large parts of the new ocean floor consist of pillow lavas with amorphous volcanic glass on the pillow surface. The microscopy and culturing evidently suggest that microorganisms involved in the weathering of this volcanic glass (Fisk *et al.*, 1998). Channels, cavities and holes in the micrometer range penetrate the glass and the tips of the tunnels commonly stain with nucleic acid specific stains, suggesting the presence of microorganisms. The circulation of salt-water through the basement and the weathering activities of barophilic microorganisms in the flow paths therefore may have a significant impact on the composition of saltwater (Thorseth *et al.*, 1995).

The intraterrestrial microorganisms dwell in the pores of consolidated sediments and coarser, unconsolidated material, in fractures in hard rock and in fluid inclusions. Almost all very deep environments are anaerobic, with the exception of places where radioactivity may cause radiolysis of water, producing hydrogen and oxygen (Pedersen, 2000). The research of continental sedimentary rocks shows that living microorganisms are present and active down to the deepest levels studied (~3000 m). Large phylogenetically and culturally diverse communities of Archaea and Bacteria in North American deep subsurface continental sediments have been shown as well as other investigated underground sites in Africa and Scandinavia (Chandler *et al.*, 1997; Crozier *et al.*, 1999; Pederson *et al.*, 1996). Successful isolations of thermophiles have been performed from samples of sub-sea floor and continental oil reservoirs, indicating that Bacteria and Archaea (*Archaeoglobus fulgidus*, *Thermodesulforhabdus norvegicus*, *Thermotoga subterranea*) inhabit deep hot oil reservoirs (Beeder *et al.*, 1994; Beeder *et al.*, 1995; Jeanthon *et al.*, 1995).

Concerning the cultivation of microorganisms from samples of super deep wells it is important to note that the borehole windows into super deep environments are still very few and none of them have been drilled with the microbiological purpose. The deepest well SG-3, 12262 m deep, occurs in the Pechenga-Zapolyarny area of Kola Peninsula, Russia. The reports about isolation of microorganisms from such ecosystems have positive results with isolation but none of them had good negative control for the exclusion of contamination (Pedersen, 2000).

Igneous rocks are the predominant solid constituents of the earth, formed through cooling of molten or partly molten material at or beneath the earth's surface. The studies in the Swedish long-term nuclear waste disposal research programme on subterranean microbiology revealed previously unknown microbial ecosystems in igneous rock aquifers at depths exceeding 1000 m (Pedersen & Ekendahl, 1990). The study of subterranean microbial diversity led to isolation of pure cultures of methanogens, methanotrophs, and sulfate-reducing bacteria (Kotelnikova and Pedersen, 1998; Kotelnikova *et al.*, 1998; Motamedi and Pedersen, 1998; Kalyuzhnaya *et al.*, 1999).

Barophilic enzymes, currently referred to as piezostable enzymes, are stable at high pressures and have been isolated from a wide variety of organisms, most of which are either thermophilic or psychrophilic and have optimal growth at above one atmosphere (Abe & Horikoshi, 2001).

New findings from deep-sea ecosystems have important applications in medicine and biotechnology. Mesophilic *Vibrio diebolicus* isolated from a vent sample, secretes an innovative exopolysaccharide of potential medical interest for its chemical resemblance to heparin, which delays the onset of blood clotting (Raguenees *et al.*, 1997). Another new mesophilic vent strain of *Pseudomonas aeruginosa* is remarkable for its tolerance to high concentrations of cadmium (up to 5 mM). It not only tolerates but completely (>99 %) removes the cadmium from solution by precipitation onto the cell wall, making it a prime candidate for applications in heavy-metal recovery and environmental remediation (Wang *et al.*, 1997).

It was reported that a high-pressure-regulated system for the gene expression is found not only in deep-sea-adapted microorganisms, but also in bacteria growing at atmospheric pressure (Sato *et al.*, 1995; Welch and Bartlett, 1996). These results suggest that the systems developed in the high-pressure environment may be conserved in organisms growing at atmospheric pressure, possibly indicating that life emerged from the deep-sea environment a long time ago. Pressure is the only stressor known to simultaneously induce many heat shock and cold-shock proteins, sets of proteins, which are induced following exposure to opposing thermal regimes. The major pressure-inducible protein is an unknown small highly basic protein. The heat-shock proteins may also be induced in piezophiles upon decompression. A stress protein showing similarity to a heat-shock protein, was found to be induced in the deep-sea piezophilic hyperthermophile *Thermococcus barophilus* at the cultivation under atmospheric pressure (Marteinsson *et al.*, 1999). Other interesting effects of high pressure on the structural unites of biological cells have been discussed in a previous review (Bartlett, 2002).

4. Psychrophilic microorganisms.

Already in 1840 Hooker observed living algae associated with ice. In 1887 Forster was the first to report microorganisms growing well at 0 °C that were isolated from fish (Hoyoux *et al.*, 2004). The first mention of the term "psychrophile" (cold loving) was made by Schmidt-Nielsen in 1902 for the description of bacteria capable of growing at 0 °C. Later the term was also used to refer to a number of species of eukaryotic organisms; including yeasts, diatoms, algae, lichens, mosses, insects and fish. The current biological definition for psychrophilic microorganisms is the ability to grow at low and subzero temperatures, and the inability to grow with absence of metabolic activity at a temperature higher than 15 °C. True psychrophilic microorganisms have cell lysis at room temperature, and their enzymes and proteins are very sensitive and destabilized very fast without being maintained in ice. Psychrotolerant microorganisms can have high metabolic activity and be

capable of growth with a prolonged lag phase at subzero and low temperatures, but unlike psychrophiles they do not die at room temperatures and they usually have an optimum growth in the range for mesophilic microorganisms. Here we specially ignore the term “psychrotrophic”, since the exact translation means “eating cold” that is inappropriate due to the tolerance of cold, but not trophy.

In north geographical regions, the surface of large snow fields are sometimes observed colored pink or red, due to the growth of snow algae, such as a species of *Chlamydomonas nivalis*, which produce brilliant red spores. These organisms alter the albedo of the snow to induce a snowmelt and increase the availability of liquid water for other psychrophilic microorganisms. Some diatoms and other psychrophiles produce extra-cellular enzymes and ice-active substances that lead to the pitting of ice and melting the surrounding ice making the liquid water available, which is necessary for metabolic activity (Raymond *et al.*, 1994). It is now known that most (if not all) Antarctic Sea-Ice diatoms are able to synthesize these large macromolecular substances that appear to be glycoproteins (Raymond, 2000). Among higher evolved living creatures, as for example some polar fish, the production of glycerol serves as the protection against freezing (Raymond, 1995). The synthesis of such cryoprotectants leads to freeze avoidance. However, some organisms have developed mechanisms of freeze tolerance, involving drastic metabolic modifications during frost hardening, and resistance to membrane damage, solute concentration and dehydration accompanying ice crystallization (Franks, 1985).

Cryoprotectants such as glycerol or DMSO are water-miscible liquids, and they penetrate the cell and protect it from freezing by reducing the severity of dehydration effects and preventing the formation of ice crystals. In microbiological laboratories, the lyophilization (dry freezing) is a traditional way of preserving microbial cultures under vacuum at very low temperatures (-70 to -196 °C), that means at specific conditions biological cells are indeed capable of surviving deep freezing, and this feature probably is universal for life.

Ecosystems and microbial diversity of psychrophiles.

The ecosystems of Earth with permanently low temperatures include the regions of the Arctic and Antarctic with polar ice sheets, glaciers and permafrost, the snow-caps and glaciers of high mountains, and the deep water and marine sediments of the Oceans. In the polar regions, enormous areas of frozen ground are formed – termed permafrost, which are defined as the thermal condition in which soils and sediments remain at or below 0 °C for two or more years in succession (Wagner *et al.*, 2001). The terrestrial permafrost, which underlies more than 20 % of the world’s land area, is above all controlled by climatic factors and characterized by extreme terrain conditions and landforms. On Earth the thickness of the permafrost can reach several hundreds of meters, e.g., it is ~600 to 800 m thick in East Siberia (Central Yakutia). During a relatively short period of the polar summer, a thin surface zone (active region) of the permafrost sediments undergoes thawing.

The largest low-temperature ecosystem on Earth is the deep sea floor since oceans cover three quarters of the surface of the planet. The psychrophiles that inhabit this global-scale ecosystem (with a constant temperature of 4 °C below a depth of 1,000 m) are true extremophiles as they are adapted not only to low temperatures, but also to further environmental constraints (Feller and Gerday, 2003). In the ocean depths and sediments they are faced with extremely high pressures, and therefore must be piezo-psychrophiles (or baro-psychrophiles). The microbial communities that are found in sea ice, which comprise bacteria, algae, fungi and protozoa, are exposed to salt concentrations of several molar in brine veins at -20 °C, and are therefore halo-psychrophiles (Staley and Gosink,

1999). On the snow surface of glaciers and polar caps, psychrophiles are exposed to strong ultraviolet radiation (Carpenter *et al.*, 2000). The endolithic microbial communities that are found in rocks of the Antarctic dry deserts, which comprise lichens, yeasts, Cyanobacteria and heterotrophic bacteria, survive low water and nutrient availability (Friedmann, 1982). In alpine caves and cracks, microorganisms also evolve in a poor environment in the absence of light (trogl-psychrophiles).

The physiological and phylogenetic diversity of psychrophilic and psychrotolerant microorganisms that was detected and isolated from cold ecosystems is very broad. This includes many eukarya and almost all main physiological groups of eubacteria and some archaea. Isolation of diatoms and other algae, cyanobacteria, fungi and yeast were also reported as common findings in cold environments. Active anaerobes that have a truly psychrophilic nature occurred in the following physiological groups: fermentative bacteria, methanogens, acetogens, sulfate-reducers, iron-reducers, and nitrate-reducers. In the last decade more attention has been paid to the search for psychrophilic strains of methanogens, since many (if not most) of methanogenic habitats occurred in cold climates (swamps of Siberia in Russia and of Canada). Also methane-oxidizing aerobic bacteria were found in the cold swamp water of the Arctic region (Trotsenko and Khmelenina, 2002). The first Gram-positive sulfate-reducing bacterium from Antarctica *Desulfotomaculum antarcticum* strain No.64 was isolated in 1968 (Iizuka *et al.*, 1969). However, truly psychrophilic Gram-negative sulfate-reducers capable of growth at -1.8°C and within optima of growth at $7-10^{\circ}\text{C}$ were isolated only in 1999 from permanently cold marine sediments off the coast of Svalbard (Knoblauch *et al.*, 1999).

The producers (or synthetics) of organic compounds in cold environmental ecosystems could include not only photosynthetics (Cyanobacteria, diatoms, green and brown algae), but also chemolithotrophic bacteria. Unique photosynthetic processes were found on the deep sea floor near black-smoker hydrothermal vents (Van Dover *et al.*, 1996; Kolber *et al.*, 2001); bacteria that could perform photosynthesis by using light in infrared wavelengths from high-temperature vents. These bacteria could switch photosynthesis on and off and are photoheterotrophic (require organic compounds for anabolism). These bacteria are not psychrophilic, but they could provide a primary biomass for the decomposition by organotrophic truly psychrophilic species that inhabit the deep-sea floor.

Mechanisms of adaptation to low temperatures.

Every microorganism is adapted for optimum functioning in natural ecological environments and it is reflected in its physiological characteristics: range and optima of temperature, pH, salinity, etc. Any extreme change in environmental conditions from the optimum inflicts a stress on an organism. The intensity and time duration of the change usually determines whether the organism is killed, ceases growth, or has an increased lag phase of growth and reduced biomass yield (Ray 1986; Russel *et al.*, 1995). Most bacteria are able to tolerate small changes in an environmental parameter and can adapt over the time scale of minutes, hours, or days. Microorganisms could perform this by both yielding to the stress conditions and making suitable provisions for survival or attempting to resist the stress. For most microorganisms, this tolerance can be pushed to maximum limits if the cell is provided with a sufficient opportunity to sense and adapt to the deteriorating environment. Psychrophiles have adapted their lifestyles to prefer (and sometimes to require) these extreme environments. Changes in environmental conditions away from the optimal value can cause the induction of many elaborate stress responses. These strategies are generally directed at survival rather than growth. According to recent reviews, the mechanisms of adaptation to cold temperatures

could be connected with the changing of capacities of proteins (more flexible structural and conformation changes), increasing the fluidity of membranes by the changing of the unsaturation degree of fatty acids, modifications to *ante-iso-iso-* branching patterns, and by shortening in the fatty acid chain length. Also, the synthesis of antifreeze glycoproteins and peptides can further depress the freezing point of cellular water.

Unsaturation of fatty acid chains is the change that is most commonly found to occur when the temperature is reduced. This increases the fluidity of the membrane because unsaturated fatty acid groups create more disturbances to the membrane than saturated chains and it is achieved by desaturases situated in the membrane itself and thus are able to react quickly. Also, the average fatty acid chain length may be shortened, which would have the effect of increasing the fluidity of the cell membrane. After a fall in temperature, an increase in the amount and/or kind of branched fatty acids may occur. Sometimes, there may be a reduction in the proportion of cyclic fatty acids and thus an increase in mono-unsaturated straight chain fatty acids.

Sudden decreases in temperature initiates a specific alteration in gene expression – cold-shock response. It involves the induction and the synthesis of cold-shock proteins. Cold shock response can involve the expression of up to 50 different cold shock proteins depending on the species, as well as the rate and extent of temperature drop. There are two groups of proteins produced at the gene expression during cold shock response. Cold shock proteins (Csps) are synthesized at low temperature, and the larger the severity of the shock, the more Csps are produced. A second group of cold-induced proteins, the cold acclimation proteins (Caps) has also been described that are comparable to Csps and are continuously synthesized during prolonged growth at low temperatures, and differentiate psychrophiles from mesophiles. Cold shock proteins can stabilize mRNA and re-initiate protein production. Others are also linked to maintaining the fluidity of the membrane such as inducible desaturases.

The regulation of Csp synthesis is multifactorial, and controlled at the levels of transcription and translation, as well as mRNA and protein stabilities. The main function of Csp are in the regulation of cellular protein synthesis, at the level of transcription as well as the initiation of translation, and in mRNA folding, acting as a chaperone preventing the formation of an mRNA secondary structure. The cellular function of acclimation proteins remains not well understood.

Other adaptive strategies developed by psychrophilic organisms involve the regulation of ion channel permeability, seasonal dormancy and microtubule polymerization (Detrich *et al.*, 2000). The key adaptive strategy of psychrophiles is the modification of enzyme kinetics, allowing the emergence of metabolic rates compatible to life at low temperatures. The psychrophilic enzymes could be highly active at low temperatures and could be 10 times higher than that of their mesophilic homologues.

5. Halophilic microorganisms.

Halophilic microorganisms require salt in media for growth. Halotolerant microorganisms are not dependent upon the salts in growth media but able to tolerate up to 15% salinity. Halophiles are found in each of the three taxonomic domains: Archaea, Bacteria, and Eukarya. The metabolic diversity of halophiles is very wide: oxygenic and anoxygenic phototrophs, aerobic heterotrophs, fermenters, denitrifiers, sulfate-reducers, and methanogens (Oren, 2002). With increasing salinity the diversity of metabolic types decreases dramatically. The maximal salinity limit, at which each

dissimilatory process occurs, is correlated with the amount of energy generated and energetic cost of osmotic adaptation in environment.

Halobacteria include red-pigmented extremely halophilic archaea, members of the *Halobacteriaceae* family, the only family in the Halobacteriales order. Most halobacteria require 1.5 M NaCl both to grow and maintain the structural integrity of the cells. Halobacteria can be differentiated from halophilic bacteria on the basis of their archaeal characteristics, in particular the presence of ether-linked lipids. Most halobacteria are colored red or orange in cell vesicle due to the presence of C-50 carotenoids, but some strains are colorless, and those with gas vesicles form opaque, white, or pink colonies. A purple hue may be seen in halobacteria that form the bacteriorhodopsin-containing purple membrane. Halobacteria exhibit features characteristic of the archaea, including eukaryotic-like transcription and translation mechanisms, ether-linked lipids and like some bacteria, a cell wall S-layer composed of glycoprotein. A unique feature of halobacteria is the purple membrane, specialized regions of the cell membrane that contain a two-dimensional crystalline lattice of a chromoprotein, bacteriorhodopsin. Bacteriorhodopsin contains a protein moiety (bacterioopsin) and covalent bound chromophore (retinal), and it acts as a light-dependent transmembrane proton pump. The membrane potential generated can be used to drive ATP synthesis and support a period of phototrophic growth. Bacteriorhodopsin is induced by low oxygen tension and high light intensity. Also, halobacteria contain large quantities of red-orange carotenoids, which have been shown to be necessary for stimulating an active photorepair system to repair thymine dimers resulting from ultraviolet radiation. The most abundant carotenoids are C-50 bacterioruberins, although smaller amounts of biosynthetic intermediates such as β -carotene and lycopene are also present. Retinal is produced by oxidative cleavage of β -carotene, a step that requires molecular oxygen. Several retinal proteins, in addition to bacteriorhodopsin, are also synthesized in halobacteria, including halorhodopsin, which is an inwardly directed light-driven chloride pump, and two sensory rhodopsins that mediate the phototactic response (swimming towards green light and away from blue and ultraviolet light). Like many aquatic bacteria, halobacteria produce buoyant gas vesicles that are hollow proteinaceous structures surrounding a gas filled space. The function of gas vesicles is to enable the cells to float to the more oxygenated surface layers since halobacteria are primarily aerobic and live in concentrated brines in which the solubility of molecular oxygen is low (especially at high temperatures). This also increases the availability of light for purple membrane-mediated photophosphorylation (Joo & Kim, 2005),

Adaptation to the hypersaline environment. There are two fundamentally different strategies within the microbial world that enable microorganisms to cope with the osmotic stress inherent to the presence of high salt concentrations. The first strategy is when cells maintain high intracellular salt concentrations, osmotically at least equivalent to the external concentrations (the “salt-in” strategy). In this case all intracellular systems should be adapted to the presence of high salt concentrations. The second strategy is when cells maintain low salt concentrations within their cytoplasm (the “compatible-solute” strategy). In this case the osmotic pressure of the medium is balanced by organic compatible solutes, and here no special adaptation of the intracellular systems is required (Oren, 1999). Microorganisms that balance the osmotic pressure of the medium with high concentrations of organic osmotic solutes also have to expend energy in ion pumps to keep intracellular ionic concentrations low and to counteract the diffusion of inorganic salts through their membranes. Na^+/H^+ antiporters have been characterized from moderately halophilic bacteria such as *Salinivibrio costicola* that mainly use organic solutes. These antiporters play a key role in keeping

the intracellular Na^+ concentrations low. The activity of the Na^+/H^+ antiporters in the cytoplasmic membrane of halotolerant alga *Dunaliella salina* have been increased at the cultivation of cells in higher NaCl concentrations of medium. The need for Na^+ pumping thus may increase the energy cost of the “compatible-solutes” strategy beyond that of the synthesis of the osmolytic compounds only. The “salt-in” strategy is applied in two phylogenetically unrelated groups: aerobic extremely halophilic archaea of the order *Halobacteriales* and anaerobic halophilic bacteria of the order *Haloanaerobiales*. No organic osmotic solutes have been found in representatives of these groups, and most studies report that the intracellular ionic concentrations are similar to those of the surrounding media. The single exception of these observations is *Desulfovibrio halophilus* that could accumulate salt inside of cytoplasm and at the same time it could provide the osmotic balance by trehalose and glycine (Welsh et al., 1996). The intracellular salt composition is usually represented by molar concentrations of KCl.

The energetically cheaper salt-in option requires extensive adaptations of the intracellular machinery, and these can be achieved only in a long and complex evolutionary process. This option is used by a few specialized groups of prokaryotes only. All other organisms use organic osmotic solutes, which are expensive to produce (Oren, 1999). When organic solutes are accumulated, there is a clear preference at the highest salinities for the solutions with smaller molecules, which are energetically easier to produce.

A survey of the halophilic microorganisms shows that not all known metabolic types function in the highly mineralized environments. While oxygenic and anoxygenic photosynthesis, aerobic respiration, fermentation, and denitrification can occur at or close to NaCl saturation, other physiological groups of microorganisms have never been shown to thrive at high salinities. Autotrophic methanogens, acetoclastic methanogens, dissimilatory sulfate-reducers with complete oxidation, autotrophic ammonia and nitrite oxidizers have not been found at salinities above 15 %. Only one halophilic autotrophic sulfur oxidizer *Thiobacillus halophilus* has been isolated in pure culture; this organism oxidizes thiosulfate, tetrathionate, and elemental sulfur in media with up to 24 % NaCl (optimum is at 5-6 % NaCl).

In most cases a good correlation exists between the upper salt concentration, at which the different microbial processes have been shown *in situ* to occur and the ability of the corresponding microbial pure cultures to grow at high salt concentrations *in vitro*. Most microorganisms that produce organic osmolytes are also able to take up such solutes from the environment when it is available, thereby reducing the energy cost of life at high salt concentrations. Such a situation could occur in complex microbial communities where different metabolic types of microorganisms coexist, some of which may produce and excrete the osmolytic molecules in the environment. The inclusion of different kinds of osmolytes in the enrichment media for isolation of new halophiles may be recommended. That may explain why the “missing” physiological groups of halophiles remain undiscovered.

Usual proteins at high salt concentrations are in general destabilized due to enhanced hydrophobic interactions. Halophilic proteins have evolved specific mechanisms that allow them to be both stable and soluble in the high KCl cytoplasmic concentration. Halophilic proteins have specific structural elements to fold into proper native form at high salt concentrations. Since all soluble halophilic enzymes have a highly negative surface charge and once folded properly their flexibility may be achieved by the repulsion forces between close charges. The instability caused by the high surface charge should somehow be compensated to avoid unfolding and it is believed that the role of high salt concentrations is to shield this high surface charge. All halophilic proteins are highly negatively

charged with hydrated carboxyl groups, which maintain the solubility of proteins at high salt concentrations. The electrostatic repulsion offsets destabilization from the hydrophobic effect enhanced by salt. In the case of malate dehydrogenase (hMDH) from *Haloarcula marismortui* whose three-dimensional structure has been determined by X-ray crystallography, the requirement for high NaCl or KCl concentrations for the stabilization can simply be explained by a specific low affinity binding of salt ions to the folded protein. Therefore, molar amounts of salt are necessary to saturate these binding sites.

The osmolytes that allow halophiles and halotolerant microorganisms maintain their active physiology in a highly saline environment (the “compatible-solute” strategy), were comprehensively reviewed by Roberts (2005).

Some of the extremely halophilic archaea produce bacteriocins called halocins; they inhibit a broad spectrum of bacteria. Bacteriocidal action of halocins includes inhibition of transcription, translation, Na^+/H^+ antiporter, DNA and RNA nuclease activity, pore formation, bacteriolysis and disruption of cellular membranes.

Recently it was shown that within *Halobacteriaceae* there are many species capable of *p*-hydroxybenzoic acid degradation. The forty-four new halophilic archaea were found to be able to grow in 0.4 mM *p*-hydroxybenzoic acid as the sole carbon and energy source (Cuadros-Orellana *et al.*, 2006).

Due to the importance in biotechnology of the search for new molecules from halophiles the intensive work with isolation of new strains from hyper-saline ecosystems is continuing. One hundred and sixty-five halophilic archaea were isolated from three different types of hypersaline lakes (Erliannor, Shangmatale and Xilin soda lake) in Inner Mongolia (Pan *et al.*, 2006). During the study of prokaryotic biodiversity in El Golea salt lake, in Algerian Sahara, extreme halophilic microorganisms were isolated from a site containing 290 g l^{-1} NaCl (Hacène *et al.*, 2004). The investigation of halophilic archaea in the crystallizers of Adriatic Sečovlje salterns by using gene fragments encoding 16S rRNA and bacteriorhodopsin as molecular markers resulted in 180 clones that were screened, and as a result 15 different 16SrRNA and 10 different bacteriorhodopsin phlotypes were detected (Pašić *et al.*, 2005). Archaeal diversity along a transient soil salinity gradient at Salt Spring in British Columbia, Canada, was investigated with cultivation-independent approach (Walsh *et al.*, 2005).

The biotechnological application of halophiles is continuously increasing. Halophilic microorganisms produce stable enzymes (DNAses, lipases, amylases, gelatinases, and proteases) capable of functioning under high concentrations of salt, which have precipitated or denatured most known proteins, and most of these enzymes inactivated and denatured at NaCl concentrations below 1M. Compatible solutes now used as stabilizers of enzymes, DNA, membranes, and whole cells. Ectoine for example, stabilize polymerase chain reaction. Trehalose is used as a cryoprotectant in freeze-drying processes of biomolecules. Biosurfactants and halophilic exopolysaccharides enhance the remediation of oil-contaminated soil and water. Ether-linked lipids from archaeal halophiles have a high chemical stability and resistance against esterases and thus a higher survival rate than liposomes based on fatty acid derivatives. Halophiles used in bioremediation of oil-polluted saline ecosystems. Some halophiles are used for the bioremediation of pollutants by halogenated organic compounds. *Lactobacillus plantarum* is used for the fermentation of pickles and sauerkraut. *Anabaena torulosa* is used in amelioration of soil salinity during crop growth.

The significance of halophiles for paleontology is of great concern as it has been reported that some halophiles were found alive in Permian, Jurassic, or even more ancient salt crystals as has been reviewed by McGenity *et al.* (2000). Also halophiles are important to the question of possible biodeterioration of containers that store toxic and nuclear waste, buried in deep underground salt caves.

6. Alkaliphilic microorganisms.

Alkaliphilic microorganisms are microorganisms that require a growth media with pH higher than 8.0, and cannot grow at neutral pH. There are some species that have two optima pH in dependence upon growth conditions, particularly nutrition. The most alkaliphilic microorganisms are cyanobacteria that are capable of growth at pH 12-13. Alkalitolerant organisms have optimal pH for growth at neutral pH, but are able to grow at pH 10-11. According to Horikoshi (1999) all alkaliphiles are divided on two main physiological groups of microorganisms: alkaliphiles and haloalkaliphiles. Haloalkaliphiles require high pH and they also require the presence of NaCl in the growth media. Some of the alkaliphiles and haloalkaliphiles are obligately dependent upon CO_3^{2-} ions, especially microorganisms isolated from soda lakes.

It is interesting that alkaliphilic microorganisms could be isolated from neutral or even acidic soils and feces. Two anaerobes *Anoxybacillus pushchinensis* and *Desulfotomaculum alkaliphilum* were isolated from manure with neutral pH (Pikuta *et al.*, 2000a; Pikuta *et al.*, 2000b). Historically aerobic alkaliphiles were primarily isolated on media with high pH adjusted by NaOH solution; later after the study of chemical composition of highly mineralized soda lakes the media for isolation of truly alkaliphilic microorganisms were based on $\text{NaCO}_3/\text{NaHCO}_3$ buffers. With developing an anaerobic technique additional reducers (sulfide, thiosulfate etc.) were applied for isolation of alkaliphilic anaerobes. And as it was for extreme halophiles the application of highly mineralized media (in case for alkaliphiles it was carbonate/di-carbonate composition) led to the discovery of numerous phylogenetically distant new genera and families (Zavarzin *et al.*, 1999; Zavarzin & Zhilina, 2000).

The first alkaliphilic strain *Streptococcus faecalis* was isolated in 1928 (Downie and Cruickshank, 1928), but much earlier the communications about alkalitolerant nitrifying bacteria were reported (Meek and Lipman, 1922). The works of Horikoshi showed the presence of extreme alkaliphiles (*Clostridium* and *Bacillus*) in soils with neutral pH. In soda lakes the dominant microorganisms are cyanobacteria including following genera: *Spirulina*, *Chroococciopsis*, *Nostoc*, *Cyanospira*, *Gleocapsa*, *Nodularia*, *Synechococcus*, *Synechocystis*, *Calothrix*, *Scytonema*, *Anabaena*, *Lamprocystis*, *Thiocapsa*, *Thiocystis*, *Chromatium*, *Amoebobacter*, *Thiospirillum*, *Rhodobacter*, *Ectothiorhodospira*, *Aphanothece*, *Chamaesiphon*, *Rhabdoderma*, *Phormidium*, *Oscillatoria*, etc. Tindall with co-researchers described aerobic, alkaliphilic, extremely halophilic archaea belonging to genera *Natronobacterium* and *Natronococcus* (Tindall *et al.*, 1984). *Methanobacterium thermoalkaliphilum* and *Methanosalsus zhilinaeae* are examples of anaerobic archaea isolated from soda lakes. The aerobic heterotrophic spore-forming alkaliphilic eubacterium (belonging to the genus *Bacillus*) was isolated in 1983 from highly mineralized lake Wadi Natrun in Egypt. This bacterium was tolerant to salinity of more than 4 M NaCl (Weisser and Trüper, 1985). Alkaliphilic anaerobic sugarlytic spirochetes also were found to be halotolerant, and some of them require NaCl for growth (Zhilina *et al.*, 1996; Hoover *et al.*, 2003). The discovery of alkaliphilic

sulfate-reducing bacteria finally had confirmed the long-time existing hypothesis of Abd-el-Malek about functioning of sulfidogenes in hypersaline environments (Abd-el-Malek and Rizk, 1963). After the application of specific media with carbonate/bicarbonate buffers the presence of sulfate-reducers (*Desulfonatrenovibrio hydrogenovorans*, *Desulfonatronum lacustre*, *D. thiodismutans*, and *D. cooperativum*) was shown in soda lakes of African, Asian, and American continents (Zhilina *et al.*, 1997; Pikuta *et al.*, 1998; Pikuta *et al.*, 2003c; Zhilina *et al.*, 2005). One of the alkaliphilic sulfate-reducers was found positioned very distantly from known phylogenetic lineages and represented separate family *Desulfonatronaceae* (Kuever *et al.*, 2005).*

The alkaliphilic enzymes from some alkaliphiles have optimal activity at pH much higher than physiological pH growth optimum of organisms, and this capacity is used quite well in biotechnology (detergents and wash powders).

Alkaliphilic microorganisms develop at high concentrations of Na⁺ ions and, similarly to halophiles, have to accumulate osmoprotective compounds to resist the osmotic stress. Due to a high environmental pH the alkaliphiles must maintain their intracellular pH and realize different pathways for ion transport. All these properties of alkaliphilic organisms suggest that their adaptation to extreme environmental conditions during evolution had occurred at the system level (Zavarzin *et al.*, 1999). According to academician Zavarzin, the basic difference between the alkaliphilic and halophilic microbial communities consists of the fact that soda mineralization resulted from the formation of athalassic inland water bodies rather than thalassic ones. For this reason, the prokaryotic alkaliphilic community can be regarded as a relic analogue of the continental biota in the early Proterozoic. In order to answer the question of whether the alkaliphilic community can indeed reflect the terrestrial source that gave rise to the biological diversity of prokaryotes, one should find out if the biodiversity of this community is sufficient to include representatives of all major branches of the universal phylogenetic tree of prokaryotes and show that the alkaliphilic community is not a dead-end line of evolution, like hyperthermophiles, out of which it is impossible to derive total diversity.

A recent review compared the mechanisms of pH homeostasis in neutrophilic bacteria and alkaliphilic bacilli (Padan *et al.*, 2005). It was shown that active alkaline pH homeostasis that is important for both survival and growth, significantly depends upon monovalent cation/proton antiporters, the identification of which is generally achieved by combining genetic and biochemical approaches.

7. Microorganisms resistant to radiation.

The beginning of a study of microorganisms resistant to high radiation was initiated after the remarkable discovery of *Deinococcus radiodurans* in 1956 during the process of food sterilization by X-ray radiation. But in contrast to other types of extremophiles among radio-resistant microorganisms there are no known microorganisms that are true radiophiles. It is indeed difficult to prove that such types exist, since true radiophiles theoretically should not be able to live at certain radiation level, but should require high irradiation levels for growth. Life without a certain level of radiation is impossible, and this feature of life unfortunately was never mentioned in any biological definition of life. The evolution of organisms that are able to grow continuously at 6 kilorads (60

* The original spelling, *Desulfonatronumaceae* (sic), has been corrected on validation according to Rule 61. Reference: Lists editor, *IJSEM*: Validation list n° 107 (footnote). *Int.J.Syst.Evol.Microbiol.*, 2006, 56: 1-6.

Gy) h⁻¹ or survive acute irradiation doses of 1,500 kilorads is remarkable, given the apparent absence of highly radioactive habitats on Earth over geologic times. Notwithstanding a few natural fission reactors like those that gave rise to the Oklo uranium deposits (Gabon) two billions years ago, the radiation levels in the Earth's surface environments, including its waters containing dissolved radionuclides, have provided only about 0.05 to 20 rads/year over the last four billions years (Makarova *et al.*, 2001). DNA damage is readily inflicted on organisms by a variety of common physicochemical agents (UV light or oxidizing agents) or non-static environments (cycles of desiccation and hydration or cycles of high and low temperatures) and it seems more likely that radiation resistance evolved in response to chronic exposure to non-radioactive forms of DNA damage. In the exponential growth phase, cells of *D. radiodurans* do not die in response to ionizing irradiation up to 0.5 megarad and they show 10 % survival at 0.8 megarad (Moseley and Evans, 1983). The resistance to UV for *D. radiodurans* cells is 33-fold more than for *E. coli* cells. After complete study of *D. radiodurans* R1 genome (Makarova *et al.*, 2001) the following conclusions were made: 1) The expanded Nudix hydrolase superfamily and the homologs of plant desiccation resistance-associated proteins are likely to contribute to both the extreme radiation and the desiccation resistance of *Deinococcus*; 2) A variety of other proteins, particularly those that belong to expanded families, are likely to be involved in the unusual phenotype of this bacterium; 3) The unexpectedly numerous nucleotide repeats may play a role in stress response; 4) The transferred entire gene systems, such as genes encoding pili-associated functions, could contribute to the formation of surface structures involved in secretory systems similar to the type III secretion pathway (since there is no experimental data concerning pili formation in *D. radiodurans* cells). The conclusion after other proteomic research of *D. radiodurans* (Mrázek, 2002) suggests the application of this organism for bioremediation of radioactive waste sites. Earlier Daly showed a realistic approach for engineering *D. radiodurans* for bioremediation, metal remediation, and toxic organic compound degradation (Daly, 2000). The latest comparative genomic research of *D. radiodurans* and *Thermus thermophilus* was performed by Omelchenko *et al.* (2005).

Other radiation resistant bacteria were found during food sterilization (Hastings *et al.*, 1986; Lacroix and Lafortune, 2004). Seven strains of *Lactobacillus sake* were more resistant to gamma-irradiation in the exponential (log) phase (0.82 kGy) than in the stationary phase of growth (0.64 kGy). *Pantoea agglomerans* was found to be a natural contaminant of carrots and radiation resistant bacterium (it survived after 0.15 kGy doses).

Much research dealing with the isolation of radiation-resistant bacteria without exposure to irradiation were performed. Some strains were selected by the resistance to desiccation, and other by hot spring habitats (Sanders and Maxcy, 1979; Albuquerque *et al.*, 2005). A radiation resistant moderately thermophilic species *Truepera radiovictrix* was described during the study of isolates from hot spring runoffs on the Island of São Miguel in the Azores; this species phylogenetically was very distant from known genera and families, and exhibited 60 % survival after exposure to 5.0 kGy of gamma irradiation.

Anellis with colleagues performed experimental measurements on the survival of different strains during Cobalt gamma irradiation at different temperature regimes (Anellis *et al.*, 1973). It was shown that at low temperatures (-80 °C) during ionization the most resistant microorganisms were: *D. radiodurans*, *Streptococcus farcalis*, and *Alcaligenes faecalis*.

Experiments with the desiccation-tolerant cyanobacterium *Chroococcidiopsis* demonstrated high resistance of this microorganism to 15 kGy X rays irradiation (Billi *et al.*, 2000).

The authors hypothesized that the capacity of *Chroococcidiopsis* cells to repair radiation-induced DNA damage and probably also desiccation-induced DNA damage is related to the ability use redundant genetic information, as *D. radiodurans* does.

Other investigation have established that a vacuum of 10^{-6} Pa decreased the survival of desiccation-resistant microorganisms, and that exposure to space vacuum caused a greater decline in survival when compared to ground vacuum. The exposure to space vacuum ($\sim 10^{-6}$ Pa) decreased cell survival by two and four orders of magnitude for *Bacillus* sp. PD3D and *D. radiodurans* respectively (Saffary *et al.*, 2002).

Another interesting experimental work was performed with measurement of metals inside cells of radio-resistant bacteria (Ghosal *et al.*, 2005). Researchers had shown that radioresistant bacteria accumulate exceptionally high intracellular manganese and low iron levels. The opposite effect was found for dissimilatory metal-reducing *Shewanella oneidensis* that accumulates Fe but not Mn, and is extremely sensitive to radiation. The authors proposed that for Fe-rich/Mn-poor cells, death by radiation is caused by released Fe(II) from proteins, that leads to additional cellular damage by Fe(II)-dependent oxidative stress. In contrast, Mn(II) ions accumulated in cells of *D. radiodurans* might have function of antioxidants that reinforce enzymic systems, which defend against oxidative stress during recovery. The conclusion made is: It is necessary to include considerations of respiration, tricarboxylic acid cycle activity, peptide transport and metal reduction as potential targets to control recovery after radiation injury.

The effect of UV-B radiation on aquatic ecosystems was comprehensively discussed in Häder's review (Häder, 2000). It was concluded that the consequences of significantly enhanced levels of exposure to UV-B radiation include losses of biomass, such as food sources for human, changes in species composition, decrease in availability of nitrogen compounds and reduced sink capacity for atmospheric carbon dioxide, resulting in the potential augmentation of global warming. According to the author, there is significant evidence that increased UV-B exposure is harmful to aquatic organisms, however damage to ecosystems is still uncertain. Responses to increased solar radiation will not be limited to simple decreases in primary production. In fact, shifts in community structure may be more common and result in little detectable differences in ecosystem biomass.

The number of species in the genus *Deinococcus* was recently increased by additional 9 species isolated from the Sonoran Desert in Arizona (Rainey *et al.*, 2005), and currently this genus includes 20 species. The genera *Acinetobacter*, *Hymenobacter*, *Kineococcus*, *Kocuria*, *Methylobacterium* (eubacteria), and *Thermococcus*, *Pyrococcus* (euryarchaeotae) also include species and strains that are resistant to ionizing radiation.

8. Extremophiles in soils.

The diversity of extremophilic organisms in soils is quite high, exceeding our knowledge and what we once expected. This field is just beginning to be explored and appreciated. With the advancement in molecular techniques and the increasing availability of environmental rDNA sequences, future research can be done to look into the community structure as well as the microbial processes in which these microbes are engaged.

Microorganisms were known as the most abundant species in soil for some time, and their diversity was enormous. The paper published by Torsvik *et al.* (1990) indicated there were 4,600 distinct prokaryotic genomes per gram of soil based on DNA reassociation kinetics. A decade later,

they surveyed different types of soil and estimated 10^{10} cell/cm³ with up to 8,800 genome equivalents in pasture soil (Torsvik, 2002). The effort to study the composition and distribution, however, has only met with limited success. This is mainly due to the difficulties in cultivating and characterizing soil microbes by traditional, culture-based methods. As a result, cultivable species only represented less than 0.1% of the diversity (Bintrim et al., 1997).

In recent years significant progress has been made to assess microbial communities based on a variety of culture-independent methods and techniques that were developed. A number of studies utilized ribosomal RNA operon, typically the small subunits such as 16S and sometimes 5S rRNA, to analyze the population in soils (Stackebrandt, 1993; Ueda, 1995; Hofle, 1999; Borneman, 1997; Zhou, 1997). Environmental sequencing projects targeting 16S rRNA also allowed look at the diversity of uncultured organisms (DeLong and Pace, 2001; Tringe *et al.*, 2005). Several techniques based on PCR amplification of DNA extracted from soil samples also became available. These include ribosomal intergenic spacer analysis (RISA), denaturing gradient gel electrophoresis (DGGE), temperature gradient gel electrophoresis (TGGE), single-strand conformation polymorphism (SSCP), random amplified polymorphic DNA (RAPD) and amplified ribosomal DNA restriction analysis (ARDRA). The fingerprints generated from these methods enable the investigation of community compositions as well as study the differences due to treatment or biogeographic changes (Kent, 2002; Fierer, 2006).

The PCR-based methods are generally easy to perform and they have made significant contribution in soil community analysis. In spite of their popularity and frequent usage, these methods are not without limitations. The procedures often introduce variables, which result in biases in detecting certain populations. Some limitations include cell lysing methods possibly affect DNA recovery from certain bacteria, extracting DNA from soil samples not always capturing microbes in low numbers, and various inhibitors in soil impacting the amplifying capability of PCR.

In order to circumvent and minimize biases, different approaches have developed in recent years to improve detection and resolve amplified DNA fragments. These include length heterogeneity PCR (LH-PCR) (Ritchie, 2000) and terminal restriction fragment length polymorphism (T-RFLP) (Kitts 2001; Marsh, 2000; Osborne, 2000). Alternatives to PCR methods have also been used; examples include Fatty Acid Methyl Ester (FAME) profiles of cell membrane (Ibekwe, 1998) and fluorescence in situ hybridization (FISH) to perform direct microscopic examination (Amann 2001). The latter technique also allows specific groups of interest to be targeted by customizing the fluorescent probes for hybridization (Amann, 1995; Raaijmakers, 1998).

Microbial diversity in soils.

Many environmental factors influence prokaryotic diversity in soil. These include: soil moisture, particle size, pH, organic carbon concentration, carbon to nitrogen ratio, and vegetation type. Fierer & Jackson's study (2006) indicated bacterial diversity was unrelated to site temperature, latitude and other variables that typically affect plant and animal diversity, and the composition of a community was not dependent on geographic distances. Instead, the pH of soil was the best predictor of bacterial diversity and richness, as well as community composition. Bacterial diversity was highest in neutral soil, and the level of diversity and richness decrease as pH moves towards acidic conditions. The authors did not know whether this phenomenon extended to the other extreme, because basic soil (pH>8.5) samples were not easy to obtain. So it is unclear if diversity plateaus in soils with neutral pH.

Microbial diversity is influenced by the particle size of soil (Ranjard, 2000). One study indicated α -Proteobacteria dominated in large particle soil, whereas *Holophaga/Acidobacterium* were the most common organisms found in clay particles (Sessitch, 2001).

There were many studies showing bacterial diversity, or community composition, from different parts of the world (Liles et al., 2003; Nemergut et al., 2005; Wery et al., 2003; Nagy et al., 2005). Liles et al. (2003) analyzed the diversity of microbial genomes by constructing a metagenomic DNA library from soil samples collected in Madison Wisconsin. The rDNA genes indicated the majority of the bacteria belonged to the Proteobacteria, Acidobacteria, Cytophaga-Flexibacter-Bacteroides (CFB), Firmicutes and Verrucomicrobia groups (76-86 % within a three-year span). The predominant population belonged to members in Acidobacteria. Within the Proteobacteria, all subdivisions of alpha, beta, gamma and delta-Proteobacteria were present. Their proportions varied depending on the molecular approaches being used. The bacterial sequences obtained from a sandy ecosystem near Darmstadt, Germany showed similar diversity as to the phyla and groups being detected (Ochsenreiter, et al., 2003).

Acidobacteria was the predominant group in the Colorado Rocky Mountains alpine tundra soil (Nemergut et al., 2005). They were most abundant during spring, summer and winter months; and they also presented in high numbers in both wet and dry meadow samples during springtime. Another study which surveyed the prokaryotic diversity in Arizona Sonoran Desert soil crusts showed Acidobacteria-like clones ranked highest in arid soils, and their DNA was among one of the major groups being amplified and identified. On the other hand, these bacteria were low representatives in agriculture soils (Nagy et al., 2005). These findings were consistent with those of Dunbar et al. (1999) where *Acidobacterium*-related organisms comprised nearly half of the identifiable bacteria in their clone libraries from four arid soils in Northern Arizona. Furthermore, a wheat field from Holland also yielded similar results, showing *Acidobacterium* was one of the dominant groups of uncultured organisms (Smit et al., 2001).

Acidobacteria are considered extremophiles, capable of growing under unusual conditions when compared with other soil inhabitants such as Proteobacteria, Actinobacteria, Firmicutes. The type species, *Acidobacterium capsulatum*, was an acidophilic chemoorganotroph isolated from acidic mineral environment. Since its name was validated in 1991, members of this genus have been found in a variety of soils as mentioned above, and not only limited to acidic habitats.

Other extremophilic organisms existed in soil, especially if the environmental conditions of the soil are considered extreme; i.e. Antarctic soil, alkaline sediments, acidic forest cambisol (Wery, et al., 2003; Dunfield et al., 2003; Wiegel and Kevbrin, 2004). Many of these extremophiles belong to the Bacteria Domain. It is not surprising to find a number of psychrotrophs in Antarctic soils; for example, species of *Psychrobacter* and *Planococcus* as well as other ubiquitous genera (i.e. *Chryseobacterium*, *Brevundimonas*, *Paenibacillus*) that could tolerate low temperatures (Wery, 2003). Then there are halophilic organisms residing in saline soils; *Marinococcus halotolerans* (Li et al., 2005), *Streptomonospora alba* (Li et al., 2003), *Alkalibacterium iburiense* (Nakajima et al., 2005) just to name a few from recently published names. Thermophilic organisms continued to be discovered in soils with higher temperature; for example, *Anaerobranca gottschalkii* is a novel thermoalkaliphile isolated from a humid soil of a hot lake inlet in Kenya (Prowe & Antranikian, 2001), and *Geobacillus tepidamans* originated from a geothermally heated soil in Yellowstone National Park (Schäffer et al., 2004). Examples in acidic niches are *Methylocella silverstris*, a

methanotroph from an acidic forest cambisol (Dunfield et al., 2003), and two Clostridia *Cl. akagii* and *Cl. acidisoli* from acidic peat-bog soil (Kuhner et al., 2000).

Then there existed some extremophiles in unexpected places. Echigo et al. (2005) reported halophilic bacteria inhabiting in non-saline environment in areas around Tokyo, and most of them belong to the *Bacillus* and related genera. A couple of thermophilic spore formers were isolated from cool soil environment in Northern Ireland (Banat et al., 2004). Alkaliphiles have been isolated from a variety of mesobiotic and neutral soils and sediments (Wiegel and Kevbrin, 2004). Several studies have found sequences from mesophilic soil habitats worldwide, and they belong to the Crenarchaeota phylum in Archaea Domain by phylogenetic association (Sliwinski and Goodman, 2004; Simon et al., 2005; Ueda, 1995).

Archaea in Soils.

Crenarchaeota. The presence and significance of Archaea in soils were not recognized until a decade ago. This was mainly due to the fact that Archaeal members were mostly isolated from extreme environments or unique ecological niches, and they could not be easily cultivated and studied by traditional microbiological methods. In 1997, Bintrim with colleagues demonstrated the presence of archaeal population in soil, and they were more widespread than once realized (Bintrim et al., 1997). By amplifying and cloning the small subunit ribosomal RNA genes of Archaea from soil samples collected in Madison Wisconsin, a cluster of these organisms was found to represent a component of the microbial communities in these soils. Phylogenetic analysis of these clone sequences indicated this lineage was within the Crenarchaeota phylum, but its branch was distant to any cultivated member of the Archaea. In fact, it affiliated with the Group I marine Archaea, which composed of members in the planktonic division of the Crenarchaeota.

Subsequent studies confirmed this unexpected finding, that nonthermophilic members of Crenarchaeota were ubiquitous and abundant in soil. This particular lineage of Crenarchaeota was consistently present in the different soils samples from diverse geographic areas in Europe and Asia, as well as from two microbial mats (Ochsenreiter, et al., 2003). A survey from soil samples in Darmstadt, Germany revealed the relative abundance of crenarchaeotal rDNA was between 0.5-3% in bulk soil. In the rhizosphere of a sandy ecosystem and an agricultural setting, these archaeal sequences represented 0.16 and 0.17 % of the populations respectively. Thus, soil crenarchaeota constitute a stable and unique component of the inhabitants in terrestrial environments (Ochsenreiter, et al., 2003).

Furthermore, these crenarchaeota were found in association with plants. Their small subunit rRNA gene sequences were present in rice rhizosphere (Großkopf et al., 1998b), plant roots such as maize and tomato (Chelius and Triplett, 2001; Simon et al., 2005), and mycorrhizospheres of pine seeding (Bomberg et al., 2003). Using epifluorescence microscope and culture-independent recovery of small subunit rRNA gene sequences, Simon et al. (2000) showed soil Crenarchaeota colonized both young and senescent plant roots at unexpected high frequency. In fact, they were particularly good colonizers of senescent roots, thus, these organisms had important implications for root development and ecology.

Sliwinski & Goodman used PCR-single stranded conformation polymorphism (PCR-SSCP) to compare the crenarchaeal consortia inhabiting the rhizosphere with those present in bulk soil at different locations in Wisconsin. They showed the profiles from rhizosphere were distinct from those obtained in corresponding bulk soil. This difference seemed to be resulting from the increased richness in rhizosphere as compared to bulk soil, and it was not dependent on plant lineage.

Therefore, the authors concluded that the crenarchaeal associations with plants in native environments reflected the interactions between plant and microbes, and their relationship was mediated by environmental factors (Sliwinski & Goodman, 2004).

Archaea also reside in upland pastures, as shown by Nicol *et al.* (2004). Managed pastures with inorganic nitrogen fertilizer input had a different crenarchaeal community than those from unmanaged upland pastures (i.e. native grassland) where their nitrogen sources were largely from grazing sheep. The archaeal communities were very stable in these soil microcosms, and they were not affected by nitrogen sources amendments and pH increases.

All these results indicate mesophilic soil crenarchaeotes play an important ecological role (Bintrim *et al.*, 1997) and their association with plant roots is significant. Until recently, all the studies were done by culture-independent methods and no member has been isolated and identified. Some progress began to surface lately, as the first evidence of culturing mesophilic crenarchaeotes was reported. These organisms responded positively to root extract enrichment although the actual substrates supporting their growth were still unknown (Simon *et al.*, 2005). As novel cultivation methods continue to be developed, the axenic culture of crenarchaeotes will hopefully appear in the near future.

Euryarchaeota. Other archaeal diversity has been found in different types of soil in addition to Crenarchaeota. By using 16S rRNA PCR amplification, Utsumi *et al.* (2003) surveyed peat soil samples from geographically distinct wetlands, ranging from the cold West Siberia bog to the subtropical Okefenokee swamp in the US. Most of the sequences were related to the methanogens in the Euryarchaeota phylum. This finding was not surprising, since methanogens were among the earliest members recognized in the Archaea Domain. Many genera and species have been described since, in spite of their obligate anaerobic characteristics (Euzéby, 2006).

Most of the archaeal community studies on Euryarchaeota were done in paddy field soil, especially from flooded rice paddies where these organisms exist as one of the major biogenic sources of atmospheric methane. Methanogens are the most functionally important archaea in these environments due to the anoxic condition caused by flooding (Liesack *et al.*, 2000). These organisms utilize H₂-CO₂ or acetate as substrates for methanogenesis, and they can survive during the dry and oxic periods. The majority of methanogens are found in the anoxic bulk soil of flooded rice microcosms; in fact, between 10⁶ to 10⁷ cells of H₂-CO₂ utilizing methanogens were detected per gram of dry soil in Italy (Großkopf *et al.*, 1998a). Reports on Italian and Japanese rice paddy soils seem to indicate *Methansarcinaceae* (i.e. species *M. barkeri*, *M. mazei*) and *Methanosaetaceae* (species *M. concilii*) were the predominant acetate-utilizing methanogens; whereas members of the *Methanobacteriaceae* family (species *M. bryantii*, *M. formicicum*) were the major group found in hydrogenotrophic methanogens (Liesack *et al.*, 2000; Großkopf *et al.*, 1998a; Min *et al.*, 1997). These diverse patterns of methanogenic organisms were influenced by environmental factors such as acetate concentration, temperature, and soil particles size (Liesack *et al.*, 2000; Großkopf *et al.*, 1998a; Chin *et al.*, 1999; Ramakrishnan *et al.*, 2000). In addition to the known methanogenic species, Liesack *et al.* (2000) noticed several novel archaeal lineages that have been detected in the anoxic bulk soil, rice roots of flooded rice microcosms, and in slurries of Italian rice paddy soil. Pure cultures were not available from these lineages and only a phylogenetic branching pattern was noted. One of the rice clusters grouped on the same branch as *Thermoplasma acidophilum* and marine group II, but it only had a distant relationship. The other rice cluster formed a novel main line of descent without any affiliation to any other euryarchaeal groups (Großkopf *et al.*, 1998b). Obviously

there are members in the Euryarchaeota phylum waiting to be discovered in order to study and elucidate their ecological roles.

9. Astrobiological significance of extremophilic microorganisms.

The study of metabolically active microorganisms at low and/or subzero temperatures is of major interest to astrobiology, since most bodies of our Solar System are frozen worlds.

It has also been shown that microorganisms can remain viable for geological periods of time when cryopreserved in ice (Abyzov *et al.*, 2004; Pikuta *et al.*, 2005). It is well known that microbes are generally capable of remaining viable for very long periods of time when freeze-dried (lyophilization) and that some organisms can maintain metabolic activity at temperatures as low as -20 °C. Representatives of the genus *Trichococcus* have been found to be capable of replication and growth at -5 °C at both aerobic and anaerobic conditions (Pikuta *et al.*, 2006).

The microflora of the cryosphere of planet Earth provides the best analogs for life forms that might be found in the permafrost or polar ice caps of Mars, near the surface of the nuclei of comets, or in the liquid water and the ice crusts of icy moons of Jupiter and Saturn (Hoover *et al.*, 2002; Hoover and Pikuta, 2004; Abyzov *et al.*, 2006). It is now well established that water ice exists today in a vast deposit at the North Polar Ice Cap of Mars. (Zuber *et al.*, 1998; Bass, and Paige, 2000; Bass *et al.*, 2000; Kieffer *et al.*, 1976; Greve 2000; Haberle, and Jakosky, 1990). The Mars Global Surveyor had produced images of double-rimmed polygons on Mars that are consistent with the double-rimmed polygons in permafrost of Earth that are produced by expansion and contraction effects associated with the seasonal water-ice freeze-thaw cycles (Paepe *et al.*, 2001). The High Energy Neutron detector aboard the Mars Odyssey spacecraft (Litvak *et al.*, 2006) has also found that there are large water-rich permafrost areas, with contents up to 50 % water by mass fraction, at both the North and South polar regions of Mars. In the northern latitudes, these water-ice layers of permafrost are very near the surface, but in the south there are covered by a thicker (10-20 cm) layer of soil. The water content of the Mars regolith was found to exceed 11% at latitudes greater than 60 degrees in both hemispheres. Two regions near the equator (Arabia Terra and Medusae Fossae) even exhibited 9-10 % water (Mitrofanov *et al.*, 2004) with the lower layer of soil in the moistest region (30 E, 10 N) having soil with over 16% of water by mass which is comparable to the water content (13-19 %) of tropical rain forest soil on Earth (Odokuma and Dickson, 2003).

The possibility that microbial extremophiles might thrive today (or be present in a cryopreserved state) in the water ice and permafrost in the modern Martian regolith is of great significance to Astrobiology. While it is generally accepted that the one common bond for all life known on Earth is the universal requirement for water, it is clearly not necessary that the water be found in liquid state in large bodies, such as seas or lakes. Many microorganisms grow within thin films of water in permafrost and in frozen rocks of the polar regions (cryptoendoliths). Microorganisms also inhabit acidic (acidophiles) and brine channels (halophiles) in glaciers (Hoover *et al.*, 2002; Pikuta and Hoover, 2004b; Hoover and Gilichinsky, 2001; Mancinelli *et al.*, 2004). On Earth, solar heating of low albedo rocks entrained in glaciers results in microenvironments (cryoconite ecosystems) with trapped liquid water, gasses, organic chemicals, and minerals that provide ideal conditions for the growth of microbial communities that include both prokaryotic and eukaryotic microorganisms. The study of the species of Earth's microflora that thrive in the coldest regions of our planet is of great importance to the design and development of

robotic systems that will be needed to search for biomarkers and evidence of past or present life on Mars, Europa, and other prime targets for Astrobiology research.

The importance of studying acidophilic microorganisms is primarily concerned with acidic thermal systems such as may exist on bodies of our solar system that have volcanic or tectonic activities. Evidence for ancient volcanic activity on Mars and Venus, and recent volcanic activity on Io has been revealed by the *Viking*, *Pathfinder*, *Global Surveyor*, *Mars Odyssey*, *Voyager*, *Venera*, and *Magellan* space missions with optical, ultraviolet, infrared, and radar images and IR, gamma-ray, and neutron spectral data (Pikuta and Hoover, 2004b). Much volcanic activity has also occurred on Venus, which is known as the most acidic planet of the Solar System. Venus is thought of as a twin planet of Earth although it rotates in the opposite direction. However it is much closer to the Sun and it has a surface temperature of around 470 °C, and an atmospheric pressure is equivalent 100 times that of Earth's atmosphere. After the discovery (Harris *et al.*, 2002) of active microorganisms in Earth's atmosphere at an altitude of 44 km (level of stratosphere) the point of view about possible existence of life forms on acidic droplets in the cooler dense clouds of the Venus atmosphere has been discussed.

The Jovian moon Io presents the most active volcanic body of our Solar System with conditions that may be acceptable for the development of sulfur-dependent microorganisms. Io's volcanic activity could also provide acidic geysers and springs suitable for the development of acidophilic sulfur-metabolizing lithotrophs. There is also much evidence that beneath the fractured ice crust of Europa, the liquid water ocean exists. The Jovian tidal effects and/or hydrothermal vents could provide the heat necessary for the liquid water ocean underneath the crust of this icy moon. It is conceivable that acidophilic microorganisms might also inhabit acidic hydrothermal vents on Europa (Pikuta and Hoover, 2004a), and that psychrophiles could thrive in niches at interfaces between the liquid water and the ice sheet.

The importance of study alkaliphilic microorganisms for astrobiology was enhanced by the findings of abundant carbonates and carbonate globules rimmed with possibly biogenic magnetites in association with the putative microfossils in the ALH84001 meteorite (McKay *et al.*, 1996; Wadhwa and Lugmair, 1996). The study of this meteorite provided definitive evidence of the existence of carbonate minerals on Mars. The participation of microbial agents in formation of carbonate systems on Earth was discussed previously (Pikuta *et al.*, 2003a). Even though the ALH84001 nano-structures were not of sufficient size as to be certainly accepted as valid recognized microfossils, these results profoundly stimulated the study of meteorites, bacterial paleontology, and microbial extremophiles. The extensive research efforts triggered by the ALH84001 results have helped to define valid biomarkers and to delineate where and how to best search for evidence of life in Earth's most hostile environments and elsewhere in the Solar System.

Recent studies have resulted in the detection of evidence for the mineralized remains of a large and complex assemblage of diverse coccoidal and filamentous microorganisms and bacterial mats in the Orgueil (CI1) and Murchison (CM2) carbonaceous meteorites (Hoover, 2005; Hoover 2006). These forms are much larger and more complex than the ALH84001 nanostructures and they exhibit features that provide clear associations with known morphotypes of all five orders of the Cyanobacteriaceae. The forms occur as spherical and irregular coccoids, sometimes in colonies with electron transparent carbonaceous mucilaginous envelopes. These morphotypes have characteristics of the Chroococcales and in pseudo-filaments with baeocyte-like forms consistent with the Pleurocapsales. These morphological simple forms are not so distinctive as the larger forms that

occur as uniseriate and multiseriate unbranched and branched filaments and mats that have been found embedded in the meteorite rock matrix. The vast majority of the filamentous forms found in freshly fractured surfaces of the meteorites are consistent in both size and detailed morphological features with the non-polarized filaments, carbon-rich sheaths, and mineralized trichomes of the Oscillatorian cyanobacteria. However, the Orgueil meteorite has also been found to contain a number of other forms that have tapered polarized filaments and exhibit cross-wall constrictions, true and false branching and highly differentiated cells. Many of the tapered filaments appear to represent benthic forms attached to the substratum by basal heterocysts (occasionally exhibiting ellipsoidal akinetes) with detailed features characteristic of morphotypes of the genus *Calothrix*. The most distinctive of the Orgueil filaments are exceptionally well-preserved forms that are interpreted as representing morphotypes of the Orders: Nostocales and Stigonomatales. These microfossils have extremely complex and distinctive biogenic characteristics that cannot be confused with minerals and other abiotic microstructures. Energy Dispersive X-ray Analysis has revealed that they have anomalous C/O, C/N, and C/S ratios that clearly establish that they cannot be attributed to recent biological contaminants and the forms have been interpreted as indigenous microfossils. The detection of microfossils in carbonaceous meteorites supports the hypothesis proposed by Hoyle and Wickramasinghe in 1981 that comets and meteorites could be carriers and distributors of living cells in the Solar System (Hoyle and Wickramasinghe, 1981; Hoover *et al.*, 1986).

Microbial extremophiles investigated for Astrobiology research.

Based on everything discussed, there are important reasons to study extremophilic microorganisms as part of Astrobiological research in our laboratory, and the following directions were chosen for investigation: 1) Study of samples from cold regions of our planet (Alaska, Siberia, Patagonia, Antarctica); 2) Study of samples from acidic ecosystems (Chena hot spring in Alaska); 3) Study of samples from alkaline ecosystems (soda lakes in California); 4) Study of samples from deep-sea hydrothermal vents (Azores, Middle Atlantic); 5) Study of samples from moderate thermal springs (Alaska).

As a result the following groups of microorganisms were isolated and studied. From halo-alkaline systems of Owens and Mono Lakes in California five strains of alkaliphilic mesophilic bacteria were isolated in pure cultures and described. Four of them have been validly published and deposited in International Microbial Culture Collections:

- 1) *Spirochaeta americana* ASpG1^T was found to be the first obligately anaerobic and alkaliphilic free-living sugarlytic spirochete on the American continent; the peculiar feature of this unique strain is production of molecular hydrogen as a major end metabolic product that remains biotechnologically potential for the alternative fuel strategy in the economy;
- 2) *Desulfonatronum thiodismutans* FML1^T was also found to be the first obligately alkaliphilic sulfate-reducing bacterial species on the American continent, and as a specific feature of this strain differentiating it from Asian strains is that it has a chemolithotrophic metabolism that allows this bacterium to grow exclusively on hydrogen and CO₂ without any organic molecules;
- 3) *Tindallia californiensis* APO^T was described as an agent responsible for decomposition of proteolysis products and some organic acids (Pikuta *et al.*, 2003d); This species performs the function of a secondary anaerobe in anaerobic alkaliphilic microbial

community of Mono Lake; This strain is capable of respiration by performing a Stickland reaction on certain pairs of amino acids;

4) *Anaerovirgula multivorans* SCA^T was found as the cellulolytic agent in the anaerobic microbial community of Owens Lake, but sequent study showed this process as very weak. The major trophic function for this species is degradation of proteolysis products molecules and some organic acids. This strain was found to be phylogenetically distant from known clostridial species and had taken a separate lineage on the separate genus level and separate species (Pikuta et al., 2006b);

5) *Spirochaeta* sp. ASpC2^T is now in preparation for publication; this obligately anaerobic and alkaliphilic strain was isolated from a cellulolytic enrichment culture from Owens Lake, and it has a number of physiological features differentiating it from other known free-living spirochetes.

From the acidic system of Chena Hot Spring in Alaska the obligately acidophilic strain AGC2^T was isolated in pure culture (Pikuta and Hoover, 2004a). This mesophilic, spore-forming strain is able to grow on the medium with high concentrations of Fe²⁺ with pH 1.5-2.0. Currently the article with taxonomic description of this strain is in preparation.

From a deep-sea hydrothermal vent the obligately sulfur-reducing archaeobacterial strain OGL-20P^T was isolated and described. This isolate belongs to a separate new species of the genus *Thermococcus* with proposed name *T. thioeducens* (*IJSEM* in peer-review).

The study of samples from cold regions led to discovery of psychrotolerant *Carnobacterium pleistocenium* FTR1^T that was alive after 32,000 years permafrost conditions in Fox Tunnel in Alaska (sample was dated as Pleistocene's geological period). This facultatively anaerobic fermenting bacterium was described as a separate new species since it was different from known species both on genetic and phenotypic levels (Pikuta et al., 2005). This bacterium is of great importance to both Astrobiology and Paleontology. It is first a validly published species of the Pleistocene period and it was cryopreserved in ancient permafrost. It is also important to cryobiology by demonstrating that it is possible to extract living microorganisms from the ice and permafrost samples of other bodies of the Solar System.

Another interesting psychrotolerant sugarlytic bacterium *Trichococcus patagoniensis* PmagG1^T was isolated from penguin guano, and it was able to grow at -5 °C at aerobic and anaerobic conditions. This is the lowest known record for growth of pure culture (Pikuta et al., 2006a). The description of the proteolytic psychrotolerant bacterium strain PPP2^T that represents a separate new genus and species is also in preparation (source of isolation is also penguin guano).

Conclusion.

If we consider the distribution of life in a matrix of pH/Temperature, salinity/Temperature, salinity/pH, radiation/Temperature by placing the names described known species of microorganisms on the corresponding matrix, the picture will be as follows. The diagram pH/Temperature (Fig. 1a) shows a gray painted square in an area of neutral pH and moderate/moderate-thermophiles limits; this square represents the region of a majority of known described microbial species that includes saprophytes, pathogens, and environmental important agents responsible for the balance of trophic chains in surrounding ecosystems. On the left side of this square the distribution of acidophilic species is shown. As we can see, none of the acidophilic

psychrophilic species were described and found, this supports the point of view about primary origin of acidophiles on hot cooling Earth. The area of hyperthermophilic and moderately thermophilic species of acidophiles is quite busy and is mostly "populated" by a species of archaea. The most thermostable species is *Pyrolobus-fumarii* that could survive 30-60 min in autoclave at 120 °C, and the neutrophilic but not validly published strain 121 that grows at a temperature higher than 120 °C and survives for a short period of time at 130 °C. The most alkalitolerant described species of archaea among the hyperthermophiles is *Thermococcus alkaliphilus*, which is able to grow at pH 10.5. On the right side of the gray painted square the distribution of mesophilic and moderately thermophilic species of microorganisms is shown; it includes bacteria, cyanobacteria, and archaeobacteria. As was discussed previously, mesophilic cyanobacteria are the leaders in alkalitolerancy among all living organisms. The recently described psychrotolerant species *Trichococcus patagoniensis* is able to grow at pH 10.0 and its other specific feature is the ability to grow at -5 °C. Before the discovery of this species the record of growth at subzero temperatures was established for psychrophilic species of sulfate-reducing bacteria

The diagram of salinity/pH (Fig.1b) shows the distribution of known species of microorganisms within this matrix. On the diagram on the left side the gray painted oval represents the area of a majority of known species of microorganisms (saprophytes, pathogens, and environmental agents). As we can see, the extreme acidophiles appear in regions of salinity before 5 %. Species of the genera *Natronobacterium* and *Natronorubrum* are able to grow at 30 % (w/v) salinity. Cyanobacteria are spread to the pH 12 area and limited by 15 % salinity. This diagram, once again, confirms the hypothesis about the later appearance of halophily during the evolution of life. The accumulation and solubility of Na⁺ and Cl⁻ ions in water became possible only on latest stages of Earth evolution.

The diagram of salinity/Temperature (Fig. 1c) shows the presence of a mesophilic and moderately thermophilic species of microorganisms in extreme saline area (25-30 %); None of the hyperthermophilic species are known at extreme salinity. Probably, again, the rule of Earth's life evolution is working: the first organisms were non-halophilic hyperthermophiles; the primordial ocean was not salty at all.

And the final diagram (fig.1d) shows the distribution of a known species in matrix of radiation/temperature. Three species (one of them an archaeobacterium) could tolerate 30 kGy irradiation level. Does that suggest that the level of irradiation on Early Earth was around this number? Or is it possible that these microorganisms originated elsewhere in the cosmos and were transported by comets and meteorites to Earth? To resolve these questions there is a need for Science to have more data (this area of extremophily has not been thoroughly studied).

Looking at these diagrams the logical question raises: What direction of biological changes (adaptation, mutations, transformation into metabolically inactive forms, etc.) to cells had been experienced during the Earth evolution? Is it appropriate to suggest that an exclusively single source of homogenous cell culture started biological evolution on Earth, or were there several physiologically different types of cell cultures that interacted and developed into present day life? So, who was the first ancestor from the whole diverse extremophilic microbial world that we observe today? If the formation of any planet required a certain high-temperature regime then it is obvious, the hyperthermophilic microorganisms have priority as the role of ancestors. Psychrophily would have to have developed after a significant temperature drop on the planet surface. Free oxygen at the formation and cooling of the planet is excluded, and this means that anaerobic types of metabolisms

would have been primary. Acidophily probably was the initial in biological history on Earth, and only after certain mineral precipitation and enough buffer concentration of CO₂ in the air the occurrence of alkaliphily became possible. Halophily perhaps developed as the latest when the arid climate dominated on the land parts of Earth. Due to the tiny sizes of bacterial and archaean cells, the pressure does not have a significant influence on them as it has for the highly organized forms of life with macroscopic sizes, but still the barophily was formed among microorganisms of deep-sea and deep-underground ecosystems. Another fundamental question in biology arises: How is life distributed after the delivery transportation or after self-origin on the planet? Was it from the surface to center, or opposite, from the center (initial requirement of high pressure and absence of light) to a surface?

The sequence and direction of mutations that allowed adaptations to occur in ancestor cells in response to the new changes of environments had a complicated character. It included not only multi-factor influence, but probably, back and forward vectors as well. During the process of evolution, biological cells could have experienced several extreme physico-chemical factors at the same time. As a result, newly formed organisms appeared that were resistant, tolerant, or even developed obligate requirement (phily) to the multiple aggressive environmental factors. This is supported by the fact that now we observe the existence of acidophilic thermophiles, halo-alkaliphilic thermophiles, barophilic hyperthermophiles, halo- or alkaliphilic psychrophiles, etc. The genetics of extremophilic microorganisms could be read as the map of environmental effects in the past and probably have enough flexibility for providing a necessary "buffer pool" towards the future stressful changes. In any case, life would easily become extinct at any global cataclysms on climatic and geophysical levels. Except for the environmental influence of the ecosystems affecting living cells, another very important factor was developed and designated as *biological*. Life itself even on the level of monotypic or pure culture always has competitive relations between cells: those that eat more and breath better, are the strongest and predominant, and the weaker forms die out. On the interspecies level this is expressed even more effectively: the specific behavior and physiological specifics such as an excretion of antibiotics, poisons, or an inhibition by trivial end metabolic products, all of these quite limit the neighborhood of surrounding species, that led to the creation of the adjusted functionally and physiologically microbial consortia, and then communities. Symbiotic relations between different kinds of microbial organisms are reflected on the cytological and genetic levels of eukaryotic organisms (mitochondria and chloroplasts) for energetic efficiency, and also in such phenomenon as the binary microbial cultures (cases when microbiologists cannot separate cells of two symbiotically (undetected trace compounds or an exchange by vitamins) related microbial cultures. It is quite possible that such relationships initially have also been developed in extreme environments.

So, the interaction of growing and accumulating organisms with the environment and other biological species consequently led to the creation of balanced ecosystems. All geo-paleontological deposits represent an encoded chronology of biological activity. The formation of minerals and fossils is the direct or indirect result of biological activity. Fossilized carbon, oil, elemental sulfur, methane, gold, iron pyrites, etc., all of them were formed in certain periods of evolution under specific circumstances, where local or wide-spread regions had experienced the demographic explosion of certain extremophilic populations, and the formation of rocks and minerals in such regions occurred due to the bioleaching processes or just to pure metabolic activity. After the global climatic changes and the geo-formation processes some populations with unique physiology were

completely replaced by others forever, this can be observed on mineral and rock deposits. For this reason, all modern extremophiles could be appropriately called relics, because they do have genome that memorized environmental history, and this feature makes the study of extremophiles important to many fundamental sciences. Indeed, the modern face of the Earth is a product of the interaction of biological agents (mostly microorganisms) and minerals. Soil formation was the key moment for the distribution of plants on Earth, and it still remains an unsolved problem in biology. In this process, again the extremophilic microorganisms probably played a pioneering role. It is now well established that eukaryotic organisms cannot exist without prokaryotes; it is equally true for both plants (rhizosphere of root system, protective surface flora) and animals (flora of intestine and skin surfaces). The participation of extremophilic bacteria in such ecosystems is not excluded (bacterium developing at extreme acidity in stomach resulting in chronic disease, microbial process of methanogenesis in intestine, etc.).

So, what is the situation with the cultivation of extremophilic microorganisms today? The molecular biologists have pronounced that microbiologists have successfully cultivated *in vitro* only one percent (or less) of all microbial diversity that exists in nature. Based on the data of genomic libraries constructed by 16 S rRNA gene sequence analysis, approximately 99 % of whole gene pool of prokaryotes still remains unknown. At the present time around 7300 species of bacteria and archaea are validly published. From the time when Dr. Petri introduced the method of cultivation on agar solid media using dishes, which are now named for him, one hundred years have past, but hyperthermophilic anaerobes were discovered only in the past twenty five years or so. Of course, the percent of cultivated extremophilic microorganisms compared to other "normal" bacteria and archaea is significantly lower. The development of Biotechnology has tremendously stimulated the investigation of extremophiles. Today many laboratories in different countries are extensively engaged in the study of these fascinating and critically important microorganisms.

References

- Abd-el-Malek, Y., and S.G. Rizk. 1963. Bacterial sulphate reduction and the development of alkalinity. *J.Appl.Bacteriol.* 26: 7-26.
- Abe, F., and K. Horikoshi. 2001. The biotechnological potential of piezophiles. *Trends Biotechnol.* 19: 102-108.
- Abyzov, S.S., Hoover, R.B., Imura, S., Mitskevich, I.N., Naganuma, T., Poglazova, M.N., and M.V. Ivanov. 2004. Comparative Results of Using Different Methods for Discovery of Microorganisms in Very Ancient Layers of the Central Antarctic Glacier above Lake Vostok. *Adv.Space Res. Cospar.* 33: 1222-1230.
- Abyzov, S.S., Duxbury, N.S., Bobin, N.E., Fukuchi, M., Hoover, R.B., Kanda, H., Mitskevich, I.N., Mulyukin, A.L., Naganuma, T., Poglazova, M.N., and M.V. Ivanov. 2006. Super-long anabiosis of ancient microorganisms in ice and terrestrial models for development of methods to search for life on Mars, Europa and other planetary bodies. *Adv.Space Res. Cospar* 38: 1191-119.

Albuquerque, L., Simões, C., Nobre, M.F., Pino, N.M., Battista, J.R., Silva, M.T., Rainey, F.A., and M.S. da Costa. 2005. *Truepera radiovictrix* gen. nov., a new radiation resistant species and the proposal of *Trueperaceae* fam. nov. *FEMS Microbiol.Let.* 247: 161-169.

Amann, R.I. 1995. Fluorescently labeled, ribosomal RNA-targeted oligonucleotide probes in the study of microbial ecology. *Mol.Ecol.* 4:543-553.

Amann, R., Fuchs, B.M., and S. Behrens. 2001. The identification of microorganisms by fluorescence in situ hybridization. *Curr.Opin.Biotechnol.* 12:231-236.

Amend, J.P., and A. Teske. 2005. Expanding frontiers in deep subsurface microbiology. *Palaeogeography, Palaeoclimatology, Palaeoecology* 219: 131-155.

Anellis, A., Berkowitz, D., and D. Kemper. 1973. Comparative resistance of nonsporogenic bacteria to low-temperature gamma irradiation. *Appl.Microbiol.* 25: 517-523.

Banat, I.M., Marchant, R., and Rahman, T. 2004. *Geobacillus debilis* sp. nov., a novel obligately thermophilic bacterium isolated from a cool soil environment, and reassignment of *Bacillus pallidus* to *Geobacillus pallidus* comb. nov. *Int.J.Syst.Evol.Microbiol.* 54: 2197-2201.

Barns, S.M., Fundyga, R.E., Jeffries, M.W., and N.R. Pace. 1994. Remarkable archaeal diversity detected in a Yellowstone National Park hot spring environment. *Proc.Natl.Acad.Sci.USA.* 91: 1609-1613.

Bartlett, D.H. 2002. Pressure effect on in vivo microbial processes. *Biochem.Biophys.Acta* 1595: 367-381.

Bass, D.S., and D.A. Paige. 2000. Variability of Mars' north polar water ice cap. II. Analysis of Viking IRTM and MAWD data. *Icarus* 144: 397-409.

Bass, D.S., Herkenhoff, K.E., and D.A. Paige. 2000. Variability of Mars' north polar R. water ice cap. I. Analysis of Mariner 9 and Viking orbiter imaging data. *Icarus* 144: 382-396.

Beeder, J., Nilsen, R. K., Rosnes, J. T., Torsvik, T., and T. Lien. 1994. *Archaeoglobus fulgidus* isolated from hot north sea oil field waters. *Appl.Environ.Microbiol.* 60: 1227-1231.

Beeder, J., Torsvik, T., and T. Lien. 1995. *Thermodesulforhabdus norvegicus* gen., sp. nov., a novel thermophilic sulfate-reducing bacterium from oil field water. *Arch.Microbiol.* 164: 331-336.

Billi, D., Friedmann, E.I., Hofer, K.G., Caiola, M.G., and R. Ocampo-Friedmann. 2000. Ionizing-radiation resistance in the desiccation-tolerant cyanobacterium *Chroococcidiopsis*. *Appl.Env.Microbiol.* 66: 1489-1492.

- Bintrim, S.B., Donohue, T.J., Handelsman, J., Roberts, G.P., and R.M. Goodman. 1997. Molecular phylogeny of *Archaea* from soil. *Proc.Natl.Acad.Sci.* 94: 277-282.
- Blöechl, E., Rachel, R., Burggraf, S., Hafenbradl, D., Jannasch, H.W., and K.O. Stetter. 1997. *Pyrolobus fumarii*, gen. and sp. nov., represents a novel group of archaea, extending the upper temperature limit for life to 113 °C. *Extremophiles* 1: 14-21.
- Bomberg, M., Jurgens, G., Saano, A., Sen, R., and S. Timonen. 2003. Nested PCR detection of archaea in defined compartments of pine mycorrhizospheres developed in boreal forest humus microcosms. *FEMS Microbiol.Ecol.* 43: 153-171.
- Borneman, J., and E.W. Triplett. 1997. Molecular microbial diversity in soils from eastern Amazonia: evidence for unusual microorganisms and microbial population shifts associated with deforestation. *Appl.Environ.Microbiol.* 63: 2647-2653.
- Brock, T.D., and H. Freeze. 1969. *Thermus aquaticus* gen. n. and sp. n. a non-sporulating extreme thermophile. *J.Bacteriol.* 98: 289-297.
- Burggraf, S., Heyder, P., and N. Eis. 1997. A pivotal archaea group. *Nature* 385: 780.
- Carpenter, E.J., Lin, S., and D.G. Capone. 2000. Bacterial activity in South Pole snow. *Appl.Environ.Microbiol.* 66: 4514-4517.
- Chandler, D.P., Li, S.M., Spadoni, C.M., Drake, G.R., Balkwill, D.L., Fredrickson, J.K., and F.J. Brockman. 1997. A molecular comparison of culturable aerobic heterotrophic bacteria and 16S rDNA clones derived from a deep subsurface sediment. *FEMS Microbiol.Ecol.* 23: 131-144.
- Chelius, M.K., and E.W. Triplett. 2001. The diversity of archaea and bacteria in association with the roots of *Zea mays*. *Lett.Microb.Ecol.* 41: 252-263.
- Childers, S. E., and D.R. Lovley. 2001. Differences in Fe(III) reduction in the hyperthermophilic archaeon, *Pyrobaculum islandicum*, versus mesophilic Fe(III)-reducing bacteria. *FEMS Microbiol.Lett.* 195: 253-258.
- Chin, K., Lukow, T. and R. Conrad. 1999. Effect of temperature on structure and function of the methanogenic archaeal community in an anoxic rice field soil. *Appl.Environ.Microbiol.* 65: 2341-2349.
- Cowan, D.A. 2004. The upper temperature for life – where do we draw the line? *TRENDS Microbiology* 12: 58-60.
- Crozier, R.H., Agapov, P.M., and K. Pedersen. 1999. Towards complete biodiversity assessment: an evaluation of the subterranean bacterial communities in the Oklo region of the sole surviving natural nuclear reactor. *FEMS Microbiol.Ecol.* 28: 325-334.

Cuadros-Orellana, S., Pohlschröder, M., and L.R. Durrant. 2006. Isolation and characterization of halophilic archaea able to grow in aromatic compounds. *Int.Biodeterior.Biodegrad.* 57: 151-154.

Daly, M. 2000. Engineering radiation-resistant bacteria for environmental biotechnology. *Cur.Opin.Biotech.* 11: 280-285.

De, G.C., Oliver, D.J., and B.M. Pesic. 1996. Effect of silver on the ferrous iron oxidizing ability of *Thiobacillus ferrooxidans*. *Hydrometallurgy* 41: 211-229.

DeLong, E.E., and N.R. Pace. 2001. Environmental diversity of *Bacteria* and *Archaea*. *Syst.Biol.* 50: 470-478.

Detrich, H.W., Parker, S.K., Williams, R.C., Jr., Nogales, E., and K.H. Downing. 2000. Cold adaptation of microtubule assembly and dynamics. Structural interpretation of primary sequence changes present in the α - and β -tubulins of Antarctic fishes. *J.Biol.Chem.* 275: 37038-37047.

Dew, D.W., Muhlbauer, R., and C. van Buuren. 1999. Bioleaching of copper sulphide concentrates with mesophiles and thermophiles. In *Alta Copper 99*. Brisbane, Australia.

Downie, A.W., and J. Cruickshank. 1928. The resistance of *Streptococcus fecalis* to acid and alkaline media. *Br.J.Exp.Pathol.* 9: 171-173.

Dopson, M., Baker-Austin, C., Koppineedi, P.R., and P.L. Bond. 2003. Growth in sulfidic mineral environments: metal resistance mechanisms in acidophilic micro-organisms. *Microbiol.* 149: 1959-1970.

Dunbar, J., Takala, S., Barns, S.M., Davis, J.A., and C.R. Kuske. 1999. Levels of bacterial community diversity in four arid soils compared by cultivation and 16S rRNA gene cloning. *Appl.Enviorn.Microbiol.* 65: 1662-1669.

Dunfield, P.F., Khmelenina, V.N., Suzina, N.E., Trotsenko, Y.A., and S.N. Dedysh. 2003. *Methylocella silvestris* sp. nov., a novel methanotroph isolated from an acidic forest cambisol. *Int.J.Syst.Evol.Microbiol.* 53: 1231-1239.

Echigo, A., Hino, M., Fukushima, T., Kamekura, T.M., and R. Usami. 2005. Endospores of halophilic bacteria of the family *Bacillaceae* isolated from non-saline Japanese soil may be transported by Kosa event (Asian dust storm). *Saline Systems* 20: 1-8.

Edmond, J., L'Haridon, S., Corre, E., Forterre, P., and D. Prieur. 2003. *Thermococcus gammatolerans* sp. nov., a hyperthermophilic archaeon from deep-sea hydrothermal vent that resists to ionizing radiation. *Int.J.Syst.Evol.Microbiol.* 53: 847-851.

Ehrlich, H.L. 1984. Growth of *Thiobacillus ferrooxidans* in the presence of silver compounds. In *Abstracts of the 84th Annual Meeting of the American Society for Microbiology*, p. 197. Washington, DC: American Society for Microbiology.

Euzéby, J.P. 2006. List of Prokaryotic names with standing in nomenclature. <http://www.bacterio.cict.fr/>. Last update: February 10, 2006.

Feller, G., and C. Gerday. 2003. Psychrophilic enzymes: hot topics in cold adaptation. *Nature Rev. Microbiology* 1: 200-208.

Fisk, M. R., Giovannoni, S. J., and I.H. Thorseth. 1998. Alteration of oceanic volcanic glass: textural evidence of microbial activity. *Science* 281: 978-980.

Fierer, N., and R.B. Jackson. 2006. The diversity and biogeography of soil bacterial communities. *Proc. Natl. Acad. Sci.* 103: 626-631.

Franks, F. 1985. *Biophysics and biochemistry at low temperatures*. Cambridge University Press, Cambridge.

Friedmann, E.I. 1982. Endolithic microorganisms in the Antarctic cold desert. *Science* 215: 1045-1053.

Fujiwara, S. 2002. Extremophiles: developments of their special functions and potential resources. *J. Bioscience Bioengineering* 94: 518-525.

Ghosal, D., Omelchenko, M.V., Gaidamakova, E.K., Matrosova, V.Y., Vasilenko, A., Venkateswaran, A., Zhai, M., Kostandarithes, H.M., Brim, H., Makarova, K.S., Wackett, L.P., Fredrickson, J.K., and Daly, M.J. 2005. How radiation kills cells: Survival of *Deinococcus radiodurans* and *Shewanella oneidensis* under oxidative stress. *FEMS Microbiol. Rev.* 29: 361-375.

Gilichinský, D., Rivkina, E., Bakermans, C., Shcherbakova, V., Petrovskaya, L., Ozerskaya, S., Ivanushkina, N., Kochkina, G., Laurinavichus, K., Pecheritsina, S., Fattakhova, R., and J.M. Tiedje. 2005. Biodiversity of cryopegs in permafrost. *FEMS Microbiol. Ecology* 53: 117-128.

Gonzalez, J.M., Kato, C., and K. Horikoshi. 1995. *Thermococcus peptonophilus* sp. nov., a fast-growing, extremely thermophilic archaeobacterium isolated from deep-sea hydrothermal vents. *Arch. Microbiol.* 164: 159-164.

Gonzalez, J.M., Masuchi, Y., Robb, F.T., Ammerman, J.W., Yanagibayashi, M., Tamaoka, J., and Kato, C. 1998. *Pyrococcus horikoshii* sp. nov., a hyperthermophilic archaeon isolated from a hydrothermal vent at Okinawa Trough. *Extremophiles* 2: 123-130.

Greve, R. 2000. Waxing and Waning of the Perennial North Polar H₂O Ice Cap of Mars over Obliquity Cycles. *Icarus* 144: 419-431.

Großkopf, R., Janssen, P.H., and W. Liesack. 1998a. Diversity and structure of the methanogenic community in anoxic rice paddy soil microcosms as examined by cultivation and direct 16S rDNA sequence retrieval. *Appl. Environ. Microbiol.* 64: 960-969.

Großkopf, R., Stubner, S., and W. Liesack. 1998b. Novel euryarchaeotal lineages detected on rice roots and in the anoxic bulk soil of flooded rice microcosms. *Appl. Environ. Microbiol.* 64: 4983-4989.

Guay, R., Ghosh, J., and A.E. Torma. 1989. Kinetics of microbiological production of ferric ion for heap and dump leaching. In *Biotechnology in minerals and metals processing*, eds.: B.J. Scheiner, F.M. Doyle, and S.K. Kawatra, 95-106. Littleton, CO: Society of Mining Engineers.

Haberle, M., and B.M. Jakosky. 1990. Sublimation and transport of water from the north residual polar cap on Mars. *J. Geophys. Res.* 95: 1423-1437.

Hacène, H., Rafa, F., Chebhouni, N., Boutaiba, S., Bhatnagar, T., Baratti, J.C., and B. Ollivier. 2004. Biodiversity of prokaryotic microflora in El Golea Salt lake, Algerian Sahara. *J. Arid Environments* 58: 273-284.

Häder, D.-P. 2000. Effects of solar UV-B radiation on aquatic ecosystems. *Adv. Space Res.* 26: 2029-2040.

Hallberg, K.B., and D.B. Johnson. 2003. Novel acidophiles isolated from moderately acidic mine drainage waters. *Hydrometallurgy* 71: 139-148.

Haney, P.J., Badger, J.H., Buldak, G.L., Reich, C.L., Woese, C.R., and G.J. Olsen. 1999. Nucleotide thermal adaptation analyzed by comparison of protein sequences from mesophilic and extremely thermophilic *Methanococcus* species. *Proc. Natl. Acad. Sci.* 96: 3578-3583.

Harris, M.J., Wickramasinghe, N.C., Lloyd, D., Narlikar, J.V., Rajaratnam, P., Turner, M.P., Al-Mufti, S., Wallis, M.K., Ramadurai, S., and F. Hoyle. 2002. The detection of living cells in stratospheric samples. In *Instruments, Methods, and Missions for Astrobiology IV*, eds. R.B. Hoover, A. Rosanov, and R. Paepe. Proc.SPIE, San Diego, 4495: 192-198.

Hastings, J.W., Holzapfel, W.H., and J.G. Niemand. 1986. Radiation resistance of Lactobacilli isolated from radurized meat relative to growth and environment. *Appl. Environ. Microbiol.* 52: 898-901.

Hofle, M.G., Haas, H., and K. Dominik. 1999. Seasonal dynamics of bacterioplankton community structure in a eutrophic lake as determined by 5S rRNA analysis. *Appl. Environ. Microbiol.* 65: 3164-3174.

- Holden, J.F., and M.W. Adams. 2003. Microbe-metal interactions in marine hydrothermal environments. *Curr.Opinion Chem.Biol.* 7: 160-165.
- Hoover, R.B., Hoyle, F., Wickramasinghe, N.C., Høover, M.J., and S. Al-Mufti. 1986. Diatoms on Earth, Comets, Europa, and in Interstellar Space. *Earth, Moon, and Planets* 35: 19-45.
- Hoover, R.B., and D. Gilichinsky. 2001. Significance to Astrobiology of Microorganisms in Permafrost and Ice. In *Permafrost Response on Economic Development, Environmental Security and Natural Resource Potential*, ed. R. Paepe, 553-580. NATO-ARW held in NOVOSIBIRSK, SIBERIA, 12-16 Nov. 1998. Kluwer Publishing, New York.
- Hoover, R.B., Pikuta, E.V., Marsic, D., and J. Ng. 2002. Anaerobic psychrophiles from Alaska, Antarctica, and Patagonia: Implications to possible life on Mars and Europa. In *Methods, and Missions for Astrobiology IV*, eds. R.B. Hoover, G.V. Levin, R.R. Paepe, A.Yu. Rozanov, Proc. SPIE, 4495: 313-324.
- Hoover, R.B., Pikuta, E.V., Bej, A.K., Marsic, D., Whitman, W.B., Tang, J., and P. Krader. 2003. *Spirochaeta americana* sp. nov., a new haloalkaliphilic, obligately anaerobic spirochete isolated from soda Mono Lake in California. *Int.J.Syst.Evol.Microbiol.* 53: 815-821.
- Hoover, R.B., and E.V. Pikuta. 2004. Microorganisms on comets, Europa and the polar ice caps of Mars. In *Methods, and Missions for Astrobiology VII*, ed(s) R.B. Hoover, A.Yu. Rozanov, Proc. SPIE 5163: 191-201.
- Hoover, R.B., Pikuta, E.V., Wickramasinghe, N.C., Wallis, M.K., and R.B. Sheldon. 2005. Astrobiology of comets. In *Instruments, methods, and missions for astrobiology VIII*, eds. R.B. Hoover, A.Yu. Rozanov, and G.V. Levin, Proc. SPIE 5555: 93-106.
- Hoover, R.B. 2005. Microfossils, biominerals, and chemical biomarkers in meteorites. In *Perspectives in Astrobiology, NATO Science Series: Life and Behavioural Sciences*, eds. R.B. Hoover, R.R. Paepe, and A. Yu. Rozanov, IOS Press, Amsterdam, Netherlands, 336: 43-65.
- Hoover, R.B. 2006. Fossils of prokaryotic microorganisms in the Orgueil meteorite. In *Instruments, Methods and Missions for Astrobiology IX*, eds.R.B. Hoover, A.Yu. Rozanov, and G.V. Levin, SPIE, 630902: 1-17.
- Horikoshi, K. 1999. Alkaliphiles: some applications of their products for biotechnology. *Microbiol.Molecular Biology Reviews* 63: 735-750.
- Hoyle, F., and C. Wickramasinghe. 1981. Comets - A Vehicle for Panspermia. In *Comets and the Origin of Life*, ed. C. Ponnampurama, 222-239. D. Reidel Publ. Co., Dordrecht.
- Horikoshi, K., and T. Akiba. 1982. *Alkalophilic microorganisms: a new microbial world*. Springer-Verlag KG, Heidelberg, Germany.

Hoyoux, A., Blaise, V., Collins, T., D'Amico, S., Gratia, E., Huston, A.L., Marx, J.-C., Sonan, G., Zeng, Y., Feller, G., and C. Gerday. 2004. Extreme catalysts from low-temperature environments. *J.Biosc.Bioeng.* 98: 317-330.

Huber, H., and K.O. Stetter. 1998. Hyperthermophiles and their possible potential in biotechnology. *J.Biotechnol.* 64: 39-52.

Huber, G., Spinnler, C., Gambacorta, A., and K.O. Stetter. 1989. *Metallosphaera sedula* gen. and sp. nov. represents a new genus of aerobic, metal-mobilizing, thermoacidophilic archaeobacteria. *Syst.Appl.Microbiol.* 12: 38-47.

Huber, H., Hohn, M.J., Stetter, K.O., and R. Rachel. 2003. The phylum Nanoarchaeota: Present knowledge and future perspectives of a unique form of life. *Res.Microbiology* 154: 165-171.

Hungate, R.E. 1969. A roll tube method for cultivation of strict anaerobes. In *Methods in Microbiology*, eds. J.R. Norris, and D.W. Ribbons. 3B: 117-132. New York. Academic Press.

Ibekwe, A.M., and A.C. Kennedy. 1998. Fatty acid methyl ester (FAME) profiles as a tool to investigate community structure of two agricultural soils. *Plant Soil* 206: 151-161.

Iizuka, H., Okazaki, H., and N. Seto. 1969. A new sulfate-reducing bacterium isolated from Antarctica. *J.Gen.Appl.Microbiol.* 15: 11-18.

Itoh, T. 2003. Taxonomy of nonmethanogenic hyperthermophilic and related thermophilic Archaea. *J.Biociencia Bioengineering* 96: 203-212.

Jeanthon, C., Reysenbach, A.L., Haridon, S., Gambacorta, A., Pace, N.R., Glénat, P., and D. Prieur. 1995. *Thermotoga subterranean* sp. nov., a new thermophilic bacterium isolated from a continental oil reservoir. *Arch.Microbiol.* 164: 91-97.

Johnson, D.B., Ghauri, M.A., and M.F. Said. 1992. Isolation and characterization of an acidophilic, heterotrophic bacterium capable of oxidizing ferrous iron. *Appl.Environ.Microbiol.* 58: 1423-1428.

Johnson, D.B. 1998. Biodiversity and ecology of acidophilic microorganisms. *FEMS Microbiol.Ecol.* 27: 307-317.

Joo, W.A., and C.W. Kim. 2005. Proteomics of halophilic archaea. *J.Chromatography B* 815: 237-250.

Kalyuzhnaya, M.G., Khmelenina, V.N., Kotelnikova, S., Holmquist, L., Pedersen, K., and Yu.A. Trotsenko. 1999. *Methylomonas scandinavica*, sp. nov., a new methanotrophic psychrotrophic bacterium isolated from deep igneous rock ground water of Sweden. *Syst.Appl.Microbiol.* 22: 565-572.

- Kashefi, K., and D. Lovely. 2003. Extending the upper temperature limit for life. *Science* 301: 934.
- Kato, C., Sato, T., and K. Horikoshi. 1995. Isolation and properties of barophilic and barotolerant bacteria from deep-sea mud samples. *Biodiv. Conserv.* 4: 1-9.
- Kato, C., Masui, N., and K. Horikoshi. 1996. Properties of obligatory barophilic bacteria isolated from a sample of deep-sea sediments from the Izu-Bonin trench. *J.Mar.Biotechnol.* 4: 96-99.
- Kato, C., Li, L., Nogi, Y., Nakamura, Y., Tamaoka, J., and K. Horikoshi. 1998. Extremely barophilic bacteria isolated from the Mariana Trench, Challenger Deep, at a depth of 11,000 meters. *Appl. Environ. Microbiol.* 64: 1510-1513.
- Kieffer, H.H., Chose, S.C., Martin, T.Z, Miner, E.D., and F.D. Palluconi. 1976. Martian North pole summer temperature: Dirty water ice. *Science* 194: 1341-1344.
- Kent, A.D., and E.W. Triplett. 2002. Microbial communities and their interactions in soil and rhizosphere ecosystems. *Annu. Rev. Microbiol.* 56: 211-236.
- Kishimoto, N., Kosako, Y., and T. Tano. 1991. *Acidobacterium capsulatum* gen. nov., sp. nov.: an acidophilic chemoorganotrophic bacterium containing menaquinone from acidic mineral environment. *Curr. Microbiol.* 22: 1-7.
- Kitts, C.L. 2001. Terminal restriction fragment patterns: a tool for comparing microbial communities and assessing community dynamics. *Curr. Issues Intest. Microbiol.* 2: 17-25.
- Knoblauch, C., Sahm, K., and B.B. Jørgensen. 1999. Psychrophilic sulfate-reducing bacteria isolated from permanently cold Arctic marine sediments: description of *Desulfofrigus oceanense* gen. nov., sp. nov., *Desulfofrigus fragile* sp. nov., *Desulfobaba gelida* gen. nov., sp. nov., *Desulfotalea psychrophila* gen. nov., sp. nov. and *Desulfotalea arctica* sp. nov. *Int. J. Syst. Bacteriol.* 49: 1631-1643.
- Kobayashi, T., Kwak, Y.S., Akiba, T., Kudo, T., and K. Horikoshi. 1994. *Thermococcus profundus* sp. nov., a new hyperthermophilic archaeon isolated from a deep-sea hydrothermal vent. *System. Appl. Microbiol.* 17: 232-236.
- Kolber, Z.S., Plumley, F.G., Lang, A.S., Beatty, J.T., Blankenship, R.E., VanDover, C.L., Vetriani, C., Koblizek, M., Rathgeber, C., and Falkowski, P.G. 2001. Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* 292: 2492-94.
- Kotelnikova, S., and K. Pedersen. 1998. Distribution and activity of methanogens and homoacetogens in deep granitic aquifers at Åspö Hard Rock Laboratory, Sweden. *FEMS Microbiol. Ecol.* 26: 121-134.

- Kotelnikova, S., Macario, A.J.L., and K. Pedersen. 1998. *Methanobacterium subterraneum*, a new species of Archaea isolated from deep groundwater at the Åspö Hard Rock Laboratory, Sweden. *Int.J.Syst.Bacteriol.* 48: 357-367.
- Kuever, J., Rainey, F.A., and F. Widdel. 2005. Family IV. *Desulfonatronumaceae* fam. nov. In *Bergey's Manual of Systematic Bacteriology, 2nd edition*, eds. D.J. Brenner, N.R. Krieg, J.T. Staley, and G.M. Garrity, V.2 (The Proteobacteria), part C (The Alfa-, Beta-, Delta-, and Epsilonproteobacteria), p. 956. Springer, New York.
- Kuhner, C.H., Matthies, C., Acker, G., Schmittroth, M., Gossner, A.S., and H.L. Drake. 2000. *Clostridium akagii* sp. nov. and *Clostridium acidisoli* sp. nov.: acid-tolerant, N₂-fixing clostridia isolated from acidic forest soil and litter. *Int.J.Syst.Evol.Microbiol.* 50: 873-881.
- Lacroix, M., and R. Lafortune. 2004. Combined effects of gamma irradiation and modified atmosphere packaging on bacterial resistance in grated carrots (*Daucus carota*). *Rad.Physics and Chemistry* 71: 77-80.
- Li, W., Xu, P., Zhang, L. Tang, S., Cui, X., Mao, P., Xu, L., Schumann, P., Stackebrandt, E., and C. Jiang. 2003. *Streptomonospora alba* sp. nov., a novel halophilic actinomycete, and emended description of the genus *Streptomonospora* Cui *et al.* 2001. *Int.J.Syst.Evol.Microbiol.* 53: 1421-1425.
- Li, W., Schumann, P., Zhang, Y., Chen, G., Tian, X., Xu, L. Stackebrandt, E., and C. Jiang. 2005. *Marinococcus halotolerans* sp. nov., isolated from Qinghai, north-west China. *Int.J.Syst.Evol.Microbiol.* 55: 1801-1804.
- Li, W.F., Zhou, X.X., and Lu, P. 2005. Structural features of thermozymses. *Biotechnol.Advaces* 23: 271-281.
- Liesack, W., Schnell, S., and N.P. Revsbech. 2000. Microbiology of flooded rice paddies. *FEMS Microbiol.Rev.* 24: 625-645.
- Liles, M.R., Manske, B.F., Bintrim, S.B., Handelsman, J., and R.M. Goodman. 2003. A census of rRNA genes and linked genomic sequences within a soil metagenomic library. *Appl.Environ.Microbiol.* 69:2684-2691.
- Litvak, M.L., Mitrofanov, I.G., Kozyrev, A.S., Sanin, A.B., Tretyakov, V.I., Boynton, W.V., Kelly, N.J., Hamara, D., Shinohara, C., and R.S. Saunders. 2006. Comparison between polar regions of Mars from HEND/Odyssey data. *Icarus* 180: 23-37.
- Makarova, K.S., Aravind, L., Wolf, Yu.I., Tatusov, R.L., Minton, K.W., Koonin, E.V., and M.J. Daly. 2001. Genome of the extremely radiation-resistant bacterium *Deinococcus radiodurans* viewed from the perspective of comparative genomics. *Microbiol.Mol.Biol.Rev.* 65: 44-79.

- Makarova, K.S., and E.V. Koonin. 2005. Evolutionary and functional genomics of the Archaea. *Curr.Opin.Microbiol.* 8: 586-594.
- Mancinelli, R.L., Fahlen, T.F., Landheim, R., and M.R. Klovstad. 2004. Brines and evaporites: analogs for Martian life. *Adv.Space Res.* 33: 1244-1246.
- Marsh, T.L., Saxman, P., Cole, J., and J. Tiedje. 2000. Terminal restriction fragment length polymorphism analysis program, a web-based research tool for microbial community analysis. *Appl.Environ.Microbiol.* 66: 3616-3620.
- Marteinsson, V.T., Reysenbach, A.L., Birrien, J. L., and D. Prieur. 1999. A stress protein is induced in the deep-sea barophilic hyperthermophile *Thermococcus barophilus* when grown under atmospheric pressure. *Extremophiles* 3: 277-282.
- McKay, D.S., Gibson, E.K., Thomas-Keppta, K.L., Vali, H., Romanek, C.S., Clemett, S.J., Chillier, X.D.F., Maechling, C.R., and Zare, R.N. 1996. Search for past life on Mars: Possible relic biogenic activity in Martian meteorite ALH84001. *Science* 273: 924-930.
- Meek C.S., and C.B. Lipman. 1922. The relation of the reaction and of salt content of the medium on nitrifying bacteria. *J.Gen.Physiol.* 5: 195-204.
- Min, H., Zhao, Y.H. Chen, M.C., and Y. Zhao. 1997. Methanogens in paddy rice soil. *Nutr.Cycl.Agroecosyst.* 49: 163-169.
- Mitrofanov, I.G., Litvak, M.L., Kozyrev, A.S., Sanin, A.B., Tret'yakov, V.I., Grin'kov, V.Yu., Boynton, W.V., Shinohara, C., Hamara, D., and Saunders, R.S. 2004. Soil Water Content on Mars as Estimated from Neutron Measurements by the HEND Instrument On board the 2001 *Mars Odyssey* Spacecraft. *Solar System Research* 38: 253-357.
- McGenity, T.J., Gemmell, R.T., Grant, W.D., and Stan-Lotter. 2000. Origins of halophilic microorganisms in ancient salt deposits. *Environ.Microbiol.* 2: 243-250.
- Meek, C.S., and C.B. Lipman. 1922. The relation of the reaction and of salt content of the medium on nitrifying bacteria. *J.Gen.Physiol.* 5: 195-204.
- Morita, R.Y. 1975. Psychrophilic bacteria. *Bacteriol.Rev.* 39: 144-167.
- Motamedi, M., and K. Pedersen. 1998. *Desulfovibrio aespoensis* sp. nov., a mesophilic sulfate-reducing bacterium from deep groundwater at Åspö Hard Rock Laboratory, Sweden. *Int.J.Syst.Bacteriol.* 48: 311-315.
- Moseley, B.E., and D.M. Evans. 1983. Isolation and properties of strains of *Micrococcus (Deinococcus) radiodurans* unable to excise ultraviolet light-induced pyrimidine dimmers from DNA: evidence for two excision pathways. *J.Gen.Microbiol.* 129: 2437-2445.

Motamedi, M., and K. Pedersen. 1998. *Desulfovibrio aespoensis* sp. nov., a mesophilic sulfate-reducing bacterium from deep groundwater at Åspö Hard Rock Laboratory, Sweden. *Int.J.Syst.Bacteriol.* 48: 311-315.

Mrázek, J. 2002. New technology may reveal mechanisms of radiation resistance in *Deinococcus radiodurans*. *Proc.Natl.Acad.Sci.* 99: 10943-10944.

Nagi, A.D., and L. Regan. 1997. An inverse correlation between loop length and stability in a four-helix-bundle protein. *Fold.Des.* 2: 67-75.

Nagy, M.L., Perez, A., and F. Garcia-Pichel. 2005. The prokaryotic diversity of biological soil crusts in the Sonoran Desert (Organ Pipe Cactus National Monument, AZ). *FEMS Microbiol.Ecol.* 54: 233-245.

Nakajima, K., Hirota, K., Nodasaka, Y., and I. Yumoto. 2005. *Alkalibacterium iburiense* sp. nov., an obligate alkaliphile that reduces an indigo dye. *Int.J.Syst.Evol.Microbiol.* 55: 1525-1530.

Nemergut, D.R., Costello, E.K., Meyer, A.G., Pescador, M.Y., Weintraub, M.N., and S.K. Schmidt. 2005. Structure and function of alpine and arctic soil microbial communities. *Res.Microbiol.* 156: 775-784.

Nicol, G.W., Webster, G., Glover, L.A., and J.I. Prosser. 2004. Differential response of archaeal and bacterial communities to nitrogen inputs and pH changes in upland Pasture rhizosphere soil. *Environ.Microbiol.* 6: 861-867.

Norris, P.R., and D.B. Johnson. 1998. Acidophilic microorganisms. In *Extremophiles. Microbial life in extreme environment*, eds. K. Horikoshi, and W.D. Grant, 133-153. Wiley-Liss, New York.

Ochsenreiter, T., Selezi, D., Quaiser, A., Bonch-Osmolovskaya, E., and C. Schleper. 2003. Diversity and abundance of *Crenarchaeota* in terrestrial habitats studied by 16S RNA surveys and real time PCR. *Environ.Microbiol.* 5: 787-797.

Odokuma, L.O., and A.A. Dickson. 2003. Bioremediation of a Crude Oil Polluted Tropical Rain Forest Soil. *Global J.Environmental Sciences* 2: 29-40.

Omelchenko, M.V., Wolf, Y.I., Gaidamakova, E.K., Matrosova, V.Y., Vasilenko, A., Zhai, M., Daly, M.J., Koonin, E.V., and K.S. Makarova. 2005. comparative genomics of *Thermus thermophilus* and *Deinococcus radiodurans*: divergent routes of adaptation to thermophily and radiation resistance. *BMC Evol.Biol.* 5: 57.

Oren, A. 1999. Bioenergetic aspects of halophilism. *Microbiol.Mol.Biol.Rev.* 663: 334-348.

Oren, A. 2002. Diversity of halophilic microorganisms: environments, phylogeny, physiology, and applications. *J.Ind.Microbiol.Biotechnol.* 28: 56-63.

- Osborne, A.M., Moore, E.R.B., and K.N. Timmis. 2000. An evaluation of terminal-restriction fragment length polymorphism (T-RFLP) analysis for the study of microbial community structure and dynamics. *Environ.Microbiol.* 2: 39-50.
- Padan, E., Bibi, E., Ito, M., and T.A. Krulwich. 2005. Alkaline pH homeostasis in bacteria: New insights. *Biochim.Biophys.Acta* 1717: 67-88.
- Paepe, R., Hoover, R.B., and E. Van Overloop. 2001. Patterned Ground as Evidence of Water on Mars. In *Permafrost Response on Economic Development, Environmental Security and Natural Resource Potential, NATO-ARW held in NOVOSIBIRSK, SIBERIA, 12-16 Nov. 1998*, ed. R. Paepe, 581-588. Kluwer Publishing, New York.
- Pan, H.L., Zhou, C., Wang, H.L., Xue, Y.F., and Y.H. Ma. 2006. Diversity of halophilic archaea in hypersaline lakes of Inner Mongolia, China. *Wei Sheng Wu Xue Bao* 46: 1-6.
- Pašić, L., Bartual, S.G., Ulrich, N.P., Grabnar, M., and B.H. Velikonja. 2005. Diversity of halophilic archaea in the crystallizers of an Adriatic solar saltern. *FEMS Microbiol.Ecol.* 54: 491-498.
- Pasteur, L. 1861. Animalcules infusoires vivant sans gaz oxygène libre et déterminant des fermentations. *Compt. Rend. Acad. Sci. (Paris)*, 52, 344-347.
- Pedersen, K. 2000. Exploration of deep intraterrestrial microbial life: current perspectives. *FEMS Microbiol.Lett.* 185: 9-16.
- Pedersen, K., and S. Ekendahl. 1990. Distribution and activity of bacteria in deep granitic groundwaters of southeastern Sweden. *Microbiol.Ecol.* 20: 37-52.
- Pedersen, K., Arlinger, J., Ekendahl, S., and L. Hallbeck. 1996. 16S rRNA gene diversity of attached and unattached groundwater bacteria along the access tunnel to the Åspö Hard Rock Laboratory, Sweden. *FEMS Microbiol.Ecol.* 19: 249-262.
- Pick, U. 1999. *Dunaliella acidophila*- a most extreme acidophilic alga. In: *Enigmatic microorganisms in extreme environments*, ed. J. Seckbach, 465-47. Kluwer Academic Press, Dordrecht/The Netherlands.
- Pikuta, E.V., Zhilina, T.N., Zavarzin, G.A., Kostrikina, N.A., Osipov, G.A., and F.A. Rainey. 1998. *Desulfonatronum lacustre* sp. nov.: a new alkaliphilic sulfate-reducing bacterium utilizing ethanol. *Microbiology (English translation of Mikrobiologiya)* 67: 123-131.
- Pikuta, E.V., Lysenko, A.M., Suzina, N., Osipov, G.A., Kuznetsov, B., Tourova, T., Akimenko, V.K., and K.S. Laurinavichus. 2000a. *Desulfotomaculum alkaliphilum* sp. nov., a new alkaliphilic, moderately thermophilic, sulfate-reducing bacterium. *Int.J.Syst.Evol.Microbiol.* 50: 25-33.

- Pikuta, E.V., Lysenko, A.M., Chuvilskaya, N.A., Mendrock, U., Hippe, H., Suzina, N., Nikitin, D., Osipov, G.A., and K.S. Laurinavichius. 2000b. *Anoxybacillus pushchinensis* gen. nov., sp. nov., a novel anaerobic, alkaliphilic, moderately thermophilic bacterium from manure, and description of *Anoxybacillus flavithermus* comb. nov. *Int.J.Syst.Evol.Microbiol.* 50: 2109-2117.
- Pikuta, E.V., and Hoover, R.B. 2003. Psychrophiles and astrobiology: microbial life of frozen worlds. In *Instruments, Methods, and Missions for Astrobiology VI*, eds. R.B. Hoover, A.Yu. Rozanov, and J.H. Lipps. Proc. SPIE, 4939: 103-116.
- Pikuta, E.V., Detkova, E.N., Bej, A.K., Marsic, D., and Hoover, R.B. 2003a. Anaerobic haloalkaliphilic bacterial community of athalassic, hypersaline Mono Lake and Owens Lake in California. In: *Instruments, methods, and missions for astrobiology V*, ed(s): R.B. Hoover, A.Yu. Rozanov, and R.R. Paepe. Proceedings of SPIE, 4859: 130-144. Waikoloa, Hawaii 22-23 August 2002, USA.
- Pikuta, E.V., Hoover, R.B., Marsic, D., Bej, A.K., Tang, J., Cleland, D., and P. Krader. 2003b. Psychrotolerant anaerobic bacteria from Patagonia and the permafrost tunnel of Fox, Alaska. In *103rd General Meeting American Society for Microbiology*, May 18-22, p. 351., Washington, D.C.
- Pikuta, E.V., Hoover, R.B., Bej, A.K., Marsic, D., Whitman, W.B., Cleland, D., and P. Krader. 2003c. *Desulfonatronum thiodismutans* sp. nov., a novel alkaliphilic, sulfate-reducing bacterium capable of lithoautotrophic growth. *Int.J.Syst.Evol.Microbiol.* 53: 1327-1332.
- Pikuta, E.V., Hoover, R.B., Bej, A.K., Marsic, D., Detkova, E.N., Whitman, W.B., and P. Krader. 2003d. *Tindallia californiensis* sp. nov., a new anaerobic, haloalkaliphilic, spore-forming acetogen isolated from Mono Lake in California. *Extremophiles* 7: 327-334.
- Pikuta, E.V., and R.B. Hoover. 2004a. Astrobiological significans of chemolithoautotrophic acidophiles. In *Instruments, methods, and missions for astrobiology VII*, eds. RB Hoover, AYU Rozanov. Proceedings of SPIE 5163: 179-1490.
- Pikuta, E.V., and R.B. Hoover. 2004b. Growth of the facultative anaerobes from Antarctica, Alaska, and Patagonia at low temperatures. In *Instruments Methods and Missions for Astrobiology VIII*, Proc.SPIE 5555: 160-191.
- Pikuta, E.V., Hoover, R.B., Marsic, D., Bej, A.K., Tang, J., and P. Krader. 2005a. *Carnobacterium pleistocaenium* sp. nov., a novel psychrotolerant, facultative anaerobe isolated from Fox Tunnel permafrost, Alaska. *Int.J.Syst.Evol.Microbiol.* 55: 473-478.
- Pikuta, E.V., Itoh, T., and R.B. Hoover. 2005b. Anaerobic decomposition of cellulose by alkaliphilic microbial community of Owens Lake, California. In: *Instruments, methods, and missions for astrobiology IX*, eds.: R.B. Hoover, A.Yu. Rozanov, G.V. Levin, and G.R. Gladstone, Proc.SPIE 5906: 29-40, San Diego, USA.

- Pikuta, E.V., Hoover, R.B., Bej, A.K., Marsic, D., Whitman, W.B., Krader, P.E., and J. Tang. 2006a. *Trichococcus patagoniensis* sp. nov., a facultative anaerobe that grows at -5°C , isolated from penguin guano in Chilean Patagonia. *Int.J.Syst.Evol.Microbiol.* 56: 2055 – 2062.
- Pikuta, E.V., Itoh, T., Krader, P., Tang, J., Whitman, W.B., and R.B. Hoover. 2006b. *Anaerovirgula multivorans* gen. nov., sp. nov., a novel spore-forming, alkaliphilic anaerobe isolated from Owens Lake, California. *Int.J.Syst.Evol.Microbiol.* 53: 2623-2629.
- Pledger, R.J., Crump, B.C., and J.A. Baross. 1994. A barophilic response by two hyperthermophilic, hydrothermal vent Archaea: an upward shift in the optimal temperature and acceleration of growth rate at supra-optimal temperatures by elevated pressure. *FEMS Microbiol.Ecol.* 14: 233-242.
- Prowe, S.G., and G. Antranikian. 2001. *Anaerobranca gottschalkii* sp. nov., a novel thermoalkaliphilic bacterium that grows anaerobically at high pH and temperature. *Int.J.Syst.Evol.Microbiol.* 51: 457-465.
- Raaijmakers, J.M., and D.M. Weller. 1998. Natural plant protection by 2,4-diacetyl-phloroglucinol-producing *Pseudomonas* spp. in take-all decline soils. *Mol.Plant-Microb.Interact.* 11: 144-152.
- Rachel, R., Wyschkony, I., Riehl, S., and H. Huber. 2002. The ultrastructure of *Igniococcus*: Evidence for a novel outer membrane and for intracellular budding in an archaeon. *Archaea* 1: 9-18.
- Raguenes, G., Christen, R., Guezennec, J., Pignet, P., Barbier, G. 1997. *Vibrio diebolicus* sp. nov., a new polysaccharide-secreting organism isolated from a deep-sea hydrothermal vent polychaete annelid, *Alvinella pompejana*. *Int.J.Syst.Bacteriol.* 47: 989-995.
- Rainey, F.A., Ray, K., Ferreira, M., Gatz, B.Z., Nobre, M.F., Bagaley, D., Rash, B.A., Park, M-J., Earl, A.M., Shank, N.C., Small, A.M., Henk, M.C., Battista, J.R., Kämpfer, P., and M.S. da Costa. 2005. Extensive diversity of ionizing-radiation-resistant bacteria recovered from Sonoran Desert soil and description of nine new species of the genus *Deinococcus* obtained from a single soil sample. *Appl.Env.Microbiol.* 71: 5225-5235.
- Raj, H.D., Duryee, F.L., Deeney, A.M., Wang, C.H., Anderson, A.W., and P.R. Elliker. 1960. Utilization of carbohydrates and amino acids by *Micrococcus radiodurans*. *Can.J.Microbiol.* 6: 289-298.
- Ramakrishnan, B., Lueders, T., Conrad, R., and M. Friedrich. 2000. Effect of soil aggregate size on methanogenesis and archaeal community structure in anoxic rice field soil. *FEMS Microbiol.Ecol.* 32: 261-270.
- Ranjard, L., Poly, F., Combrisson, J., Richaume, A., and F. Gourbiere. 2000. Heterogeneous cell density and genetic structure of bacterial pools associated with various soil microenvironments as determined by enumeration and DNA fingerprinting approach (RISA). *Microb.Ecol.* 39: 263-272.

- Ray, B. 1986. Impact of bacterial injury and repair in food microbiology: its past, present and future. *J.Food Prot.* 49: 651-655.
- Raymond, J.A., Sullivan, C.W., and A.L. DeVries. 1994. Release of an ice-active substance by antarctic sea ice diatoms. *Polar Biology* 14: 71-75.
- Raymond, J.A. 1995. Glycerol Synthesis in the rainbow smelt *Osmerus mordax*. *J.Exp.Biol.* 198: 2569-2573.
- Raymond, J.A. 2000. Distribution and partial characterization of an ice-active molecule associated with sea ice diatoms. *Polar Biology* 23: 721-729.
- Ritchie, N.J., Schutter, M.E., Dick, R.P., and D.D. Myrold. 2000. Use of length heterogeneity PCR and fatty acid methyl ester profiles to characterize microbial communities in soil. *Appl.Environ.Microbiol.* 66: 1668-1675.
- Roberts, M.F. 2005. Organic compatible solutes of Halotolerant and halophilic microorganisms. *Saline Systems* 1: 5. (<http://www.salinesystems.org/content/1/1/5>)
- Rothschild, L.J., and R.L. Mancinelli. 2001. Life in extreme environments. *Nature* 409: 1092-1101.
- Russel, N.J., Evans, R.I., terSteege, P.F, Hellemons, J., Verheul, A., and Abee, T. 1995. Membranes as a target for stress adaptation. *Int.J.Food Microbiol.* 28: 255-261.
- Saffary, R., Nandakumar, R., Spencer, D., Robb, F.T., Davila, J.M., Swartz, M., Ofman, L., Thomas, R.J., and J. DiRuggiero. 2002. Microbial survival of space vacuum and extreme ultraviolet irradiation: strain isolation and analysis during a rocket flight. *FEMS Microbiol.Lett.* 215: 163-168.
- Sakuraba, H., and Ohshima, T. 2002. Novel energy metabolism in anaerobic hyperthermophilic archaea: a modified Embden-Meyerhof Pathway. *J.Bioscience and Bioengineering* 93: 441-448.
- Sanders, S.W., and R.B. Maxcy. 1979. Isolation of radiation-resistant bacteria without exposure to irradiation. *Appl.Env.Microbiol.* 38: 436-439.
- Sato, T., Kato, C., and K. Horikoshi. 1995. Effect of high pressure on gene expression by *lac* and *tac* promoters in *Escherichia coli*. *J.Mar.Biotechnol.* 3: 89-92.
- Schäffer, C., W.L. Franck, A. Scheberl, P. Kosma, T.R. McDermott, and P. Messner. 2004. Classification of isolates from locations in Austria and Yellowstone National Park as *Geobacillus tepidamans* sp. nov. *Int.J.Syst.Evol.Microbiol.* 54: 2361-2368.
- Schleper, C., Pühler, G., Kühlmorgen, B., and W. Zillig. 1995. Life at extremely low pH. *Nature* 375: 741-742.

Schleper, C., Pühler, G., Klenk, H.P., and W. Zillig. 1996. *Picrophilus oshimae* and *Picrophilus torridus* fam. nov., gen. nov., sp. nov., two species of hyperacidophilic, thermophilic, heterotrophic, aerobic archaea. *Int.J.Syst.Bacteriol.* 46: 814-816.

Schuliger, J.W., Brown, S.H., Baross, J.A., and R.M. Kelly. 1993. Purification and characterization of a novel amylolytic enzyme from ES4, a marine hyperthermophilic archaeum. *Mol.Mar.Biol.Biotech.* 2: 76-87.

Sessitsch, A., A. Weilharter, A., Gerzabek, M.H., Kirchmann, H., and E. Kandeler. 2001. Microbial population structures in soil particle size fractions of a long-term fertilizer field experiment. *Appl.Environ.Microbiol.* 67: 4215-4224.

Simon, H.M., Dodsworth, J.A., and R.M Goodman. 2000. *Crenarchaeota* colonize terrestrial plant roots. *Environ.Microbiol.* 2: 495-505.

Simon, H.M., Jahn, C.E., Bergerud, L.T., Sliwinski, M.K., Weimer, P.J., Willis, D.K., and R.M. Goodman. 2005. Cultivation of mesophilic soil *Crenarchaeotes* in enrichment cultures from plant roots. *Appl.Environ.Microbiol.* 71: 4751-4760.

Sliwinski, M.K., and R.M. Goodman. 2004. Spatial heterogeneity of Crenarchaeal assemblages within mesophilic soil ecosystems as revealed by PCR-single-stranded conformation polymorphism profiling. *Appl.Environ.Microbiol.* 70: 1811-1820.

Smit, E., Leeftang, P., Gommnas, S., Van Den Broek, J., Van Mil, S., and K. Wernars. 2001. Diversity and seasonal fluctuations of the dominant members of the bacterial soil community in a wheat field as determined by cultivation and molecular methods. *Appl.Environ.Microbiol.* 67: 2284-2291.

Stackebrandt, E.W., Liesack, W, and B.M. Goebel. 1993. Bacterial diversity in a soil sample from a subtropical Australian environment as determined by 16S rDNA analysis. *FASEB J.* 7: 232-236.

Staley, J.T., and J.J. Gosink. 1999. Poles apart: biodiversity and biogeography of sea ice bacteria. *Annu.Rev.Microbiol.* 53: 189-215.

Stetter, K O., Thomm, M., Winter, J., Wildgruber, G., Huber, H., Zillig, W., Janecovic, D., König, H., Palm, P., and S. Wunderl. 1981. *Methanothermus fervidus*, sp. nov., a novel extremely thermophilic methanogen isolated from an Icelandic hot spring. *Zentrabl. Für Bakteriologie und Hygiene I. Abteilung Originale* C2: 166-178.

Szewzyk, U., Szewzyk, R., and T. Stenström. 1994. Thermophilic, anaerobic bacteria isolated from deep borehole in granite in Sweden. *Proc.Natl.Acad.Sci.* 91: 1810-1813.

Takami, H., Inoue, A., Fujii, F., and K. Horikoshi. 1997. Microorganisms isolated from the deepest sea mud of Mariana Trench. *FEMS Microbiol.Lett.* 152: 279-285.

Thorseth, I.H., Torsvik, T., Furnes, H., and K. Muehlenbachs. 1995. Microbes play an important role in the alteration of oceanic crust. *Chem.Geol.* 126: 137-146.

Tindall, B.J., Ross, H.N.M., and W.D. Grant. 1984. *Natronobacterium* gen. nov., and *Natronococcus* gen. nov., two new genera of halophilic archaeobacteria. *Syst.Appl.Microbiol.* 5: 41-57.

Torsvik, V., Salte, K., Sorheim, R., and J. Goksoyr. 1990. Comparison of phenotypic diversity and DNA heterogeneity in a population of soil bacteria. *Appl.Environ.Microbiol.* 56: 776-781.

Torsvik, V., Ovreas, L., and T.F. Thingstad. 2002. Prokaryotic diversity-magnitude, dynamics, and controlling factors. *Science* 296: 1064-1066.

Trevors, J.T., Oddie, K.M., and B.H. Belliveau. 1985. Metal resistance in bacteria. *FEMS Microbiol.Rev.* 32: 39-54.

Tringe, S.G., Von Mering, C., Kobayashi, A., Salamov, A.A., Chen, K., Chang, H.W., Podar, M., Short, J.M., Mathur, E.J., Detter, J.C., Bork, P., Hugenholtz, P., and E.M. Rubin. 2005. Comparative metagenomics of microbial communities. *Science* 308: 554-557.

Trotsenko, Yu.A., and V.N. Khmelenina. 2002. Biology of extremophilic and extremotolerant methanotrophs. *Arch.Microbiol.* 177: 123-131.

Ueda, T., Suga, Y., and T. Matsuguchi. 1995. Molecular phylogenetic analysis of a soil microbial community in a soybean field. *Eur.J.Soil.Sci.* 46: 415-421.

Utsumi, M., Belova, S.E., King, G.M., and H. Uchiyama. 2003. Phylogenetic comparison of methanogen diversity in different wetland soils. *J.Gen.Appl.Microbiol.* 49: 75-83.

Van Den Burg, B. 2003, Extremophiles as a source for novel enzymes. *Curr.Opin.Microbiol.* 6: 213-218.

Van Dover, C.L., Reynolds, G.T., Chave, A.D., and J.A. Tyson. 1996. Light at deep-sea hydrothermal vents. *Geophysical Res.Lett.* 23: 2049-2052.

Van Heurck, H. 1909. Diatomées. *Expédition Antarctique Belge. Resultats du voyage du S. Y. Belgica en 1897-1898-1899 sous le commandement de Gerlache de Gomery. Rapports scientifiques Botanique*, 5: 1-128.

Vieille, C., and G.J. Zeikus. 2001. Hyperthermophilic enzymes: sources, uses, and molecular mechanisms for thermostability. *Microbiol.Mol.Biol.Rev.* 65: 1-43.

Wadhwa, M., and G.W. Lugmair. 1996. The formation age of carbonates in ALH 84001. *Meteoritics* 31: A145.

Wagner, D., Spieck, E., Bock, E., and E.-M. Pfeiffer. 2001. Microbial life in terrestrial permafrost: methanogenesis and nitrification in gelisols as potentials for exobiological processes. www.awi-bremerhaven.de/Publications/Wag2001a.pdf

Walsh, D.A., Papke, R.T., and W.F. Doolittle. 2005. Archaeal diversity along a soil salinity gradient prone to disturbance. *Environ.Microbiol.* 7: 1655-1666.

Wang, C.L., Michels, P.C., Dawson, S.C., Kitisakkul, S., Baross, J.A., Keasling, J.D., and D.S. Clark. 1997. Cadmium removal by a new strain of *Pseudomonas aeruginosa* in aerobic culture. *Appl.Environ.Microbiol.* 63: 4075-4078.

Welch, T. J., and D.H. Bartlett. 1996. Isolation and characterization of structural gene for *OmpL*, a pressure-regulated porin-like protein from the deep-sea bacterium *Photobacterium* species strain SS9. *J.Bacteriol.* 178: 5027-5031.

Welsh, D.T., Lindsay, Y.E., Caumette, P., Herbert, R.A., and Hannan, J. 1996. Identification of trehalose and glycine betaine as compatible solutes in the moderately halophilic sulfate reducing bacterium, *Desulfovibrio halophilus*. *FEMS Microbiol.Lett.* 140: 203-207.

Weisser, J., and H.G. Trüper. 1985. Osmoregulation in a new haloalkaliphilic *Bacillus* from Wadi Natrun (Egypt). *Syst.Appl.Microbiol.* 6: 7-11.

Wery, N., Gerike, U., Sharman, A., Chaudhuri, J.B., Hough, D.W., and M.J. Danson. 2003. Use of a packed-column bioreactor for isolation of diverse protease-producing bacteria from Antarctic soil. *Appl.Environ.Microbiol.* 69: 1457-1464.

Whitman, W.B., Coleman, D.C., and W.J. Wiebe. 1998. Prokaryotes: the unseen majority. *Proc.Natl.Acad.Sci.* 95: 6578-6583.

Wiegel, J., and V.V. Kevbrin. 2004. Alkalithermophiles. *Biochem.Soc.Trans.* 32: 193-198.

Wong, C.M., Silver, M., and D.J. Kushner. 1982. Effects of chromium and manganese on *Thiobacillus ferrooxidans*. *Can.J.Microbiol.* 28: 536-544.

Yano, J.K., and T.L. Poulos. 2003. New understandings of thermostable and peizostable enzymes. *Curr.Opinion Biotechnol.* 14: 360-365.

Yayanos, AA., Dietz, A.S., and R.V. Van Boxtel. 1979. Isolation of a deep-sea barophilic bacterium and some of its growth characteristics. *Science* 205: 808-810.

Yayanos, A.A. 1986. Evolutional and ecological implications of the properties of deep-sea barophilic bacteria. *Proc.Natl.Acad.Sci.* 83: 9542-9546.

Zavarzin, G.A., Zhilina, T.N., and V.V. Kevbrin. 1999. The alkaliphilic microbial community and its functional diversity. *Microbiology* 68: 503-521. (Translated from *Mikrobiologiya* 68: 579-599).

↔ Zavarzin, G.A., and T.N. Zhilina. 2000. Anaerobic chemotrophic alkaliphiles. In: *Journey to Diverse Microbial Worlds*, ed. J. Seckbach, pp. 191-208. Kluwer Academic Publishers, Netherlands.


Zhilina, T.N., Zavarzin, G.A., Rainey, F., Kevbrin, V.V., Kostrikina, N.A., and A.M. Lysenko. 1996. *Spirochaeta alkalica* sp. nov., *Spirochaeta africana* sp. nov., and *Spirochaeta asiatica* sp. nov., alkaliphilic anaerobes from the continental soda lakes in Central Asia and the East African Rift. *Int.J.Syst.Evol.Microbiol.* 46: 305-312.

Zhilina, T.N., Zavarzin, G.A., Rainey, F.A., Pikuta, E.V., Osipov, G.A., and N.A. Kostrikina. 1997. *Desulfonatronovibrio hydrogenovorans* gen. nov., sp. nov., an alkaliphilic, sulfate-reducing bacterium. *Int.J.Syst.Evol.Microbiol.* 47: 144-149.


Zhilina, T.N., Zavarzina, D.G., Kuever, J., Lysenko, A.M., and G.A. Zavarzin. 2005. *Desulfonatronum cooperativum* sp. nov., a novel hydrogenotrophic alkaliphilic sulfate-reducing bacterium, from a syntrophic culture growing on acetate. *Int.J.Syst.Evol.Microbiol.* 55: 1001-1006.

Zhou, J., Davey, M.E., Figueras, J.B., Rivkina, E., Gilichinsky, D., and J.M. Tiedje. 1997. Phylogenetic diversity of a bacterial community determined from Siberian tundra soil DNA. *Microbiology* 143: 3913-3919.


Zuber, M.T., Smith, D.E., Solomon, S.C., Abshire, J.B., Afzal, R.S., Aharonson, O., Fishbaugh, K., Ford, P.G., Frey, H.V., Garvin, J.B., Head, J.W., Ivanov, A.B., Johnson, C.L., Muhleman, D.O., Neumann, G.A., Pettengill, G.H., Phillips, R.J., Sun, X., Zwally, H.J., Banerdt, W.B., and T.C. Duxbury. 1998. Observations of the north polar region of Mars from the Mars Orbiter Laser Altimeter. *Science* 282: 2053-2060.




a.



b.



c.



d.

Fig.1. Diagram showing distribution of microorganisms in matrix: (a) pH/Temperature, (b) salinity/pH, (c) salinity/Temperature, (d).radiation/ Temperature.