

# Spectral Analysis of the Accretion Flow in NGC 1052 with Suzaku

L. W. Brenneman<sup>1</sup>, K. A. Weaver<sup>2</sup>, M. Kadler<sup>3,4,5</sup>, J. Tueller<sup>2</sup>, A. Marscher<sup>6</sup>, E. Ros<sup>7</sup>, A. Zensus<sup>7</sup>, Y.Y. Kovalev<sup>7,8</sup>, M. Aller<sup>9</sup>, H. Aller<sup>9</sup>, J. Irwin<sup>9</sup>, J. Kerp<sup>10</sup>, S. Kaufmann<sup>11</sup>

## ABSTRACT

We present an analysis of the 101 ks, 2007 *Suzaku* spectrum of the LINER galaxy NGC 1052. The 0.3 – 10 keV continuum is well-modeled by a power-law continuum modified by Galactic and intrinsic absorption, and exhibits a soft, thermal emission component below 1 keV. Both a narrow core and a broader component of Fe-K $\alpha$  emission are robustly detected at 6.4 keV. While the narrow line is consistent with an origin in material distant from the black hole, the broad line is best fit empirically by a model that describes fluorescent emission from the inner accretion disk around a rapidly rotating black hole. We find no direct evidence for Comptonized reflection of the hard X-ray source by the disk above 10 keV, however, which casts doubt on the hypothesis that the broad iron line is produced in a standard accretion disk. We explore other possible scenarios for producing this spectral feature and conclude that the high equivalent width and full width half maximum velocity of the broad iron line ( $v \geq 0.37c$ ) necessitate an origin within  $d \sim 8r_g$  of the hard X-ray source. Based on the confirmed presence of a strong radio jet in this source, the broad iron line may be produced in dense plasma at the base of the jet, implying that emission mechanisms

---

<sup>1</sup>NPP Postdoctoral Fellow (ORAU); NASA's GSFC, mail code 662, Greenbelt MD 20771 USA

<sup>2</sup>NASA's GSFC, mail code 662, Greenbelt MD 20771 USA

<sup>3</sup>Dr. Karl-Remeis-Sternwarte, Astronomisches Institut der Universität Erlangen-Nürnberg, Sternwartstr. 7, 96049 Bamberg, Germany

<sup>4</sup>CRESST/NASA Goddard Space Flight Center, 662 Greenbelt, MD 20771 USA

<sup>5</sup>Universities Space Research Association, 10211 Wincopin Circle, Suite 500 Columbia, MD 21044 USA

<sup>6</sup>Institute for Astrophysical Research, Boston University, 725 Commonwealth Ave., Boston, MA 02215 USA

<sup>7</sup>MPIfR, Postfach 2024, D-53010 Bonn, Germany

<sup>8</sup>Astro Space Center of Lebedev Physical Institute, Profsoyuznaya 84/32, 117997 Moscow, Russia

<sup>9</sup>Department of Astronomy, University of Michigan, 500 Church St., Ann Arbor, MI 48109 USA

<sup>10</sup>Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

<sup>11</sup>Landessternwarte, Universität Heidelberg, Königstuhl 12, D-69117 Heidelberg, Germany

in the centralmost portions of active galactic nuclei are more complex than previously thought.

*Subject headings:* galaxies:individual – accretion, accretion disks – black holes – galaxies:nuclei – X-rays:spectra

## 1. Introduction

Since their first identification as an interesting class at optical wavelengths (Heckman 1980), the nature of Low Ionization Nuclear Emission Regions (LINERs) has remained elusive. Many such regions are associated with active galactic nuclei (AGN), though their X-ray luminosities are highly sub-Eddington and way below those of their Seyfert galaxy counterparts. The reason for this modest nuclear energy output remains unknown: possibilities include heavy absorption of the nuclear emission, comparably inefficient radiation of the accretion flow, or some combination of both.

The notion of an advection-dominated accretion flow (ADAF) may apply in LINERs, wherein the ions lose thermal contact with the electrons and only a small fraction of the dissipated energy is radiated (Rees et al. 1982; Narayan & Yi 1994). Much controversy remains as to whether this picture is correct (e.g., Guainazzi et al. (2000)), and the debate has become more pointed since the report by Ho et al. (1997) that some 20 – 30% of all galaxies are members of the LINER class.

X-ray spectroscopy can help in determining whether the accretion disk is in an ADAF state or not. A LINER galaxy harboring an AGN produces a characteristic hard X-ray continuum. If the accretion flow close to the black hole conforms to the conventional thin disk archetype (Shakura & Syunyaev 1973), the cool gas orbiting in the inner disk should intercept and reprocess a significant fraction of the continuum, also producing the Fe-K $\alpha$  line (among many fluorescent emission lines from other species as well). This spectral feature should have a broadened, skewed profile created by the relativistic properties of the spacetime close to the black hole, and a significant Comptonized reflection component should be seen in the continuum above 10 keV as well. If, however, the accretion flow in this region takes the form of an ADAF, the ion temperature is so high that the iron ions should be totally stripped of electrons. Furthermore, the trapped energy will puff up the disk, reducing its density and optical depth. Both of these conditions render the reflected emission, including the emission of Fe-K $\alpha$ , virtually non-existent in the inner accretion disk. In this case, only contributions to the Fe-K $\alpha$  line from farther out (e.g., from the outer disk, broad line region or putative molecular torus) will manifest, and the line will have a much narrower profile as a result.

The LINER galaxy NGC 1052 is an interesting case in which to study the dynamics of the accretion flow. The AGN is housed by a nearby elliptical host galaxy with a redshift of  $z = 0.0049$

(Knapp et al. 1978), implying a distance of 20.7 Mpc using WMAP cosmology. The source has a well-studied LINER optical spectrum (e.g., Gabel et al. (2000)), and is also reasonably bright in the X-ray band for a LINER, with an average 2 – 10 keV flux of  $F_{2-10} \sim 5 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ . The source is known to be heavily absorbed based on the presence of H<sub>2</sub>O megamasers in its core (Braatz et al. 1994), which lie along the direction of a radio jet also present in the galaxy (Claussen et al. 1998). A prominent, modestly broad Fe-K $\alpha$  line has been observed in the source by *ASCA* (Weaver et al. 1999), *XMM-Newton*, *Chandra* (Kadler et al. 2004b) and *BeppoSAX*. This spectral feature, coupled with detailed sub-parsec VLBI observations of the jet (Kadler et al. 2004a,b), may offer a unique opportunity to study the connection between the accretion flow and jet production in AGN.

A link between accretion and jet activity in AGN has been noted in the radio galaxies 3C 120 (Kataoka et al. 2007; Marscher 2006; Marscher et al. 2002) and 3C 111 (Marscher 2006). While Marscher et al. have noted dips in the X-ray light curves of these sources correlated with ejections of superluminal knots in the radio jet, suggesting that a portion of the inner disk is disturbed during an ejection event, these results fall short of mapping out the physical structure of the disk at the time. However, Kataoka et al. (2007) observed a broad Fe-K $\alpha$  line in 3C 120 with *Suzaku* in both the high and low flux states of the source. This discovery demonstrates the viability of studying the accretion flow and the jet activity of radio-loud AGN simultaneously, and will hopefully pave the way for new insights into the fundamental link between these two processes.

Here we discuss the analysis of our 2007 *Suzaku* observation of NGC 1052, which offers the highest quality X-ray spectrum of this galaxy to date and gives us an unprecedented view of the 0.5 – 35 keV energy range. NGC 1052 is one of a small handful of AGN sufficiently bright in both the X-ray and radio bands for tests of a jet/disk connection. Additionally, it is the only source for which we will have a full package of multi-epoch X-ray spectral data, X-ray and radio light curves, and VLBA imaging within a common time window and with an analysis by a single collaborative group. A forthcoming paper will provide full details of this ongoing campaign, placing our *Suzaku* results in the context of prior X-ray and ongoing radio analysis of this source (Kadler et al., in prep.).

The X-ray spectrum of NGC 1052 is quite flat and appears to be dominated at lower energies by extended thermal and/or photoionized emission (Weaver et al. 1999; Kadler et al. 2004a). The source of this emission is likely the interaction of the jet with the surrounding ISM (Kadler et al. 2004a). Radio continuum and *RXTE* monitoring both show that the source varies strongly on time scales of weeks to months. The 2 – 10 keV flux, in particular, varies between  $\sim 4 - 9 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  (Kadler et al., in prep.), and though there are large uncertainties in the *RXTE* data, the spectrum appears to switch from soft to hard states, perhaps analogous to the high/soft vs. low/hard states observed in Galactic microquasars and black hole binaries (Fender et al. 2005).

Our *Suzaku* observation of NGC 1052 and the data reduction will be detailed in Section 2. Our approach to the spectral fitting of the time-averaged data and the results of this analysis for both the XIS and the HXD/PIN instruments are presented in §3. We examine the Fe-K $\alpha$  region at length in §3 as well and discuss variability over the course of the observation in §4. Our results and conclusions comprise §5.

## 2. Observations and Data Reduction

We observed NGC 1052 for a total of  $\sim 101$  ks from 16-18 July 2007 with *Suzaku*. Reduction of the *Suzaku* data followed the procedures outlined in §4.7 and §4.8 of the *Suzaku* ABC Guide, available online<sup>1</sup>. Calibration files used were the latest version as of the time of this writing (September 2008). This calibration was incorporated into the X-ray Imaging Spectrometer (XIS) data reduction using the `xispi` task on the unfiltered data. Once cleaned, the event files for the XIS0, XIS1 (XIS2 became defunct as of Nov. 2006) and XIS3 detectors were loaded into XSELECT for reprocessing. Images were extracted that enabled identification of source (a circle  $\sim 250''$  in radius) and background regions. After spatial filtering was applied, spectra and light curves were extracted for each of the remaining three detectors. Response matrices and ancillary response files were generated using the `xisresp` task, and the spectra, backgrounds and responses for the front-illuminated chips (XIS0 and XIS3) were co-added to increase the signal-to-noise using `addascaspec`. The spectra, backgrounds and responses were then rebinned by a factor of eight (512 channels) using the `rbnpha` and `rbnrmf` tasks to speed spectral fitting. Finally, the spectra, backgrounds and responses were grouped together using the `grppha` task. Only spectral channels with a minimum of 20 cts/bin were used for fitting in order to ensure the validity of the chi-squared statistic. This corresponded to an energy range of 0.7 – 9 keV for the XIS0+3 data, 0.5 – 7 keV for the XIS1 data and 12 – 35 keV for the HXD/PIN data.

*Suzaku* did observe NGC 1052 with the silicon diode PIN instrument of the Hard X-ray Detector (HXD), but the GSO crystal scintillator instrument did not have enough counts to provide useful data. The PIN data were reduced using the “tuned” calibration files for the non-X-ray background (NXB) and the response files that had been generated for the period surrounding the observation. These files were downloaded from the *Suzaku* Guest Observer Facility (GOF) for use. We first needed to ensure that only the portion of the NXB coincident with our observation was used in processing, so we merged the good-time interval (GTI) of the NXB with that of our screened event file to yield a common GTI using `mgtime`. With XSELECT, we then filtered the data and background using this common GTI and extracted our spectra for the source and NXB.

---

<sup>1</sup><http://heasarc/docs/suzaku/analysis/abc/node1.html>

We corrected for dead time in the observation using the `hxdtdcor` task, and increased the exposure time of the NXB data by a factor of ten to compensate for the inflated event rate, which had been similarly multiplied in an effort to suppress Poisson noise. After reduction and filtering, the combined XIS+PIN spectrum had 22172 cts from 0.3 – 60 keV.

The cosmic X-ray background (CXB) has not been taken into account in PIN observations, so to model it appropriately, we simulated the CXB spectrum in `XSPEC` (Arnaud 1996) using a power-law model of  $\Gamma = 1.29$  with a high energy cutoff of `foldE`= 40 keV (initial parameters are given in the ABC Guide). We then folded this model through the response file for the PIN flat field for an exposure time of  $10^6$  s to generate the simulated CXB data. This spectrum was then read back into `XSPEC` with the PIN response for our observation and the original model was refitted over 12 – 35 keV to yield the actual parameters of the CXB during this time. In our case, the simulated CXB spectrum contributed a count rate of  $\sim 2.28 \times 10^{-2}$  cts  $s^{-1}$  to the X-ray background from 12 – 35 keV. In modeling the PIN data with `XSPEC`, we applied the best-fit CXB parameters as a constant with each fit. The CXB followed a power-law form with a high-energy cutoff:  $\Gamma = 1.30$ ,  $N_{po} = 8.13 \times 10^{-4}$  ph  $cm^{-2}$   $s^{-1}$ , `foldE` = 40.53 keV.

### 3. Spectral Analysis

We began our analysis of the 2007 *Suzaku* spectrum of NGC 1052 by examining the data from the operational XIS detectors. We use the 0.5 – 9.0 keV data to extract information about the underlying continuum, any soft excess emission, the Fe-K line complex and any intrinsic absorption in the system. NGC 1052 is thought to exhibit evidence for an AGN disk (the broad Fe-K $\alpha$  line) and jets (radio observations), as well as substantial, extended soft emission. As such, we expected the spectrum to be a conglomerate of all of these features since *Suzaku* lacks the spatial resolution necessary to physically separate them.

We also analyzed the HXD/PIN spectrum from 12 – 35 keV in order to learn more about the nature of the continuum and to constrain the amount of reflection seen in the source. Because a broad iron line has been observed in NGC 1052 in the past (Ros et al. 2007), we expected that residual emission above a power-law should remain at high energies due to the presence of the so-called ‘‘Compton Hump.’’ This feature has been noted in many AGN with broad iron lines (for a review, see (Reynolds & Nowak 2003)), representing Compton down-scattering of the hard X-ray photons by the material in the disk. This Compton hump typically peaks at  $\sim 20 - 30$  keV, and is thought to go hand-in-hand with the presence of a broad iron line, as both spectral features arise from the same physical process of reflection.

### 3.1. XIS Continuum

Before the iron line region could be considered, it was necessary to first accurately model the underlying continuum of NGC 1052. We began by ignoring the energy range where the iron line is thought to be important (3 – 7 keV) and modeled the 0.5 – 3 and 7 – 9 keV energy bands with a simple power-law modified by Galactic photoabsorption ( $N_{\text{H}} = 2.83 \times 10^{20} \text{ cm}^{-2}$ , as per Kalberla et al. (2005)). The result was a poor fit:  $\chi^2/\text{dof} = 840/361$  (2.33), where “dof” indicates the number of degrees of freedom of the model. Large residuals remained at both high and low energies (see Fig. 1), and a very hard photon index for the power-law component was seen:  $\Gamma = 0.34$ . To account for differences in calibration between the XIS0+3 and the XIS1 detectors, we multiplied each data set by a constant for normalization. This constant was held at 1.00 for XIS0+3, and achieved a best-fit value of 0.94 for the XIS1 data.

Especially noteworthy is an apparent soft excess clearly visible below  $\sim 1.5$  keV (see Fig. 1). To mitigate this residual feature we incorporated a thermal `mekal` component (Mewe et al. 1985). The fit improved dramatically:  $\chi^2/\text{dof} = 381/358$  (1.06), with  $kT \sim 0.6$  keV and a solar abundance for the `mekal` component. This component has been noted in previous X-ray analyses of NGC 1052 by Kadler et al. (2004b) with *Chandra* and Weaver et al. (1999) with *ASCA* (hereafter W99), and the temperature and flux obtained from our *Suzaku* fit are comparable to both values in both previous works. This thermal emission most likely originates in extended emission around the nucleus from the interaction of the jets with the ISM (Kadler et al. 2004b). Residuals still remained at low energies, however, so we added in a second `mekal` component (with an abundance equal to that of the first) to try to account for these features. Although marginal improvement in the overall goodness-of-fit was seen ( $\Delta\chi^2/\Delta\text{dof} = -12/2$ ), the addition of this second thermal component rendered the parameters of both `mekal` components unconstrained in an error analysis. We therefore removed the second thermal component. Though modest residuals remained at select energies, no statistically significant improvement in fit was achieved with the addition of discrete Gaussian components, as one might expect in a photoionized plasma, or with allowing the abundance of the `mekal` component(s) to vary. Alternatively, we also tried to account for this excess soft emission with a second power-law, with and without further absorption. Both models exhibited significantly poorer fits than that of the single `mekal` component.

Because NGC 1052 is also classified as a Seyfert 2 galaxy, it is reasonable to assume that some intrinsic absorption exists in or around the nuclear region. Such absorption could play a significant role in the spectral curvature seen, so it must be properly accounted for in the continuum model. We employed a partial-covering absorber (`zpcfabs`) intrinsic to the source to model this component, applying it only to the power-law and other components of nuclear AGN emission and not to the extended `mekal` components. A statistically significant improvement in fit was seen:  $\chi^2/\text{dof} = 335/356$  (0.94), with  $N_{\text{H}} = 1.08 \times 10^{23} \text{ cm}^{-2}$  and a covering fraction of 84%. With the




Fig. 1.— The XIS spectrum of NGC 1052 fit with a power-law modified by Galactic photoabsorption over the 0.5 – 3 and 7 – 9 keV bands uninfluenced by the fluorescent Fe-K $\alpha$  line. The fit is poor:  $\chi^2/\text{dof} = 840/361$  (2.33). Note especially the large residuals at both the hard and soft ends. Black crosses represent the combined XIS0+3 spectrum while the black line represents the model. Red crosses and line represent the XIS1 data and model. The lower panel of the figure shows the data-to-model residuals, with the green line representing a perfect fit with a data-to-model ratio of one.

inclusion of this intrinsic absorption, the power-law photon index increased to  $\Gamma = 0.80$ . This was a significant increase in photon index from the fit without intrinsic absorption.

Physically, this continuum model and its parameters are roughly comparable to those which were fit for the 1996 *ASCA* observation of the source (W99), although the photon index of the power-law is considerably harder and the intrinsic absorption less dense in this 2007 observation. A model with a dual neutral absorber modifying intrinsic plus scattered power-law components as in W99 did not improve the fit, statistically; moreover, the parameter values of the second power-law and absorber could not be constrained.

Even after modeling the neutral absorption in the system, some residual spectral curvature still remained around 2 – 3 keV, indicating that some further component of absorption likely remained unmodeled. We applied the simple ionized absorber model `absori` (Zdziarski et al. 1995) to the nuclear emission and noted a marked visual improvement to the fit from 2 – 3 keV, as well as a small but significant improvement in the global goodness-of-fit to  $\chi^2/\text{dof} = 325/354$  (0.92). This absorber was reasonably thick and moderately ionized, with a column density of  $N_{\text{H}} = 1.37 \times 10^{22} \text{ cm}^{-2}$  and an ionization parameter of  $\xi = 68 \text{ erg cm}^{-1} \text{ s}^{-1}$ . With the addition of this component, the power-law photon index increased to  $\Gamma = 1.50$ . This value lies within reasonable, expected physical limits for the X-ray continuum of an AGN. Our base continuum model therefore included a power-law modified by intrinsic absorption from both a cold, patchy absorber and a moderately ionized absorber of smaller column density, along with extended thermal emission. All model components were affected by Galactic photoabsorption.

### 3.2. The Fe-K $\alpha$ Line

Having fit the continuum, we then took the energies from 3 – 7 keV back into consideration and attempted to model the prominent residuals remaining, highlighted by a strong emission feature centered at 6.4 keV in the rest frame that was assumed to be neutral Fe-K $\alpha$  (see Fig. 2, top). We began by holding the continuum parameters constant, except for normalizations and the power-law photon index, and refitting. The strong residual at 6.4 keV remained, appearing to have a broadened red wing associated with it that extended down to  $\sim 4$  keV. For this fit,  $\chi^2/\text{dof} = 711/631$  (1.13).

To account for the strong emission feature of 6.4 keV, we initially assumed some core contribution from neutral Fe-K $\alpha$  outside of the inner accretion disk in distant material. This would produce a relatively narrow emission line, being outside the region where relativistic broadening becomes important. Inserting such a line ( $\sigma = 5 \text{ eV}$ , intentionally less than the resolution of the XIS in order to make the line truly narrow) into the model yielded a vast improvement in fit:  $\chi^2/\text{dof} = 584/629$  (0.93), with  $E = 6.40 \pm 0.01 \text{ keV}$ , indicative of neutral iron, and  $EW = 111 \text{ eV}$ .




Fig. 2.— *Top*: The residual emission feature remaining after fitting the continuum of the NGC 1052 *Suzaku*/XIS data, as per §3.1. Note the prominent peak centered at 6.4 keV in the rest frame and the excess emission extending down to  $\sim 4$  keV.  $\chi^2/\text{dof} = 711/631$  (1.13). *Middle*: Residuals remaining after fitting only the narrow Fe-K $\alpha$  core. Evidence for a broader feature is clear.  $\chi^2/\text{dof} = 584/629$  (0.93). *Bottom*: Residuals remaining after a `laor` line (Laor 1991) was used to model the broad Fe-K $\alpha$  emission line as well as the narrow core.  $\chi^2/\text{dof} = 761/792$  (0.96).

The broad residuals centered on 6.4 keV still remained, however (see Fig. 2, middle).

We attempted to model these broad residuals three separate ways: with a broad Gaussian, with a `diskline` component representing emission from the inner disk around a Schwarzschild black hole (Fabian et al. 1989), and with a `laor` component representing emission from the inner disk around a maximally-spinning Kerr black hole (Laor 1991). Each line is centered at 6.4 keV in the rest frame, corresponding to neutral Fe-K $\alpha$ . The Gaussian component improved the fit to  $\chi^2/\text{dof} = 571/627$  (0.91), and was quite broad:  $\sigma = 747$  eV. Its equivalent width of  $EW_{\text{broad}} = 201$  eV was slightly smaller than that of previous observations, e.g., W99, in which  $EW_{\text{broad}} \sim 300$  eV. Relaxing the energy constraint of the line resulted in  $E = 5.41 - 5.88$  keV, with  $\sigma = 0.78 - 1.30$  keV and  $EW_{\text{broad}} = 377$  eV, though significant residuals still remained on the red wing of the line. Because this is a purely phenomenological model, we turned to `diskline` and `laor` in the hope of achieving more physical parameter constraints.

The `diskline` model yielded  $\chi^2/\text{dof} = 574/625$  (0.92), which is not significantly different from the Gaussian fit, statistically, though it did account for more of the residuals below 6 keV. VLBI observations of the radio jet on sub-parsec scales have constrained the inclination angle of the accretion disk to be between  $57 - 72^\circ$  relative to our line of sight (Kadler et al. 2004b), so we applied these constraints to the model. Our best fit yielded an accretion disk emissivity index of  $\alpha = 1.69 - 3.69$  (where the disk emissivity  $\epsilon \propto r^{-\alpha}$ ) and an inner disk radius of  $r_{\text{in}} < 26 r_{\text{g}}$  (where  $r_{\text{g}} = GM/c^2$ ), but the inclination angle could not be further constrained. We held the outer radius constant at  $400 r_{\text{g}}$ , since it was unable to be constrained and our emissivity index dictated that this radius safely encompasses all of the relevant emission from the disk.

The `laor` model provided the best statistical fit, at  $\chi^2/\text{dof} = 561/625$  (0.90), and also appeared to be the most effective at reducing the residuals between 4 – 7 keV (see Fig. 2, bottom). Keeping the same disk inclination constraints mentioned above for the `diskline` model, we again could not further constrain this parameter, but we noted that the emissivity index of the disk was fairly centrally concentrated at  $\alpha = 1.53 - 5.10$ , while the inner radius of emission is only mildly constrained within  $r_{\text{in}} = 8.93 - 45.10 r_{\text{g}}$ . The equivalent width of the broad iron line here was  $EW_{\text{broad}} \sim 185$  eV. Interestingly, when we relaxed the constraint on the inclination angle of the disk we obtained a best-fit value of  $i = 45 \pm 5^\circ$ : this is roughly a  $\sim 2\sigma$  deviation from the radio constraints.

The `laor` fit was a slight improvement over both the Gaussian and `diskline` models, though it should be noted that the uncertainty of the  $r_{\text{in}}$  parameter effectively renders it impossible to meaningfully distinguish between the `diskline` and `laor` models based on the Fe-K $\alpha$  profile alone. If the `laor` model fit could better constrain the inner disk radius to  $r_{\text{in}} < 6 r_{\text{g}}$ , it would suggest that the black hole at the center of NGC 1052 is rapidly rotating, as we would expect for a source that powers relativistic jets. We cannot make this statement unequivocally based on our

data, however. The data fit with this `laor` model are shown in Fig. 3. The final best-fit model parameters, incorporating the HXD/PIN data as well, are presented in the following Section within Table 1 as Model 1.

Absorption has often been suggested as an alternative to broad iron emission when attempting to model enhanced spectral curvature in the 4 – 7 keV range (e.g., Kinkhabwala (2003)). To address this possibility, we compared our best `laor` fit above to a model in which we removed the broad iron line and replaced it with a second absorbed power-law component. The resulting fit yielded  $\chi^2/\text{dof} = 582/627$  (0.93), as opposed to the  $\chi^2/\text{dof} = 561/625$  (0.90) achieved with the final `laor` model, and left significant residuals around 6.4 keV and below. Moreover, the parameter values for both the partial-covering absorber and the power-law components are not constrained in the fit. The models employing relativistic disk emission lines unquestionably provide better statistical and visual fits to the data.

### 3.3. Addition of the HXD/PIN Data

Adding the HXD/PIN data on to the combined XIS0+3 and XIS1 spectra, we noticed that the signal-to-noise ratio deteriorated significantly above  $\sim 35$  keV. As such, we restricted our energy range for this detector to 12 – 35 keV, applying a constant normalization factor as we did with the XIS1 data set. In this case, we used a parameter value of 1.09 in keeping with the value expected for the HXD nominal pointing position.

We considered the combined data refitted with Model 1, including no reflection component above 9 keV. Fig. 4 shows the data, model and residuals from this fit, which yields  $\chi^2/\text{dof} = 660/685$  (0.96). The inclusion of the high-energy data left the 2 – 9 keV parameters effectively unchanged from their values in the fit without the PIN data. If a significant flux contribution from reflection is present, as one might assume if the broad iron line has an origin in a relativistic accretion disk (see discussion in §3), we would expect to see some excess in the emission over 9 keV relative to Model 1. No such feature was detected, however.

Table 1 shows the full 0.5 – 35 keV fit with 90% confidence errors.

### 3.4. Alternative Continuum Models

Though no evidence for Comptonized reflection is apparent to the eye above 9 keV, it is nonetheless important to try to place upper limits on its possible contribution to the overall spectrum. With this in mind, we utilized two popular, well-regarded models of reflection from an




Fig. 3.— The best-fitting model for the XIS data. This model includes a power-law continuum modified by intrinsic absorption from a partial-covering neutral material as well as a mildly ionized gas, extended thermal emission, and both broad and narrow components of neutral Fe-K $\alpha$ . The spectrum is also modified by Galactic photoabsorption.  $\chi^2/\text{dof} = 561/625$  (0.90). The red, black and green colors/lines are the same as in Fig. 1.




Fig. 4.— HXD/PIN data included from 12 – 35 keV and refitted with Model 1, which does not include any type of continuum disk reflection. Inclusion of this data does not significantly alter the model parameter values from the 0.5 – 9 keV fit. Though a broad line is still required at 6.4 keV, no accompanying evidence of reflection from the disk above 9 keV is seen. Red and black colors follow the designations of Fig. 1. In the top panel, the green crosses and line represent the PIN data and model, respectively. In the bottom panel the horizontal green line still represents a data-to-model ratio of one, while the green crosses show the residuals between the PIN data and model.  $\chi^2/\text{dof} = 660/685$  (0.96).

accretion disk around a black hole in an attempt to quantify the amount of reflection that might be present in the spectrum of NGC 1052.

The `pexrav` model of Magdziarz & Zdziarski (1995) is one of the most widely used public models for AGN emission which incorporates reflected continuum from a neutral accretion disk. While it represents a more physical model than the simple power-law alone, it does not include line emission from the disk (e.g., Fe-K $\alpha$ ) and it is difficult, in practice, to constrain the three most important parameters simultaneously: photon index, reflection fraction and disk inclination angle.

Adding a `pexrav` component to our power-law continuum (and keeping the inclination angle of the disk constant at the `laor` value, the power-law cutoff at 100 keV, and the abundances fixed at their solar values), we found  $\chi^2/\text{dof} = 660/683$  (0.97), only a very marginal change in fit from that of Model 1 over the 0.5 – 35 keV energy range. The photon index fit between  $\Gamma = 1.45 - 1.56$  while the reflection fraction was constrained to be quite small:  $R < 0.01$ . As we expect, based on the residuals from fitting Model 1, reflection is not a significant contributor to the overall observed spectrum. The emissivity index of the `laor` line for Fe-K $\alpha$  had very similar constraints to those found in Model 1:  $\alpha = 1.74 - 4.56$ . Also, to 90% confidence, the inner radius of emission in the disk was constrained to  $r_{\text{in}} \leq 45 r_{\text{g}}$ , as we found in Model 1. The equivalent width of the broad Fe-K $\alpha$  line in this model also remained consistent with that found in Model 1, at  $EW = 187$  eV. The `pexrav` model will hereafter be known as Model 2.

The `reflion` model of Ross & Fabian (2005) is another widely available code used to describe the reflected spectrum of an accretion disk around a black hole. The advantage of this model over the `pexrav` model is that it is self-consistent: fluorescent emission lines from many ionized species are included in addition to the continuum, most notably those of the Fe-K complex. The ionization parameter, incident power-law photon index and iron abundance are specified by the user. The disadvantage is that photon indices below  $\Gamma = 1$  are outside the allowed parameter range of the model, and there is not a parameter for the reflection fraction, as in `pexrav`. The user is therefore obliged to estimate it using, e.g.,

$$R_{\text{refl}} = \frac{K_{\text{refl}} F_{\text{refl}}}{K_{\text{po}} F_{\text{po}}} \left( \frac{\xi}{30} \right)^{-1}. \quad (1)$$

Here  $K_{\text{refl}}$  and  $K_{\text{po}}$  denote the normalizations of the `reflion` and power-law components from our best-fit spectral model,  $\xi$  is the best-fit ionization parameter characterizing the disk reflection (where  $\xi = 30$  denotes a neutral disk), and  $F_{\text{refl}}$  and  $F_{\text{po}}$  are the total fluxes contained in the `reflion` and power-law components for unity normalization over the full wavelength range (0.001 – 1000 keV).

We began by removing the `laor` line from the Model 1 fit and adding in a `reflion` component, with the iron abundance held at the solar value and the photon index linked to that of the

power-law. As expected, this fit mimicked that of the continuum plus a narrow 6.4 keV Gaussian, leaving residuals on the low-energy end of the line. Statistically,  $\chi^2/\text{dof} = 679/687$  (0.99). Allowing the iron abundance to fit freely yielded a small improvement in fit, with  $\chi^2/\text{dof} = 667/686$  (0.97) with  $\text{Fe}/\text{solar} < 3.53$ , but doing so rendered the ionization parameter and normalization unconstrained. As such we elected to keep the iron abundance constant at the solar value.

If NGC 1052 harbors a rapidly spinning black hole, as our best `laor` fit seems to indicate, we must take into account the relativistic effects on emission originating from the inner accretion disk. We incorporated this smearing of spectral features using the `laor`-based convolution model `kdblur`. Our best fit was  $\chi^2/\text{dof} = 660/684$  (0.96), comparable to our best `laor` and `pexrav` fits. However, we were unable to constrain the emissivity index of the accretion disk, though we did obtain constraints on the inner radius of emission that are consistent with those found in the `laor` model:  $r_{\text{in}} < 45 r_{\text{g}}$ . The estimated reflection fraction is quite low at  $R_{\text{refl}} < 0.006$ , consistent with the results of the `pexrav` model fit and partly attributable to the moderately high ionization parameter ( $\xi = 109$ ) in the `reflion` model. The `kdblur(reflion)` model will hereafter be known as Model 3. For comparison, the data-to-model ratios and model components for the Models 1-3 fits are plotted in Fig. 5, and their best-fit parameters are listed in Table 1.

Models 2-3 constrain the amount of reflection from the disk to be  $R_{\text{refl}} < 0.01$ , consistent with the lack of a discernible Compton hump above 10 keV in the HXD/PIN data. Moreover, as we describe in detail later in §5, if the broad iron line results from fluorescence in the disk caused by incident hard X-ray emission, then the equivalent width of the Fe-K $\alpha$  line should be directly related to the normalization of the reflection component in our model as compared to that of the power-law. Yet this is not the case in our data. Based on the low normalizations for reflection seen in both Model 2 and Model 3, we should expect a broad Fe-K $\alpha$  equivalent width of  $EW \sim 80$  eV, much less than the observed  $EW \sim 185$  eV in both Models 1-2. In other words, our broad iron line does not appear to have a correspondingly strong amount of continuum reflection associated with it. It is possible that this “missing” reflection is actually present to some greater degree in the data, but is perhaps swamped out by the power-law continuum. It is also possible that the reflection is simply absent in this source. This scenario presents several intriguing possibilities for physical interpretations, which we discuss in §5.

#### 4. Timing Analysis and Variability

The spectral results presented thus far have focused solely on the time-averaged data from the XIS and HXD/PIN instruments, but because many types of AGN have been observed to vary on timescales ranging from hours to years, we address the spectral and temporal variability of




Fig. 5.— The relative contributions of each model component in the three relevant fits to the NGC 1052 XIS and PIN data: `laor` (top, Model 1), `pexrav+laor` (middle, Model 2) and `kdblur(reflion)` (bottom, Model 3). Each model presents its own strengths and weaknesses, as discussed in the text. Models are extended to 100 keV to illustrate their projected contributions at higher energies. Red represents the power-law (affected by Galactic and intrinsic absorption), green represents the thermal mekal component, Gaussian emission lines are in dark blue and reflection features such as the `laor` line (top), the `pexrav` component (middle) and the `kdblur(reflion)` model (bottom) are in light blue. The `laor` line in Model 2 is in magenta.

Component	Parameter Value	Model 1	Model 2	Model 3
mekal	$kT$ (keV)	$0.64^{+0.04}_{-0.04}$	$0.64^{+0.04}_{-0.04}$	$0.64^{+0.04}_{-0.04}$
	$K_{kT}$ (ph cm <sup>-2</sup> s <sup>-1</sup> )	$4.14^{+1.00}_{-1.02} \times 10^{-5}$	$4.14^{+0.74}_{-0.84} \times 10^{-5}$	$4.14^{+0.64}_{-0.57} \times 10^{-5}$
	$F_{kT}$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$9.00^{+2.17}_{-2.22} \times 10^{-14}$	$9.48^{+1.69}_{-1.92} \times 10^{-14}$	$8.78^{+1.35}_{-1.94} \times 10^{-14}$
absori	$\Gamma$	$1.50^{+0.07}_{-0.08}$	$1.50^{+0.06}_{-0.05}$	$1.50^{+0.05}_{-0.05}$
	$N_H$ ( $\times 10^{22}$ cm <sup>-2</sup> )	$1.37^{+0.54}_{-0.78}$	$1.37^{+0.27}_{-0.23}$	$1.45^{+0.63}_{-0.53}$
	$T$ (K)	$3 \times 10^4$	$3 \times 10^4$	$3 \times 10^4$
	$\xi$ (erg cm <sup>-1</sup> s <sup>-1</sup> )	$68^{+100}_{-31}$	$66^{+36}_{-46}$	$67^{+26}_{-33}$
zpcfabs	$N_H$ ( $\times 10^{22}$ cm <sup>-2</sup> )	$10.82^{+0.44}_{-0.77}$	$10.82^{+0.64}_{-0.94}$	$10.83^{+0.55}_{-0.87}$
	%cover	$0.84^{+0.03}_{-0.04}$	$0.84^{+0.02}_{-0.02}$	$0.85^{+0.02}_{-0.02}$
zpo	$\Gamma$	$1.50^{+0.07}_{-0.08}$	$1.50^{+0.06}_{-0.05}$	$1.50^{+0.05}_{-0.05}$
	$K_\Gamma$ (ph cm <sup>-2</sup> s <sup>-1</sup> )	$1.55^{+0.20}_{-0.24} \times 10^{-3}$	$1.55^{+0.21}_{-0.25} \times 10^{-3}$	$1.62^{+0.23}_{-0.35} \times 10^{-3}$
	$F_\Gamma$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$1.89^{+0.24}_{-0.29} \times 10^{-11}$	$1.79^{+0.24}_{-0.29} \times 10^{-11}$	$1.85^{+0.34}_{-0.32} \times 10^{-11}$
zgauss	$E$ (keV)	$6.40^{+0.01}_{-0.01}$	$6.40^{+0.01}_{-0.01}$	$6.40^{+0.01}_{-0.01}$
	$K_{\text{narow}}$ (ph cm <sup>-2</sup> s <sup>-1</sup> )	$1.18^{+0.19}_{-0.19} \times 10^{-5}$	$1.19^{+0.19}_{-0.19} \times 10^{-5}$	$1.26^{+0.15}_{-0.15} \times 10^{-5}$
	$F_{\text{narow}}$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$1.00^{+0.16}_{-0.16} \times 10^{-13}$	$9.54^{+1.54}_{-1.54} \times 10^{-14}$	$1.23^{+0.18}_{-0.18} \times 10^{-13}$
	$EW_{\text{narow}}$ (eV)	$111^{+18}_{-18}$	$111^{+18}_{-18}$	$121^{+16}_{-16}$
laor, kdblur	$\alpha_{\text{emis}}$	$2.40^{+2.14}_{-0.82}$	$2.40^{+2.17}_{-0.65}$	$0.95^{+9.05}_{-0.95}$
	$r_{\text{in}}$ ( $r_g$ )	$19.93^{+24.52}_{-10.04}$	$19.76^{+24.68}_{-19.76}$	$20.15^{+25.36}_{-20.15}$
	$i$ ( $^\circ$ )	$72^{+0}_{-15}$	$72^{+0}_{-15}$	$72^{+0}_{-15}$
	$K_{\text{broad}}$ (ph cm <sup>-2</sup> s <sup>-1</sup> )	$1.69^{+1.03}_{-0.69} \times 10^{-5}$	$1.71^{+0.97}_{-0.68} \times 10^{-5}$	—
	$F_{\text{broad}}$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$1.44^{+0.50}_{-0.59} \times 10^{-13}$	$1.38^{+0.78}_{-0.55} \times 10^{-13}$	—
	$EW_{\text{broad}}$ (eV)	$185^{+112}_{-75}$	$187^{+106}_{-74}$	—
pexrav, reflion	Fe/solar	—	1.0	1.0
	$\xi$ (erg cm <sup>-1</sup> s <sup>-1</sup> )	—	0.0	$109^{+2}_{-109}$
	$R_{\text{refl}}$	—	$0.01^{+0}_{-0.01}$	$5.75^{+0}_{-5.75} \times 10^{-3}$
	$K_{\text{refl}}$ (ph cm <sup>-2</sup> s <sup>-1</sup> )	—	$7.21^{+0}_{-7.21} \times 10^{-8}$	$1.30^{+1.71}_{-1.22} \times 10^{-7}$
	$F_{\text{refl}}$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	—	$4.04^{+0}_{-4.04} \times 10^{-18}$	$4.24 \times 10^{-13}$
	$\chi^2/\text{dof}$	660/685 (0.96)	660/683 (0.97)	660/684 (0.96)

Table 1: Comparison of our three best-fit models for the 0.5 – 35 keV spectrum of NGC 1052. All components of Models 1-3 are modified by Galactic hydrogen absorption with  $N_H = 2.83 \times 10^{20}$  cm<sup>-2</sup>. “K” denotes the normalization value of a given component. Flux values indicate absorbed flux from 0.5 – 35 keV. Redshifts for the model components are held constant at the cosmological value for NGC 1052:  $z = 0.0049$ . Abundances not listed are held constant at their solar values. The inclination angle of the accretion disk used in the laor and kdblur models was constrained to fall within the radio observation uncertainties of  $i = 57 - 72^\circ$ . All errors listed are at 90% confidence for one interesting parameter.

NGC 1052 here. The 2007 *Suzaku* data are discussed below, as are the 1996 *ASCA* data of W99 for comparison. The X-ray results from intervening epochs will be addressed by Kadler et al. (in prep.).

#### 4.1. 2007 *Suzaku* Data

No significant ( $> 3\sigma$ ) changes were seen in the count rates of any of the XIS instruments or the PIN instrument over the course of the observation. Fig. 6 depicts the light curves from XIS 0, XIS 1 and XIS 3, as well as the combined XIS light curve vs. that of the PIN. While small variations in count rate (over  $\sim 10^4$  s timescales) were seen in both data sets, neither varied more than a factor of two from its baseline flux throughout the observation. This flux was roughly  $0.06 \text{ cts s}^{-1}$  for the PIN and  $0.38 \text{ cts s}^{-1}$  for the combined XIS data ( $0.16$  for XIS 0,  $0.14$  for XIS 3 and  $0.21$  for XIS 1).

As mentioned in the previous Section, the overall 2 – 10 keV flux did not vary significantly between Models 1-3, averaging to  $F_{2-10} = 5.37 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  (absorbed). This was split between the continuum and the Fe-K lines, where the continuum flux was  $F_{\text{cont}} = 4.86 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  and that of the combined broad and narrow Fe-K $\alpha$  lines was  $F_{\text{Fe}} = 5.10 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ . The average unabsorbed 2 – 10 keV flux at the distance of NGC 1052,  $F_{2-10} = 8.96 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ , works out to an intrinsic luminosity of  $L_{2-10} = 4.60 \times 10^{41} \text{ erg s}^{-1}$  using WMAP cosmology.

#### 4.2. Variation Between Epochs

We considered our best 2007 *Suzaku* results (Model 1) against those obtained in the 1996 *ASCA* observation by W99. Though these authors did not present a light curve for their data, they fit a time-averaged 2 – 10 keV flux of  $F_{2-10} \sim 8 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ , corrected for a dual absorber in the best fit. This differs only slightly from the unabsorbed flux of  $F_{2-10} \sim 8.96 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  we have obtained from our data in the present epoch. The neutral hydrogen absorbing column in our 2007 best fit is roughly a factor of three less dense than that of W99: whereas we find evidence for one intrinsic neutral absorber of  $N_{\text{H}} \sim 1.08 \times 10^{23} \text{ cm}^{-2}$ , in their best-fitting model W99 employ a two-zone absorption model with  $N_{\text{H}}(1) = 3 \times 10^{23} \text{ cm}^{-2}$  and  $N_{\text{H}}(2) = 5 \times 10^{22} \text{ cm}^{-2}$ . Our model contains a component of ionized absorption that the W99 model does not, however: the `absori` model representing this absorbing material has a column density of  $N_{\text{H}} = 1.37 \times 10^{22} \text{ cm}^{-2}$  and an ionization parameter of  $\xi = 68 \text{ erg cm}^{-1} \text{ s}^{-1}$ . Even with this additional layer the W99 model contains a higher absorbing column than our 2007 model.

The heavy absorption in W99 resulted in a power-law photon index split between intrinsic

Waveband (keV)	Model 1 ( $\text{erg cm}^{-2} \text{s}^{-1}$ )	Model 2 ( $\text{erg cm}^{-2} \text{s}^{-1}$ )	Model 3 ( $\text{erg cm}^{-2} \text{s}^{-1}$ )
0.5 – 2.0	$3.42 \times 10^{-13}$	$3.42 \times 10^{-13}$	$3.68 \times 10^{-13}$
	$3.42 \times 10^{-12}$	$3.42 \times 10^{-12}$	$5.31 \times 10^{-12}$
2.0 – 10.0	$5.53 \times 10^{-12}$	$5.25 \times 10^{-12}$	$5.33 \times 10^{-12}$
	$8.96 \times 10^{-12}$	$8.96 \times 10^{-12}$	$9.20 \times 10^{-12}$
10.0 – 60.0	$3.64 \times 10^{-11}$	$3.64 \times 10^{-11}$	$3.79 \times 10^{-11}$
	$3.68 \times 10^{-11}$	$3.68 \times 10^{-11}$	$3.82 \times 10^{-11}$

Table 2: Extrapolated model fluxes for the 2007 *Suzaku* observation of NGC 1052 in three wavebands. Absorbed fluxes are on the top line, unabsorbed fluxes are on the line below. For the softest energies the flux from XIS 1 was used, while the XIS 0+3 data was used for the 2 – 10 keV band. PIN data were used for the flux above 10 keV.




Fig. 6.— *Left*: Light curves from the three operational XIS instruments for the 2007 *Suzaku* observation of NGC 1052. *Right*: Light curve of the combined XIS data (black) as well as that of the PIN (red) over the course of the observation. All light curves shown are background-subtracted, including the CXB for the PIN data. Note the overall lack of significant variability of the source.

and scattered components, where a power-law photon index of  $\Gamma = 1.7$  was assumed and 11% was scattered, while 70% of the intrinsic flux was absorbed by the larger column and 30% by the smaller column. We attempted such a neutral dual-absorber fit (see §3.1-3.2) and found that the parameters of the second absorber could not be constrained, and that a single intrinsic neutral absorber and an ionized along with Galactic photoabsorption modifying a simple, albeit fairly hard power-law ( $\Gamma \sim 1.5$ ) provided an excellent fit to the continuum.

W99 also found evidence for a thermal `mekal` component to explain the soft excess noted below 2 keV, which was best modeled with  $kT = 0.53^{+0.34}_{-0.26}$  keV and a 0.1 – 2.0 keV flux of  $F_{\text{mekal}} = 5.8^{+2.5}_{-2.3} \times 10^{-14}$  erg cm<sup>-2</sup> s<sup>-1</sup>. This component was also detected by Kadler et al. (2004a) with *Chandra*, and we found a similar thermal component in the *Suzaku* data. Though the XIS detectors lose effective area rapidly below 0.3 keV, extrapolating Model 1 down to 0.1 keV yielded  $kT = 0.64 \pm 0.04$  keV and  $F_{\text{mekal}} = 9.38^{+2.27}_{-2.31} \times 10^{-14}$  erg cm<sup>-2</sup> s<sup>-1</sup> over the same energy range. Our results showed a slight increase in temperature and flux for the `mekal` component but were consistent with the 1996 results within error bars.

In the favored dual-absorber *ASCA* model of W99, a broad Fe-K $\alpha$  line component was not required in the fit and hence these authors found no definitive evidence for reflection from an optically-thick accretion disk in their data. They did require a narrow Fe-K $\alpha$  line, however: it was centered at 6.37 keV and has an intrinsic width of  $\sigma = 20$  eV, an equivalent width of  $270 \pm 120$  eV and a normalization of  $1.8 \pm 0.8 \times 10^{-5}$  ph cm<sup>-2</sup> s<sup>-1</sup>. In our Model 1 for the *Suzaku* data, we found that both broad and narrow components were necessary in order to eliminate the residuals between 4 – 7 keV. Our narrow line, with centroid energy constrained at  $6.4 \pm 0.01$  keV in the rest frame with an intrinsic width held at  $\sigma = 5$  eV, had an equivalent width of  $111 \pm 18$  eV and a normalization of  $1.18 \pm 0.19 \times 10^{-5}$  ph cm<sup>-2</sup> s<sup>-1</sup>. The presence of a broad component to the Fe-K $\alpha$  line likely rendered this narrow feature significantly weaker than its 1996 counterpart.

Our broad Fe-K $\alpha$  line at  $E = 6.4$  keV was best fit with a `laor` emission feature originating from the accretion disk around a rapidly rotating black hole, though the inner radius of emission within the disk was only broadly constrained to  $r_{\text{in}} < 45 r_g$ , as discussed in §3.2. This feature was stronger than the narrow Fe-K $\alpha$  line: its normalization was  $1.69^{+1.03}_{-0.69} \times 10^{-5}$  ph cm<sup>-2</sup> s<sup>-1</sup> and its equivalent width was  $185^{+112}_{-75}$  eV. Replacing the `laor` line with a broad Gaussian, we found that  $\sigma = 1.04 \pm 0.26$  keV, which translates to a lower-limit FWHM of  $v \sim 0.37c$  and implies an origin for this feature very close to the black hole. Though the fit of the `laor` line itself is suggestive of a broad, fluorescent Fe-K $\alpha$  origin in the inner accretion disk, the lack of reflected continuum flux above 10 keV argues against this interpretation, as we discuss below and in the following Section.

W99 considered a reflection-based `hrefl` model to fit the *ASCA* spectrum, but were unable to obtain meaningful constraints on the reflection parameters without making several assumptions. These included fixing the value of the power-law photon index of the central X-ray source to be  $\Gamma =$

1.7 and also fixing the inclination angle of the disk to  $i = 60^\circ$ . The latter was consistent with our own angle constraints based on the VLBI observations of NGC 1052, but our *Suzaku* observation indicated a somewhat harder photon index (see Table 1). For an assumed reflection fraction of  $R_{\text{refl}} = 1$  and the observed *ASCA* spectrum, W99 found that only a small fraction of the direct emission from the X-ray source was visible, leading them to infer that a thick absorbing column in the nucleus ( $N_{\text{H}} > 3 \times 10^{23} \text{ cm}^{-2}$ ) blocked the X-ray source from view, such that reflected emission dominated the spectrum. However, the equivalent width of the Fe-K $\alpha$  line,  $EW \sim 40 \text{ eV}$ , was over an order of magnitude smaller than the same line seen in other heavily obscured AGN. To account for this line with a reflection model would require an implausibly small iron abundance of  $Z_{\text{Fe}}/\text{solar} \leq 0.05$ . This reflection model was therefore argued against by W99.

The 2007 *Suzaku* data concur with this assessment of a lack of reflection: though Model 2 produced a comparable global goodness-of-fit to Model 1, the reflection fraction was found to be quite small at  $R_{\text{refl}} < 0.01$ . Attempting to model any reflection present with Model 3, we were unable to constrain the emissivity index of the accretion disk, though we did achieve similar constraints for the equivalent width of the narrow iron line and the photon index of the power-law component. Based on this spectral fitting, we conclude that in 2007, NGC 1052 does show evidence for a broadened iron line; this line is not particularly broad or strong, however, and is *not* accompanied by the expected Compton Hump above 10 keV. This may indicate that the line arises outside of the innermost portions of the disk and/or that reflection simply does not play a large role in creating the overall X-ray spectrum.

## 5. Discussion

We have performed detailed spectral fits to the XIS and HXD/PIN data from the 101 ks, 2007 *Suzaku* observation of the LINER galaxy NGC 1052. Three different spectral models were employed: (1) an absorbed power-law continuum with contributions from a soft thermal component along with narrow and broad Fe-K $\alpha$  lines; (2) the same model, but with an added component of reflected continuum emission from a neutral accretion disk (`pexrav`); (3) the same base model, again, but with a more advanced, self-consistent reflection model which includes fluorescent emission lines from the disk and takes ionization into account (`reflion`; see Ross & Fabian (2005) for details). In Model 3 we applied relativistic smearing using the `kdblur` model. Because fluorescent lines are included in `reflion`, there is no need for an additional broad iron line component as in Model 1. We summarize our findings below:

- The three models are quite similar in their statistical goodness-of-fit, though some of the disk reflection parameters in Models 2-3 prove especially difficult to constrain.

- The continuum is well described by a component with a power-law spectrum of photon index  $\Gamma \sim 1.5$ , thermal emission with  $kT \sim 0.64$  keV, a neutral intrinsic absorbing column of  $N_{\text{H}} \sim 10^{23} \text{ cm}^{-2}$  and a covering fraction of  $\sim 84\%$ , and an ionized intrinsic absorber with  $N_{\text{H}} = 1.37 \times 10^{22} \text{ cm}^{-2}$  and  $\xi = 68 \text{ erg cm}^{-1} \text{ s}^{-1}$ . These findings are relatively consistent with those of the 1996 *ASCA* observation of W99, though our photon index is harder and our absorbing column lower in density in 2007.
- As in the *ASCA* data, a narrow Fe-K $\alpha$  line is present:  $EW = 111 \text{ eV}$ .
- Strong evidence exists for a broadened Fe-K $\alpha$  emission component ( $EW = 185 \text{ eV}$ ). Though the best statistical fit is achieved with a `laor` line profile, however, the rather broad constraint on the inner radius of  $r_{\text{in}} < 45 r_{\text{g}}$  renders it impossible to distinguish between a spinning and non-spinning black hole.
- The PIN instrument did detect hard X-ray emission above 10 keV from NGC 1052. This emission is well fit by the power-law component of the 2 – 10 keV continuum and does not show any evidence for reflected emission via the Compton hump that is commonly associated with other AGN harboring broad iron lines, and expected based on the presence of the broad iron line if the two features arise from reflection onto a standard accretion disk. Neither of our reflection models found any statistical evidence for a strong reflection component.
- Though VLBI observations of the jet in NGC 1052 have constrained the orientation angle of the disk to be between  $i = 57 - 72^\circ$ , we find that removing this constraint in our fitting yields a best-fit inclination angle of  $i = 45 \pm 5^\circ$ , though this does not significantly improve the global goodness-of-fit. Nonetheless, this result does mark a  $2\sigma$  deviation from the radio results.

These findings motivate three main questions: (1) what is the physical origin of the broad iron line in our 2007 observation; (2) how is it that we detect a broad iron line yet no evidence for disk reflection above 10 keV, and (3) why does the `laor` line inclination, when kept as a free parameter in the fit, not agree with the disk inclination constraints established in radio observations of the jet in NGC 1052 on sub-parsec scales?

As it turns out, the answers to these questions may be closely related.

Broad iron lines are expected to be associated with pronounced Compton humps. Comptonized reflection originates via irradiation of the optically thick inner accretion disk by the hard X-ray source, perhaps in some sort of lamp-post geometry, e.g., the base of a jet (Miniutti et al. 2007). This same process will also produce fluorescent line emission from the disk, most notably the Fe-K $\alpha$  line, which will experience significant broadening and skewness by virtue of its origin from the spacetime proximal to the black hole (Reynolds & Nowak 2003). The absence of a

Compton Hump, however, casts doubt on the presence/contribution of reflection from the disk to the overall X-ray spectrum. From our `pexrav` model fit, we can estimate the equivalent width of the iron line we should expect if both the Compton hump and the broad line originate from disk reflection. Using the ratio of the normalization of the power-law to that of the reflected component in our fit, we infer that the iron line equivalent width should be  $EW \sim 80 \text{ eV}$ . This low value is highly inconsistent with the observed strength of the broad iron feature.

Relativistic disk smearing is not the only process by which this spectral line can be broadened, however. In what follows we consider a number of other physical scenarios, including: transition to an advection-dominated accretion flow (ADAF) in the inner disk, resonant scattering from the putative torus region, origin of the line in an outflow located in the broad line region (BLR), and Fe-K $\alpha$  produced by Cerenkov line-like radiation rather than fluorescence.

One explanation for the lack of an observed reflected continuum could be that the accretion disk transitions to a radiatively-inefficient state within some radius, e.g., Narayan & Yi (1994). In this ADAF picture, which has often been invoked as a potential explanation for low-luminosity sources (especially those with observed jet activity), the disk traps radiation rather than releasing it, puffing up to become optically thin and geometrically thick as its temperature and ionization state rises. Because of these properties, the gas would not be expected to produce much, if any, reflection, either in the form of a Compton hump or discrete emission lines. If a contribution to the broad iron line did arise from this region, we would expect it to be highly ionized, yet this is not seen in the data: our broad and narrow iron line components both have energies robustly less than 6.41 keV, suggesting that they arise from species at or below Fe XVIII. So while an ADAF may indeed be a viable explanation for the lack of a Compton hump in this source, it is likely not a significant contributor to the broadening of the Fe-K $\alpha$  feature.

The lack of a Compton hump and the mismatch between the unconstrained `laor` disk inclination and the VLBI value causes us to rule out a relativistic disk origin for the broad iron line. We also eliminate a possible origin in an ADAF due to ionization constraints and optical depth issues. Investigating other possible sources of iron emission in the central AGN leads us to consider the role of resonant scattering in the torus region, which is thought to contribute heavily to the flux of the narrow Fe-K $\alpha$  line. However, both the large equivalent width and the large FWHM velocity of the broad line component we observe (both substantially greater than the  $v \sim 2500 \text{ km s}^{-1}$  and  $EW \lesssim 100 \text{ eV}$  typically associated with the narrow core, e.g., Yaqoob & Padmanabhan (2004)) render this explanation implausible.

An AGN outflow originating in the BLR is explored as a possible explanation for the similar spectrum of NGC 7213 (Bianchi et al. 2008). Here a broad iron line with no corresponding Compton hump is also observed, and these authors find that the equivalent width of the Fe-K $\alpha$  feature is consistent with predictions made by Yaqoob et al. (2001) for an origin in the BLR or perhaps

a Compton-thin torus. Further, measurements of the velocity broadening of the optical H $\alpha$  line concur with those measured from the broad Fe-K $\alpha$  line ( $v = 2500 \text{ km s}^{-1}$ ), suggesting a similar origin for the emission. Using the same diagnostic, we estimate that we should expect a broad line  $EW \sim 75 - 100 \text{ eV}$  if it originates in an outflow from this region; this underestimates our observed equivalent width by nearly a factor of ten using a simple Gaussian line to represent the broad feature, and by roughly a factor of two if the better-fitting `laor` line is used. Also, our estimated lower-limit FWHM velocity of the line ( $v \sim 0.37c$ ) is more than an order of magnitude greater than the polarized optical H $\alpha$  FWHM measurement of Barth et al. (1999) ( $v \sim 5000 \text{ km s}^{-1}$ ). This is a strong indication that the origin of the broad Fe-K $\alpha$  line we observe comes from a location closer to the black hole than the BLR. Moreover, variability of the iron line on years-long timescales has been noted (Kadler et al. 2004c; Ros et al. 2007); this type of variation would not be expected from the typical symmetric BLR, as changes in flux due to passing clouds in this region should theoretically average out over time.

Finally, we consider a more exotic possibility: perhaps the broad iron line is produced by Cerenkov line-like radiation in the centralmost regions of the AGN. You et al. (2003) put forward the intriguing notion that in dense regions where the refractive index of the material is large, relativistic electrons impinging upon this material can produce Cerenkov radiation in a narrow wavelength range very close to the intrinsic atomic wavelength of the material. The combination of absorption and emission causes the final emission feature to be redshifted and skewed, making it appear very like a relativistic line produced by fluorescence in the inner accretion disk. Furthermore, because electrons are producing the emission rather than photons, no reflected continuum is expected. The cool, dense, iron-rich gas in the disk of NGC 1052 could provide the medium for this process, while the relativistic electrons in the corona or base of the jet would make for an ideal bombardment population. Further, this physical picture makes for a much more consistent match with our estimate of the distance of the broad line region of origin from the hard X-ray source: our lower-limit FWHM velocity for the broad Gaussian fit to the line yields  $v \sim 0.37c$ , which corresponds to an upper-limit distance of  $d \sim 8 r_g$ . This is well inside the BLR, and would instead be consistent with a lamp-post model for the corona or the base of the jet in NGC 1052.

Given our constraint on the distance of the broad iron line emitter from the hard X-ray source, iron fluorescence from the base of the jet itself is also a potential candidate for the emission mechanism. However, considering the orientation of the jets observed in NGC 1052 to our line of sight, it is unclear whether any fluorescent emission from the jet base could escape the system without being absorbed. Correlated radio flux measurements from the base of the jet and X-ray spectra of this source must be examined over several epochs, simultaneously whenever possible, in order to assess the likelihood of this scenario as well as that of the Cerenkov line-like radiation proposed by You et al. (2003). Only through a coordinated study of the jet and the accretion flow in NGC 1052 can we hope to understand the connection between these two vital physical processes in AGN.

## 6. Conclusions

Our 2007 *Suzaku* spectrum of the LINER galaxy NGC 1052 is consistent with X-ray spectra of this source from previous epochs, with a fairly flat power-law continuum that is heavily absorbed. Both intrinsic neutral and ionized columns are detected, along with evidence of Galactic photoabsorption. An extended thermal component is also detected, likely due to the interaction of the jets with the surrounding ISM. Both narrow and broad iron lines are observed, though interestingly there is little to no reflection seen in the spectrum above 10 keV: two different models used to characterize this feature both require  $R_{\text{refl}} < 0.01$ .

We are thus faced with a complex scenario for NGC 1052 in which the broad iron line is not consistent with the lack of observed continuum reflection from the disk. While it is possible that the reflection has been drowned out in some way by the power-law continuum, we must also consider the physical implications if reflection is indeed absent above 10 keV. It could be that we are witnessing the transition of the accretion flow to a radiatively-inefficient state at some critical radius in the disk, which would eliminate the Compton hump from the spectrum. A broad iron line is still produced, however, and the velocity width of the line is consistent with an origin close to the black hole. If the inner disk is an ADAF, it is possible that the broad Fe-K $\alpha$  line is emitted from material at the base of the jet, or perhaps from a yet more exotic mechanism such as the Cerenkov-like line radiation postulated by You et al. (2003). Further, coordinated observations of both the inner jet(s) (radio) and the inner accretion flow (X-ray) in order to solve this puzzle.

## Acknowledgments

LB and MK thank the NASA Postdoctoral Program, administered by ORAU, for their support at NASA's GSFC. KW gratefully acknowledges support from NASA grant NNX08AC226, MA from NASA grant NNX08AC22G. YYK is a Research Fellow of the Alexander von Humboldt Foundation. LB and KW also appreciate excellent advice from Tahir Yaqoob, Chris Reynolds, Julian Krolik, Demos Kazanas and Tim Kallman. KW would like to dedicate this work to Professor Andrew Wilson, who prompted her interest in LINERs and NGC 1052 in the 1990s.

## REFERENCES

- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17–+
- Barth, A. J., Filippenko, A. V., & Moran, E. C. 1999, *Astrophys. J. Lett.*, 515, L61

- Bianchi, S., La Franca, F., Matt, G., Guainazzi, M., Jimenez Bailón, E., Longinotti, A. L., Nicastro, F., & Pentericci, L. 2008, *Mon. Not. R. astr. Soc.*, L74+
- Braatz, J. A., Wilson, A. S., & Henkel, C. 1994, *Astrophys. J. Lett.*, 437, L99
- Claussen, M. J., Diamond, P. J., Braatz, J. A., Wilson, A. S., & Henkel, C. 1998, *Astrophys. J. Lett.*, 500, L129+
- Fabian, A. C., Rees, M. J., Stella, L., & White, N. E. 1989, *Mon. Not. R. astr. Soc.*, 238, 729
- Fender, R., Belloni, T., & Gallo, E. 2005, *Astrophys. & Space Sci.*, 300, 1
- Gabel, J. R., Bruhweiler, F. C., Crenshaw, D. M., Kraemer, S. B., & Miskey, C. L. 2000, *Astrophys. J.*, 532, 883
- Guainazzi, M., Oosterbroek, T., Antonelli, L. A., & Matt, G. 2000, *Astr. Astrophys.*, 364, L80
- Heckman, T. M. 1980, *Astr. Astrophys.*, 87, 152
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *Astrophys. J.*, 487, 568
- Kadler, M., Kerp, J., Ros, E., Falcke, H., Pogge, R. W., & Zensus, J. A. 2004a, *Astr. Astrophys.*, 420, 467
- Kadler, M., Ros, E., Lobanov, A. P., Falcke, H., & Zensus, J. A. 2004b, *Astr. Astrophys.*, 426, 481
- Kadler, M., Ros, E., Weaver, K., Kerp, J., & Zensus, J. A. 2004c, in *Bulletin of the American Astronomical Society*, Vol. 36, *Bulletin of the American Astronomical Society*, 823–+
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, *Astr. Astrophys.*, 440, 775
- Kataoka, J., Reeves, J. N., Iwasawa, K., Markowitz, A. G., Mushotzky, R. F., Arimoto, M., Takahashi, T., Tsubuku, Y., Ushio, M., Watanabe, S., Gallo, L. C., Madejski, G. M., Terashima, Y., Isobe, N., Tashiro, M. S., & Kohmura, T. 2007, *Publ. astr. Soc. Japan*, 59, 279
- Kinkhabwala, A. A. 2003, PhD thesis, AA(COLUMBIA UNIVERSITY)
- Knapp, G. R., Faber, S. M., & Gallagher, J. S. 1978, *Astr. J.*, 83, 139
- Laor, A. 1991, *Astrophys. J.*, 376, 90
- Magdziarz, P. & Zdziarski, A. A. 1995, *Mon. Not. R. astr. Soc.*, 273, 837
- Marscher, A. P. 2006, *Astron. Nachr.*, 327, 217

- Marscher, A. P., Jorstad, S. G., Gómez, J.-L., Aller, M. F., Teräsranta, H., Lister, M. L., & Stirling, A. M. 2002, *Nature*, 417, 625
- Mewe, R., Gronenschild, E. H. B. M., & van den Oord, G. H. J. 1985, *Astr. Astrophys. Suppl.*, 62, 197
- Miniutti, G., Fabian, A. C., Anabuki, N., Crummy, J., Fukazawa, Y., Gallo, L., Haba, Y., Hayashida, K., Holt, S., Kunieda, H., Larsson, J., Markowitz, A., Matsumoto, C., Ohno, M., Reeves, J. N., Takahashi, T., Tanaka, Y., Terashima, Y., Torii, K., Ueda, Y., Ushio, M., Watanabe, S., Yamauchi, M., & Yaqoob, T. 2007, *Publ. astr. Soc. Japan*, 59, 315
- Narayan, R. & Yi, I. 1994, *Astrophys. J. Lett.*, 428, L13
- Rees, M. J., Begelman, M. C., Blandford, R. D., & Phinney, E. S. 1982, *Nature*, 295, 17
- Reynolds, C. S. & Nowak, M. A. 2003, *Phys. Reports*, 377, 389
- Ros, E., Kadler, M., Kaufmann, S., Kovalev, Y. Y., Tueller, J., & Weaver, K. A. 2007, in *Highlights of Spanish Astrophysics IV*, ed. F. Figueras, J. M. Girart, M. Hernanz, & C. Jordi, 165–+
- Ross, R. R. & Fabian, A. C. 2005, *Mon. Not. R. astr. Soc.*, 358, 211
- Shakura, N. I. & Syunyaev, R. A. 1973, *Astr. Astrophys.*, 24, 337
- Weaver, K. A., Wilson, A. S., Henkel, C., & Braatz, J. A. 1999, *Astrophys. J.*, 520, 130
- Yaqoob, T., George, I. M., Nandra, K., Turner, T. J., Serlemitsos, P. J., & Mushotzky, R. F. 2001, *Astrophys. J.*, 546, 759
- Yaqoob, T. & Padmanabhan, U. 2004, *Astrophys. J.*, 604, 63
- You, J. H., Liu, D. B., Chen, W. P., Chen, L., & Zhang, S. N. 2003, *Astrophys. J.*, 599, 164
- Zdziarski, A. A., Johnson, W. N., Done, C., Smith, D., & McNaron-Brown, K. 1995, *Astrophys. J. Lett.*, 438, L63