



# **RENAL STONE RISK DURING SPACEFLIGHT: ASSESSMENT AND COUNTERMEASURE VALIDATION**

**PRINCIPAL INVESTIGATOR: Peggy A. Whitson, Ph.D., CB/NASA/JSC**

## **CO-INVESTIGATORS:**

**Robert A. Pietrzyk, M.S., SK/Wyle**

**Clarence F. Sams, Ph.D., SK/NASA/JSC**

**Jeffery A. Jones, M.D., SD/NASA/JSC**

**Scott M. Smith, Ph.D., SK/NASA/JSC**

## **SCIENCE TEAM:**


**Ed K. Hudson, Ph.D., SK/JES Tech**

**Mayra Nelman-Gonzalez, SK/Wyle**





# Risk



- **Mission risk and Impacts:**
  - ❖ Potential risk condition exists during the pre-, in- and postflight phases
  - ❖ Risk to crewmember for both acute and chronic health effects
  - ❖ Potential for significant impact to mission operational objectives
    - Early termination of mission
    - Significant impact to affected crewmember's performance
    - Significant impact to other crewmembers for medical care and treatment of affected crewmember




## EVIDENCE

- **As of 2008, 15 symptomatic urinary calculi have been experienced by 13 U.S. astronauts (Pietryzk, et al, 2006; Jones et al, 2008)**
- **Multiple stone events among cosmonauts reported by Russian medical investigators**
- **One in-flight episode nearly causing a mission termination but was resolved by spontaneous stone passage**





# NEPHROLITHIASIS – A MULTIFACTORIAL DISEASE






# SYMPTOMS/SIGNS



- **Severe / agonizing pain in the flank (back just below the ribs spreading around to the front of the abdomen) often extending into the groin area.**
- **Usually nausea and often vomiting**
- **Fever chills and sepsis, if infection is present**
- **Gross or microscopic blood in the urine**
- **Progression if not treated, hydronephrosis, renal shutdown**





# Urolithiasis and Stone Passage

Stone Size	Chance of Spontaneous Passage	Time to Pass Stone	Require surgical intervention
<2 mm	>85%	4.5-8 days	5%
<5 mm	78-80	7 – 14.5 days	17%
5-7 mm	20-50% (35% avg)	5.5-22 days	50%
>7 mm	< 10% (8% avg)	53 days - never	>80%

**Stones 3 mm in size can cause transient or complete obstruction**  
**Recurrence approx 5-10%/year up to 75% at 20 years**



# **STUDY OBJECTIVES**

- **Quantitate the pre-, in- and postflight risk of renal stone formation associated with space flight.**
- **Determine the efficacy of potassium citrate as a countermeasure in reducing the in-flight and postflight for renal stone formation.**
- **Evaluate dietary impact on the urinary biochemistry.**



# SUBJECTS



## Placebo Group: n = 18

NASA-Mir missions  
ISS missions

12 male subjects, mission duration 129 - 208 days  
6 male subjects, mission duration 93 - 175 days

## KCIT Group: n = 12

ISS missions

11 male/1 female subjects, mission duration 93 - 175 days







## METHODS



- **24-hour urines collected pre-, in-, and post-flight**
- **Food, fluid, exercise, and medications monitored before and during the urine collection period**
- **Two potassium citrate (KCIT) pills, 10 mEq/pill, ingested daily (with the last meal of the day) from L-3 days to R+14 days**
  - **Double-blind study design except for last 3 ISS subjects**
- **Biochemical analysis of urine samples for urinary factors associated with stone formation**
- **Dietary analysis completed to assess environmental influences on the urinary biochemistry**





# INVESTIGATION RESULTS



# Potassium Citrate

The majority of oral citrate is metabolized in the liver to bicarbonate, each citrate ion producing three bicarbonate ions.



↑  $\text{HCO}_3$

↑ pH

↑ U-Citrate

↓ U-Calcium

↓ Urinary Supersaturation

↓ Renal Stone

**KCIT dosage of 20 mEq/d selected based on;**

- results from Shuttle and NASA-Mir missions
- minimize any potential for in-flight GI upset (wax matrix/ slow release prep)
- minimize the potential to exaggerate the risk for CaP stones (higher pH 7.25-7.5)
- minimize impact to crew time

**Effects on renal physiology**

65-90% of filtered citrate is reabsorbed

10-35% of citrate is excreted into the urine

**Effects of dosage used (20 mEq/d)**

expected urinary increase of 130-140 mg/d



expected rise in urinary pH of 0.2 – 0.3 units



# FLUID BALANCE

Similar fluid intake and total urine  
volumes between groups


Low urine volumes (< 2L/d)



↓ Fluid intake during flight




## Effect of Potassium Citrate on Urinary pH




↑ Urinary pH in  
KCIT crewmembers,  
but not too high



## Effect of Potassium Citrate on Uric Acid Supersaturation





↓ Risk of uric acid stone formation in KCIT crewmembers



Uric Acid Stones  
Image from Mission Pharmacal



# CALCIUM BALANCE



Dietary Ca intake below  
recommended levels

↓ Urinary Ca excretion in KCIT  
crewmembers

**PLACEBO SUBJECTS**

IN-FLIGHT CALCIUM OXALATE SUPERSATURATION (log)

PREFLIGHT CALCIUM OXALATE SUPERSATURATION (log)

## KCIT subjects maintained calcium oxalate risk at preflight levels

## A close-up photograph of a small, light brown, irregularly shaped object, possibly a seed or a piece of food, resting on a textured surface. The object has a bumpy, granular texture and is positioned in the center of the frame. The background is dark and out of focus, while the surface it sits on is lighter and has a fine, grainy texture.

**KClF SUBJECTS**

IN-FLIGHT CALCIUM OXALATE SUPERSATURATION (log)

PREFLIGHT CALCIUM OXALATE SUPERSATURATION (log)


Pre-flight Supersaturation (log)	In-flight Supersaturation (log)
0.05	0.75
0.05	0.70
0.05	1.35
0.35	0.65
0.35	0.70
0.35	1.15
0.35	1.15
0.35	-0.40
0.55	0.75
0.55	1.25
0.55	0.35
0.60	0.55
0.60	0.50
0.60	0.65
0.65	0.20
0.65	0.40
0.65	0.65
0.70	0.15
0.70	0.65
0.75	0.10
0.75	0.15
0.75	0.75
0.80	0.65
0.80	1.00
0.80	1.05
1.00	0.60
1.00	0.70
1.00	0.75
1.00	0.80
1.00	1.05






## CASE STUDY

**CREWMEMBER PARTICIPATED  
IN BOTH MIR AND ISS MISSIONS**  
**MIR – No Treatment**  
**ISS - KCIT**




**↓ IN-FLIGHT CALCIUM EXCRETION  
DURING KCIT INGESTION**

**MIR IN-FLIGHT  
HYPERCALCIURIA**






## CASE STUDY




Low urine volume during both missions






## CASE STUDY



Supersaturation values >2.0 indicate greater risk for stone formation





## **SIGNIFICANT FINDINGS**

- **KCIT treated subjects exhibited decreased urinary calcium excretion.**
- **KCIT subjects maintained the levels of calcium oxalate supersaturation risk at their preflight levels.**
- **Increased urinary pH levels in KCIT treated subjects reduced the risk of uric acid stones.**
- **Individual crewmember response may play a role in renal stone susceptibility and efficacy of countermeasures.**



A cosmic background featuring a large, detailed view of Earth's horizon on the right side, showing swirling cloud patterns. In the upper center, the Moon is visible as a dark sphere with a thin white ring of light. The rest of the image is a deep blue space filled with numerous stars and a faint, glowing nebula or galaxy structure.

# **Risk Mitigation Strategies and Recommended Actions**



# Recommendations

- Encourage increased fluid intake to increase urine volume





# Recommendations

## ➤ Use of Potassium Citrate

- urinary inhibitor of calcium-containing stones, binds with calcium reducing the amount of calcium available to form CAOx
- inhibits crystal growth, aggregation and nucleation
- alkalinizes urine and decrease urinary calcium excretion
- supported by Space Medicine
- in Transition to Medical Practice process for operational use

## ➤ Assess dietary influences

- decrease protein, sodium and oxalate intake
- maintain calcium intake to recommended levels

## ➤ Perform urinary risk assessments

- identify crewmembers who are at any elevated risk
- provides an education program to help humans remain healthy during space exploration





# FUTURE POTENTIAL







## Potential In-Flight Prediction of Stone Risk



Urine Monitoring System (UMS)

Capability to measure urine volume provided with the installation of the UMS on Flight 20A and the addition of the in-line calcium sensor for real-time data collection

Development of an oxalate sensor would be required to optimize real-time risk



## **PATIENT COMMENTS: Characterizing the symptoms**



**“ I'd rather give birth to an elephant than go through this”.**

**“ Like being hit with a two-by-four” .**

**“ Like being shot with an arrow”.**

**“ Pain came on suddenly and did not pass until doctors hopped me up on pain meds. I've had my gall bladder removed and nearly severed my thumb but I never in my life felt pain like this. Would wish it upon no one!!**

**“ I have had my leg crushed by a car backing over it and that has nothing compared to the pain of a kidney stone” !!!!**



## **ACKNOWLEDGEMENTS**

- **All the astronauts and cosmonauts who participated in this study**
- **Mission Pharmacal, San Antonio, TX. for kindly providing both the potassium citrate and placebo drugs (Space Act Agreement)**
- **Mineral Metabolism Lab, Center for Mineral Metabolism & GCRC, UT Southwestern Med.Ctr. Dallas, TX**
- **NASA Johnson Space Center Clinical Laboratory**
- **NASA Johnson Space Center Nutritional Biochemistry Laboratory**
- **ISSMP Science and Flight Hardware support teams**