



## Solution Preserves Nucleic Acids in Body-Fluid Specimens

Specimens can be stored and transported at room temperature.

*Lyndon B. Johnson Space Center, Houston, Texas*

A solution has been formulated to preserve deoxyribonucleic acid (DNA) and ribonucleic acid (RNA) in specimens of blood, saliva, and other bodily fluids. Specimens of this type are collected for diagnostic molecular pathology, which is becoming the method of choice for diagnosis of many diseases. The solution makes it possible to store such specimens at room temperature, without risk of decomposition, for subsequent analysis in a laboratory that could be remote from the sampling location. Thus, the solution could be a means to bring the benefits of diagnostic molecular pathology to geographic regions where refrigeration equipment and diagnostic laboratories are not available.

The table lists the ingredients of the solution. The functions of the ingredients are the following:

- EDTA chelates divalent cations that are necessary cofactors for nuclease activity. In so doing, it functionally removes these cations and thereby retards the action of nucleases. EDTA also stabilizes the DNA helix.
- Tris serves as a buffering agent, which is needed because minor contaminants in an unbuffered solution can exert pronounced effects on pH and

Ingredient	Weight or Volume of Ingredient	Final Concentration of Ingredient in Solution
Sodium dodecyl sulfate (SDS)	1 g	1 percent
Ethylenediaminetetraacetic acid (EDTA)	0.037 g	1.0 mM
Tris(hydroxymethyl)aminomethane (also known as "Tris" and sold under the trade name Trizma™ Base)	0.12 g	10 mM
Water Free of Deoxyribonuclease and Ribonuclease	99 mL	
	<b>Total Volume</b> 100 mL	

The **Nucleic Acid Stability Solution (NASS)** contains ingredients that perform different roles essential to the preservation of DNA and RNA in specimens. In the preparation of this solution, the ingredients are mixed in the indicated quantities (or common multiples thereof), then the solution is sterilized by passing it through a 0.2- $\mu$ m filter.

thereby cause spontaneous degradation of DNA.

- SDS is an ionic detergent that inhibits ribonuclease activity. SDS has been used in some lysis buffers and as a storage buffer for RNA after purification.

The use of the solution is straightforward. For example, a sample of saliva is collected by placing a cotton roll around in the subject's mouth until it becomes saturated, then the cotton is placed in a collection tube. Next, 1.5 mL of the solution are injected directly

into the cotton and the tube is capped for storage at room temperature. The effectiveness of the solution has been demonstrated in tests on specimens of saliva containing herpes simplex virus. In the tests, the viral DNA, as amplified by polymerase chain reaction, was detected even after storage for 120 days.

*This work was done by Duane L. Pierson of Johnson Space Center and Raymond P. Stowe. For further information, contact the Johnson Commercial Technology Office at (281) 483-3809. MSC-22891*

## Oligodeoxynucleotide Probes for Detecting Intact Cells

Cells can be detected, identified, and enumerated via chemiluminescence.

*Lyndon B. Johnson Space Center, Houston, Texas*

A rapid, sensitive test using chemiluminescent oligodeoxynucleotide probes has been developed for detecting, identifying, and enumerating intact cells. The test is intended especially for use in detecting and enumerating bacteria and yeasts in potable water.

As in related tests that have been developed recently for similar purposes, the oligodeoxynucleotide probes used in this test are typically targeted at either single-copy deoxyribonucleic acid (DNA) genes (such as virulence genes) or the multiple

copies (10,000 to 50,000 copies per cell) of 16S ribosomal ribonucleic acids (rRNAs). Some of those tests involve radioisotope or fluorescent labeling of the probes for reporting hybridization of probes to target nucleic acids. Others of those tests involve labeling with enzymes plus the use of chemiluminescent or chromogenic substrates to report hybridization via color or the emission of light, respectively. The present test is of the last-mentioned type. The chemiluminescence in the present test can be detected easily with relatively sim-

ple instrumentation.

In developing the present test, the hybridization approach was chosen because hybridization techniques are very specific. Hybridization detects stable, inheritable genetic targets within microorganisms. These targets are not dependent on products of gene expression that can vary with growth conditions or physiological states of organisms in test samples. Therefore, unique probes can be designed to detect and identify specific genera or species of bacteria or