

# First INTEGRAL and Swift observations of a giant outburst of A 0535+26

I. Caballero\*, S. Müller<sup>†</sup>, P. Bordas\*\*, C. Ferrigno\*\*, M. Kühnel<sup>‡</sup>,  
K. Pottschmidt<sup>‡</sup>, P. Kretschmar<sup>§</sup>, J. Wilms<sup>†</sup>, I. Kreykenbohm<sup>†</sup>,  
D. Klochkov<sup>¶</sup>, A. Santangelo<sup>¶</sup>, R. Staubert<sup>¶</sup>, S. Suchy<sup>||</sup>, R. Rothschild<sup>||</sup>,  
A. Camero-Arranz<sup>††</sup> and M. Finger<sup>††</sup>

\*CEA Saclay, DSM/IRFU/SAP –UMR AIM (7158) CNRS/CEA/Univ. P. Diderot –F-91191, France

<sup>†</sup>Dr. Karl Remeis-Sternwarte and ECAP, Germany

\*\*ISDC Data Centre for Astrophysics, Switzerland

<sup>‡</sup>CRESST, UMBC and NASA GSFC, Code 661

<sup>§</sup>ESAC, ESA, Spain

<sup>¶</sup>IAAT, Tübingen, Germany

<sup>||</sup>CASS UCSD

<sup>††</sup>NSSTC

## Abstract.

The Be/X-ray binary A 0535+26 has shown three giant outbursts since 2005, after a long period of quiescence. The giant outbursts in 2005 ( $\sim 5.2$  Crab, 15-50 keV range) and 2009 ( $\sim 5.6$  Crab) could not be observed by most X-ray observatories due to Sun observing constraints. Finally, a giant outburst in February 2011, that reached a flux of  $\sim 3.8$  Crab, was monitored with *INTEGRAL* and *Swift* TOO observations. We present first results these observations, with a special focus on the cyclotron lines present in the X-ray spectrum of the source.


**Keywords:** X-rays: binaries - stars: magnetic fields - stars: individual: A 0535+26

**PACS:** 97.60.Gb, 97.60.Jd, 97.80.Jp

The Be/X-ray binary A 0535+26 was discovered by *Ariel V* during a giant outburst in 1975 [6]. At a distance of  $\sim 2$  kpc, a  $P_{\text{spin}} \sim 103$  s pulsar orbits the B0 IIIe star HDE 245770 [8] in an eccentric orbit ( $e = 0.47$ ) of  $P_{\text{orb}} \sim 111.1$  d [3]. A long quiescence period of more than 11 years ended with a giant outburst in 2005. Since then, several normal and giant outbursts have been observed, all of them around the periastron passage. The source presents cyclotron lines in its X-ray spectrum at  $\sim 46$  and  $\sim 100$  keV, discovered during a giant outburst in 1989 with *HEXE* [4]. From the cyclotron lines, a magnetic field of  $B \sim 4 \times 10^{12}$  G is inferred.

A 0535+26 underwent a giant outburst in February 2011, observed with *INTEGRAL* and *Swift*. The *Swift*-BAT light curve of the outburst and the times of the observations are shown in Fig. 1 (left). These observations provide for the first time the possibility to study the cyclotron line evolution of the source during a giant outburst.

We extracted phase averaged spectra for all the *Swift* and *INTEGRAL* observations, and fitted the broad band X-ray spectrum with a cutoff powerlaw plus a cyclotron line at  $\sim 46$  keV, modeled with a Gaussian optical depth profile as in [2]. Preliminary results of the cyclotron line energy as a function of the luminosity are given in Fig. 1 (right). The *INTEGRAL*-*ISGRI* ancillary response file (ARF) is based on Crab observations. In



**FIGURE 1.** *Left:* *Swift*-BAT light curve of the February 2011 giant outburst of A 0535+26. The times of the *INTEGRAL* and *Swift* observations are shown. *Right:* cyclotron line energy evolution with the luminosity, from *RXTE*, *INTEGRAL*, and *Suzaku* observations of a normal outburst in 2005 [9, 1] and from the *INTEGRAL* observations of the 2011 giant outburst (preliminary).

most of our observations, A 0535+26 is significantly brighter than the Crab, and the ARF needs to be adjusted for these observations. For our preliminary analysis, we restricted the *ISGRI* analysis to energies below 80 keV.

Variations of the cyclotron line with the X-ray luminosity can be used to probe the change of the accretion structure and plasma properties with the mass accretion rate. Contrary to other sources (see, e.g., [5, 10, 7]), for A 0535+26 no significant variation of the cyclotron line energy with the X-ray luminosity was observed [9, 1]. The new *INTEGRAL* observations allow us to extend the studied luminosity range up to  $L_{(3-50)\text{keV}} \sim 3.4 \times 10^{37} \text{erg s}^{-1}$ . As seen in Fig. 1 (right), one of the *INTEGRAL* measurements of the cyclotron line energy is significantly lower than the other ones. The other values remain rather stable in the luminosity range  $\sim 0.037 - 3.4 \times 10^{37} \text{erg s}^{-1}$ . A more in-depth analysis of the *ISGRI* data is ongoing to account for the current uncertainties in the response matrix in order to confirm these results.

## ACKNOWLEDGMENTS

IC acknowledges financial support from the French Space Agency CNES through CNRS

## REFERENCES

1. Caballero, I., Kretschmar, P., Santangelo, A., et al. 2007, *Astron. Astrophys.*, 465, L21
2. Coburn, W., Heindl, W. A., Rothschild, R. E., et al. 2002, *Astrophys. J.*, 580, 394
3. Finger, M. H., Camero-Arranz, A., Kretschmar, P., Wilson, C., & Patel, S. 2006, in *BAAS*, Vol. 38, 359
4. Kendziorra, E., Kretschmar, P., Pan, H. C., et al. 1994, *Astron. Astrophys.*, 291, L31
5. Mihara, T. 1995, PhD thesis, Dept. of Physics, Univ. of Tokyo
6. Rosenberg, F. D., Eyles, C. J., Skinner, G. K., & Willmore, A. P. 1975, *Nature*, 256, 628
7. Staubert, R., Shakura, N. I., Postnov, K., et al. 2007, *Astron. Astrophys.*, 465, L25
8. Steele, I. A., Negueruela, I., Coe, M. J., & Roche, P. 1998, *Mon. Not. R. Astron. Soc.*, 297, L5
9. Terada, Y., Mihara, T., Nakajima, M., et al. 2006, *Astrophys. J.*, 648, L139
10. Tsygankov, S. S., Lutovinov, A. A., & Serber, A. V. 2010, *Mon. Not. R. Astron. Soc.*, 401, 1628