

**CONTROL ID:** 1209009

**TITLE:** Visible and Near-IR Reflectance Spectra of Smectite Acquired Under Dry Conditions for Interpretation of Martian Surface Mineralogy

**PRESENTATION TYPE:** Assigned by Committee (Oral or Poster)

**CURRENT SECTION/FOCUS GROUP:** Planetary Sciences (P)

**CURRENT SESSION:** P22. Phyllosilicate Formation on Mars: Connecting Experimental and Theoretical Studies with Planetary Observations

**AUTHORS (FIRST NAME, LAST NAME):** Richard V Morris<sup>1</sup>, Cherie N Achilles<sup>2</sup>, Paul D Archer<sup>1</sup>, Trevor G Graff<sup>2</sup>, David G Agrestis, Douglas W Ming<sup>1</sup>, Dadi C Golden<sup>2</sup>, Stanley A Mertzman<sup>4</sup>

**INSTITUTIONS (ALL):** 1. NASA Johnson Space Center, Houston, TX, United States.

2. ESCG Jacobs Technology, Houston, TX, United States.

3. Univ. Alabama Birmingham, Birmingham, AL, United States.

4. Franklin and Marshall College, Lancaster, PA, United States.

**ABSTRACT BODY:** Visible and near-IR (VNIR) spectra from the MEx OMEGA and the MRO CRISM hyper-spectral imaging instruments have spectral features associated with the H<sub>2</sub>O molecule and M OH functional groups (M = Mg, Fe, Al, and Si). Mineralogical assignments of martian spectral features are made on the basis of laboratory VNIR spectra, which were often acquired under ambient (humid) conditions. Smectites like nontronite, saponite, and montmorillonite have interlayer H<sub>2</sub>O that is exchangeable with their environment, and we have acquired smectite reflectance spectra under dry environmental conditions for interpretation of martian surface mineralogy. We also obtained chemical, Moessbauer (MB), powder X-ray diffraction (XRD), and thermogravimetric (TG) data to understand variations in spectral properties. VNIR spectra were recorded in humid lab air at 25-35C, in a dynamic dry N<sub>2</sub> atmosphere (50-150 ppmv H<sub>2</sub>O) after exposing the smectite samples (5 nontronites, 3 montmorillonites, and 1 saponite) to that atmosphere for up to ~1000 hr each at 25-35C, ~105C, and ~215C, and after re-exposure to humid lab air. Heating at 105C and 215C for ~1000 hr is taken as a surrogate for geologic time scales at lower temperatures. Upon exposure to dry N<sub>2</sub>, the position and intensity of spectral features associated with M-OH were relatively insensitive to the dry environment, and the spectral features associated with H<sub>2</sub>O (e.g., ~1.90 um) decreased in intensity and are sometimes not detectable by the end of the 215C heating step. The position and intensity of H<sub>2</sub>O spectral features recovered upon re-exposure to lab air. XRD data show interlayer collapse for the nontronites and Na-montmorillonites, with the interlayer remaining collapsed for the latter after re-exposure to lab air. The interlayer did not collapse for the saponite and Ca-montmorillonite. TG data show that the concentration of H<sub>2</sub>O derived from structural OH was invariant to the dry N<sub>2</sub> treatment for saponite and the montmorillonites, but the nontronites had additional structural OH after treatment. Upon exposure to dry N<sub>2</sub>, the VNIR spectra also acquired a red slope with decreasing albedo between ~0.4 and ~2.0 um. The magnitude of the effects covaries with exposure time to dry N<sub>2</sub> and heating temperature. Upon re-exposure to lab air, the slope and albedo do not completely recover to pre-exposure values. MB data show that these effects do not result from partial reduction of ferric to ferrous iron, and TG data show they do not result from loss of structural OH. Possible explanations include formation of small clusters of (superparamagnetic) ferric oxide and reduced smectite crystallinity. The difference in spectral properties between spectra acquired in humid lab air and under dry conditions are consequential for interpretation of CRISM and OMEGA spectra. For example, nontronite by itself and not nontronite plus ferrihydrite can account for the red spectral slope in martian spectra where nontronite is indicated by the Fe-OH spectral features.

**KEYWORDS:** [5470] PLANETARY SCIENCES: SOLID SURFACE PLANETS / Surface materials and properties, [5464] PLANETARY SCIENCES: SOLID SURFACE PLANETS / Remote sensing, [5415] PLANETARY SCIENCES: SOLID SURFACE PLANETS / Erosion and weathering, [3672] MINERALOGY AND PETROLOGY / Planetary mineralogy and petrology.