


Onward!
Kepler's Exoplanet Census

Kepler





Natalie Batalha
NASA Ames Research Center

● Jun 2010

● Feb 2011



● Feb 2012





Extended Mission

Earth-size planets in the HZ




Batalha et al. 2012, arXiv: 1202.5852




HZ Candidates

48 with T_{eq} between 185 and 303 K






HZ Candidates



Potential Habitable Exoplanets

(4 confirmed and 27 unconfirmed NASA Kepler Candidates)



Credit: Planetary Habitability Laboratory, UPR Arecibo (phl.upr.edu) April 2012

er-size

une-size

n-size

Equilibrium Temperature [K]

Batalha et al. 2012, arXiv: 1202.5852





HZ Candidates

48 with T_{eq} between 185 and 303 K

● Jun 2010

● Feb 2011

● Jan 2012




Equilibrium Temperature [K]

Batalha et al. 2012, arXiv: 1202.5852


The Multiples




● one - 1425 ● two - 490 ● three - 252 ● four - 108 ● five - 40 ▲ six - 6




Batalha et al. 2012, arXiv: 1202.5852




KOI-1805
KOI-864
KOI-1589
KOI-1194
KOI-906
KOI-749
KOI-723
KOI-571
KOI-921
KOI-94
KOI-70
KOI-1358
KOI-1422
KOI-711
KOI-2148
KOI-250
KOI-137
KOI-115
KOI-1078
KOI-812
KOI-884
KOI-899
KOI-869
KOI-1563
KOI-117
KOI-658
KOI-733
KOI-1860
KOI-1161
KOI-510
KOI-1432
KOI-952
KOI-1127
KOI-720
KOI-1364
KOI-248
KOI-756
KOI-1867
KOI-1436
KOI-880
KOI-941
KOI-1835
KOI-904
KOI-2169
KOI-85
KOI-834
KOI-1916
KOI-343
KOI-2220
KOI-1306
KOI-398
KOI-939
KOI-665
KOI-377
KOI-481
KOI-1557
KOI-938
KOI-1198
KOI-500
KOI-2248
KOI-191
KOI-961









Fabrycky et al. arXiv:2012.6328



- KOI-623
- KOI-1909
- KOI-520
- KOI-82
- KOI-1952
- KOI-156
- KOI-898
- KOI-1931
- KOI-907
- KOI-148
- KOI-1060
- KOI-1567
- KOI-718
- KOI-1832
- KOI-490
- KOI-1805
- KOI-864
- KOI-1589
- KOI-1194
- KOI-906








Kepler-20e





Venus



Earth



Kepler-20f




Fressin et al. 2012

Kepler-22b



$P=289.8623$ days

$R_p=2.38\pm0.13 R_\oplus$


$T_{eq}=262$ K




Borucki et al. 2012, ApJ, 745, 120





Total SNR > 140 (22 months of data)




Batalha et al. 2011, ApJ, 729, 27









Kepler-16

(2) Kepler-10


Kepler-19

Kepler-19 (2)

Kepler-19 (3)

Kepler-19 (4)

Kepler-19 (2)



Kepler-16


(2) Kepler-10

Other Systems: A TTV-Confirmation Catalog

Kepler

KO

Kepler






9:7 resonance

near 2:1

near 2:1



near 3:2




Ford et al. 2012, Steffen et al. 2012, Fabrycky et al. 2012

Kepler-19 (2)





Doyle et al 2011, Science, 333, 1602



Kepler-16

Kepler-35

Kepler-34

22

Doyle et al 2011, Science, 333, 1602

Kepler-16


(2) Kepler-10




Doyle et al 2011, Science, 333, 1602


Kepler-16


(2) Kepler-10



Doyle et al 2011, Science, 333, 1602









Kepler-16 ●
(2) Kepler-10 ●
Kepler-4 ●
TrES-2 ●
Kepler-23 (2) ●
Kepler-15 ●


Kepler-24 (2) ●
Kepler-19 (2) ●
Kepler-9 (3) ●
Kepler-10 (3) ●





KOI-872 b & c




Nesvorny et al., ScienceExpress, 12 May 10





Kepler-16 ●
 (2) Kepler-10 ●●
 Kepler-4 ●
 Kepler-23 (2) ●
 Kepler-15 ●
 TrES-2 ●



(2)


Kepler-37

Kepler-8 ●

Kepler-30 (3)


(2)

Kepler-18 (3)




(2) Kepler-10


Kepler




SOHO




From Thierry Appourchaux...






The Stellar Sample




Pinsonneault et al 2012: recalibration of KIC photometry; systematic T_{eff} error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics

The Stellar Sample




Pinsonneault et al 2012: recalibration of KIC photometry; systematic T_{eff} error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics

The Stellar Sample



Pinsonneault et al 2012: recalibration of KIC photometry; systematic T_{eff} error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics


The Stellar Sample



Muirhead et al 2012, ApJ, 747, 144

Pinsonneault et al 2012: recalibration of KIC photometry; systematic T_{eff} error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics


The Stellar Sample



Muirhead et al. 2012 arXiv:1109.1819v2

Pinsonneault et al 2012: recalibration of KIC photometry; systematic T_{eff} error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics

The Stellar Sample



Pinsonneault et al 2012: recalibration of KIC photometry; systematic Teff error
Muirhead et al 2011: IR spectra of 84 cool stars; factor of 2 smaller stellar radii
Mann et al. 2012: systematic errors in parent sample affect exoplanet statistics

Exoplanet Science in the Extended Mission




- KOI Review Team (michael.r.haas@nasa.gov)
- Stellar Characterization (your name here)
- Small Planet Completeness (jessie.l.christiansen@nasa.gov)
- Asteroseismology (gillil@stsci.edu)
- Transit Timing Variations (eford@astro.ufl.edu)
- Eclipsing Binaries (aprsa@villanova.edu)
- False Positive Tracking (Steve.Bryson@nasa.gov)
- Follow-up Observation Program (ciardi@ipac.caltech.edu)
- Other (Giant Planets, Circumbinary Planets)



Summary

- 72 Confirmed and/or characterized planets
 - rocky, earth-size, HZ, circumbinary, multiples
- 2,321 planet candidates, 1091 of which are new this year
- 16 months of data has been search for transits
- 250 earth-size candidates, one of which (KOI-2124.01) is in/near the HZ
- Power-law increase toward small planets down to $2 R_e$
- Pipeline completeness still improving
- 20% of stars observed to harbor >1 candidate
- Paucity of short-period giants in flat systems
- Small pile-up just outside of low-order resonances
- Stellar classification is high priority for reliable planet radii
- Stellar Variability (6.5-hour) is 50% higher than predicted
- Extending mission from 3.5 to 7.5 years recovers baseline performance.
- Extending mission allows for determinations of eta-Earth-size **in the HZ.**



Onward!

natalie.batalha@nasa.gov




Supplemental Slides



Distribution of Planet Diameters

For Orbital Periods < 50 Days



Howard et al. 2011, arXiv: 1103.2541




$N(Q6)/N(Q5)$

	1.25 Re < Rp < 2 Re		2.5 Re < Rp < 6 Rp
	5 < P < 50	50 < P < 150	10 < P < 125
Modeled	1.1	1.4	1.0
Observed	2.8	7.2	1.6

Gains cannot be explained solely by longer data collection.



Fractional Gains




$R_p < 2 R_e$: 197%
 $R_p > 2 R_e$: 52%

$P > 50$ days : 123%
 $P < 50$ days: 85%

Factors Affecting Completeness

Sudden Pixel Sensitivity Dropouts:

Transit detection analysis keys off on anomalous event instead of real transits. This, also, is treated in PDC-MAP.



Original flux

PDC

PDC-MAP

arXiv:1203.1383

Impact of Measured Noise



Mission design based on a peak at 20 ppm


Histogram peaks at 30 ppm


- Yields only 2% completeness for Earth analogs in baseline mission

Sensitivity improves as $\sqrt{t}$

- Noise is uncorrelated in successive transits
- Yields ~50% completeness for Earth analogs with 7.5 year mission

Noise Histogram





Kepler
Field of
View

Vega

LYRA

M57

M56

Albireo

M71

SAGITTA

AQUILA

Altair

CYGNUS

Deneb

M29

North
American
Nebula

M39


DELPHINUS



Kepler Input Catalog

- KeplerCam, 1.2-m, Mt. Hopkins
 - SDSS g, r, i, z + D51
- Archive data
 - 2MASS JHK, USNO-B
- Yields Stellar Parameters:
 - T_{eff} , $\log(g)$, $\log(Z)$, A_V , $E(B-V)$
 - Kp (kepler magnitude)
 - Radius



<http://archive.mast.edu/kepler>
Brown et al. 2012, AJ, 142, 112





- KeplerCam
 - SDSS g,
- Archive data
 - 2MASS J
- Yields Stellar
 - T_{eff} , $\log(g)$
 - Kp (kepler)
 - Radius


<http://archive.brown.edu>



Factors Affecting Completeness

Multi-Quarter Transit Search:


- Feb 2011 catalog: analysis of Q0-Q5 data based on non-pipeline tools to detect long-period events.
- Didn't benefit from pipeline whitening filters



Factors Affecting Completeness

Pre-search Data Conditioning:

First generation filters occasionally introduced high frequency noise. This has been greatly improved with PDC-MAP.




Original flux

PDC


PDC-MAP

arXiv:1203.1383


Percent Completeness



Completeness Including Transit Probability




Histogram of Total SNR





Turn-over at 9.7-sigma implies pipeline incompleteness. This should be greatly improved with PDC-MAP reprocessing.


Improved Vetting Statistics



9.1 arcsec source offset



0.2 arcsec source offset





HARPS-N: 3.-6m TNG, Canary Islands;
commissioned in the spring