

# Venus: The first Habitable world of our Solar System?


Michael Way (Uppsala, NASA/GISS)

Tony Del Genio, Nancy Kiang, Linda Sohl,  
Tom Clune, Igor Aleinov, Maxwell Kelley

[NeXSS-ROCKE3D]

2015/10/23

# Nomenclature!



4.6-3.8 Gya Hadean


3.8-2.5 Gya Archean

2.5-0.5 Gya Proterozoic

# Faint Young Sun Paradox & Habitability Through Time

- Hard to get Earth warm enough to have liquid water during early Archean = 3.8-2.5 [2.9] Gya
- Hard to get Mars warm enough for global scale oceans to persist in same period
- For Venus' early history we may be in luck
  - Low solar flux & slower rotation rate => liquid H<sub>2</sub>O

# M. Claire 2012 (Fluxes at 1 AU)



# Paleo-Venus: Water?

- Water abundance?
  - High D/H  $\sim 150 \times$  terrestrial
    - Via history of escape & re-supply in last  $\sim 10^9$  years, not a history of primordial escape (Grinspoon 1993,97,98)?
  - Likely received approx. the same amount of H<sub>2</sub>O as Earth in early solar system history
  - Timescale of H<sub>2</sub>O loss  $<\sim 100$ s My (Kasting 1984,88) to 2By (Grinspoon & Bullock 2003)
  - Current estimates are **4-530m** of liquid water
- Early Dry or Wet alternatives impossible with present knowledge - arguments for & against both (Lewis & Prinn, 1984; Yung & DeMore 1999).

# Venus slow Rotation a recent phenomenon?

## 1. Result of thick atmosphere (thermal tides):

- A. R. Dobrovolskis, A. P. Ingersoll, Atmospheric tides and the rotation of Venus. I - Tidal theory and the balance of torques. *Icarus* 41, 1–17 (1980).

## 2. Perhaps thick atmosphere is not necessary:

- J. Leconte et al. 2015



## Asynchronous rotation of Earth-mass planets in the habitable zone of lower-mass stars

Jérémie Leconte,<sup>1,2,3\*</sup> Hanbo Wu,<sup>1,4</sup> Kristen Menou,<sup>3,5</sup> Norman Murray<sup>1,4</sup>


<sup>1</sup>Canadian Institute for Theoretical Astrophysics, 60st St George Street, University of Toronto, Toronto, ON, M5S3H8,

# Venus Geology

- Cosmochemical/geochemical **models** imply Venus' bulk composition ≈Earth (Lodders & Fegley 1998)
- Complete resurfacing of Venus in past 750My-1By
- Minimal magnetic field (today)
- No present day evidence of plate tectonics
  - But: present day evidence of volcanism (Mars Express)
- Need in-situ measurements of Xe, Kr, Ar & isotopes to provide accessible record of ancient events pertinent to planetary formation & early evolutionary processes


# 1-D Simulations: Grinspoon & Bullock 2003

- 1 bar N<sub>2</sub>, CO<sub>2</sub>=.35mb, H<sub>2</sub>O=6mb
- 100% cloud cover
- Yields mean global surface temp ~300K




# GISS: 3-D General Circulation Model

- Modern Venus topography, radius, mass, spin
- N<sub>2</sub>=1bar, CO<sub>2</sub>=0.4mb, CH<sub>4</sub>=0.001mb
- 2.9Gya solar spectrum from M. Claire 2012
  - ~0.820 x present day solar luminosity




# 3-D Model Setup: Venus topography changes

fgrnd in Z72X46N\_venus001




fgrnd in Z72X46N\_venus002




# Venus land/ocean


Venus Ocean volume equivalent to 310m depth average




# Paleo-Venus Surface Temperature




# Venus preliminary results?




# High Level Cloudiness => High Albedo



# Hadley Cell is different from Earth's



# Other Worlds



# Conclusions

- Venus not Earth or Mars may have been the first habitable world of our solar system
- Can fewer assumptions be made for 1 vs another?
- Need for new in-situ observations to confirm Venus' geologic history – and water inventory