

When A Standard Candle Flickers: Hard X-ray Variations in the Crab Nebula



Colleen A. Wilson-Hodge
(NASA/MSFC)

Image Credits: X-ray: NASA/CXC/SAO/F.Seward; Optical: NASA/ESA/ASU/J.Hester & A.Loll; Infrared: NASA/JPL-Caltech/Univ. Minn./R.Gehrz


Coauthors/Collaborators

M.L. Cherry (LSU), G.L. Case (LaSierra), W.H. Baumgartner (NASA/CRESST), E. Beklen (NRAO), P.N Bhat, M.S. Briggs (UAH), R. Buehler (DESY), A. Camero-Arranz (IEECC-CSIC), V. Connaughton (UAH), R. Diehl (MPE), M.H. Finger (USRA), N. Gehrels (NASA/GSFC), J. Greiner (MPE), F. Harrison (CalTech), E. Hays (GSFC), K. Jahoda (NASA/GSFC), P. Jenke (UAH), R.M. Kippen (LANL), C. Kouveliotou (NASA/MSFC), H. A. Krimm (CRESST/NASA/GSFC/USRA), E. Kuulkers (ESA/ESAC), K.Madsen (CalTech), C. Markwardt (NASA/GSFC), C.A. Meegan (USRA), L. Natalucci (INAF-IASF), W.S. Paciesas (USRA), R. Preece (UAH), J.C. Rodi (LSU), N. Shaposhnikov, G.K. Skinner (Univ. of Birmingham), D. Swartz (USRA), A. von Kienlin, R.Diehl, X. Zhang (MPE)

MAXI data from <http://maxi.riken.jp>


Suzaku data from Kouzu et al. 2013, PASJ, 65, 74

Introduction



- 1054 AD Supernova at 2 kpc
- Consists of a pulsar, pulsar wind nebula, and a cloud of expanding ejecta
- Energy spectrum: synchrotron & inverse Compton components


Current Hard X-ray Light Curve



GBM 100-500 keV light curves


2008

2014



Predicted Time Variability


- Using MHD simulations, Volpi et al (2009) found characteristic timescales of 1-2 years at energies <0.75 MeV.
- The magnitude of the variations is larger than observed at lower energies



Volpi et al. 2009, A&A 485, 337

High Energy (MeV-GeV) Flaring

100 MeV - 300 GeV Flux (Nebula + Pulsar)



Summary & Conclusions

- The Crab Nebula shows both long-term and short timescale variability.
 - In Hard X-rays the Crab flux has returned to near 2008 levels, showing variations of up to $\sim 3.5\%/year$
 - At higher energies, 7 flares have been reported between 2007 and 2014, seen with AGILE and Fermi LAT
- The location and mechanism producing both types of variability is still mysterious.
- Crab monitoring demonstrates the value of overlapping instruments, all-sky monitoring, and need for keV-MeV coverage