




Print



Close Window



Your abstract appears below.

**Please print a copy of this page for your records.**

To return to the Submission Center and check your list of submissions; click "View Submissions" in the left menu.

#### Proof

**CONTROL ID:** 1811782

**TITLE:** Constraining Cometary Crystal Shapes from IR Spectral Features

**AUTHORS (FIRST NAME, LAST NAME):** Diane H Wooden<sup>1</sup>, Sean Lindsay<sup>2</sup>, David E Harker<sup>3</sup>, Michael S.P. Kelley<sup>4</sup>, Charles E. Woodward<sup>5</sup>, James Richard Murphy<sup>6</sup>

**INSTITUTIONS (ALL):** 1. NASA Ames Research Center, Moffett Field, CA, United States.

2. U. Tennessee, Knoxville, TN, United States.

3. UCSD/CASS, San Diego, CA, United States.

4. U. Maryland, College Park, MD, United States.

5. MN Institute for Astrophysics, Minneapolis, MN, United States.

6. NMSU, Las Cruces, NM, United States.

**ABSTRACT BODY:** A major challenge in deriving the silicate mineralogy of comets is ascertaining how the anisotropic nature of forsterite crystals affects the spectral features' wavelength, relative intensity, and asymmetry. Forsterite features are identified in cometary comae near 10, 11.05-11.2, 16, 19, 23.5, 27.5 and 33  $\mu\text{m}$  [1-10], so accurate models for forsterite's absorption efficiency ( $Q_{\text{abs}}$ ) are a primary requirement to compute IR spectral energy distributions (SEDs,  $\lambda F_{\lambda}$  vs.  $\lambda$ ) and constrain the silicate mineralogy of comets. Forsterite is an anisotropic crystal, with three crystallographic axes with distinct indices of refraction for the a-, b-, and c-axis. The shape of a forsterite crystal significantly affects its spectral features [13-16]. We need models that account for crystal shape.

The IR absorption efficiencies of forsterite are computed using the discrete dipole approximation (DDA) code DDSCAT [11,12]. Starting from a fiducial crystal shape of a cube, we systematically elongate/reduce one of the crystallographic axes. Also, we elongate/reduce one axis while the lengths of the other two axes are slightly asymmetric (0.8:1.2). The most significant grain shape characteristic that affects the crystalline spectral features is the relative lengths of the crystallographic axes. The second significant grain shape characteristic is breaking the symmetry of all three axes [17].

Synthetic spectral energy distributions using seven crystal shape classes [17] are fit to the observed SED of comet C/1995 O1 (Hale-Bopp). The Hale-Bopp crystalline residual better matches equant, b-platelets, c-platelets, and b-columns spectral shape classes, while a-platelets, a-columns and c-columns worsen the

spectral fits.

Forsterite condensation and partial evaporation experiments demonstrate that environmental temperature and grain shape are connected [18-20]. Thus, grain shape is a potential probe for protoplanetary disk temperatures where the cometary crystalline forsterite formed. The forsterite crystal shapes (equant, b-platelets, c-platelets, b-columns – excluding a- and c-columns) derived from our modeling [17] of comet Hale-Bopp, compared to laboratory synthesis experiments [18], suggests that these crystals are high temperature condensates.

By observing and modeling the crystalline features in comet ISON, we may constrain forsterite crystal shape(s) and link to their formation temperature(s) and environment(s).

#### References:

- [1] Campins, H., Ryan, E.V. 1989. ApJ, 341, 1059
- [2] Crovisier, J., et al. 1997. Science, 275, 1904
- [3] Wooden, D.H., et al. 1999. ApJ, 517, 1034
- [4] Wooden, D.H., et al. 2004. ApJL, 612, L77
- [5] Harker, D.E., et al. 2002. ApJ, 580, 579
- [6] —. 2004, ApJ, 615, 1081
- [7] Lisse, C.M., et al. 2006. Icarus 195, 941–944.
- [8] Lisse, C.M., et al. 2007. Icarus 191, 223–240.
- [9] Kelley, M.S., et al. 2010, LPSC, 41, #2375
- [10] Harker, D.E., et al. 2011, AJ, 141, 26
- [11] Draine, B.T., & Flatau, P.J. 1994, J. Opt. Soc. Am. A, 11, 1491
- [12] Draine, B.T., & Flatau, P.J. 2008, J. Opt. Soc. Am. A, 25, 2693
- [13] Fabian, D., et al., 2001, A&A, 378, 228
- [14] Tamanai, A., et al. 2006. ApJ, 648, L147
- [15] Tamanai, A., et al. 2009. ASP Conf. Ser., 414, 438
- [16] Koike, C., et al. 2010. ApJ, 709, 983
- [17] Lindsay, S.S., et al. 2013, ApJ, 766, 54
- [18] Tsuchiyama, A. 1998. Mineralogical J., 20, 59
- [19] Kobatake, H., et al., 2008. Icarus, 198, 208
- [20] Takigawa, A., et al.. 2009. ApJL, 707, L97

**INDEX TERMS:** 6015 PLANETARY SCIENCES: COMETS AND SMALL BODIES Dust, 6005 PLANETARY SCIENCES: COMETS AND SMALL BODIES Atmospheres, 5709 PLANETARY SCIENCES: FLUID PLANETS Composition, 6040 PLANETARY SCIENCES: COMETS AND SMALL BODIES Origin and evolution.

(No Image Selected)

(No Table Selected)

#### Additional Details

**Previously Presented Material:** 50% Lindsay et al. 2013, ApJ, 766, 54  
95% GRC July 2013 (Gordon Research Conference)

#### Contact Details

**CONTACT (NAME ONLY):** Diane Wooden

**CONTACT (E-MAIL ONLY):** [diane.h.wooden@nasa.gov](mailto:diane.h.wooden@nasa.gov)

**TITLE OF TEAM:**

ScholarOne Abstracts® (patent #7,257,767 and #7,263,655). © [ScholarOne](#), Inc., 2013. All Rights Reserved.  
ScholarOne Abstracts and ScholarOne are registered trademarks of ScholarOne, Inc.



Follow ScholarOne on Twitter

[Terms and Conditions of Use](#)

Product version number 4.2.0 (Build 45)  
Build date Aug 05, 2013 14:55:26. Server tss1be0013