


# Conjunction Assessment Risk Analysis



## Atmospheric Model at JSpOC



**D. Pachura, M. Hejduk**



**13 Apr 2016**




# Atmospheric Model Introduction


- **Atmospheric model used for ASW predictions composed of several stages working in sync**
  - JB2008 Model
    - Jacchia legacy
    - Driven by solar irradiance in extreme to far ultraviolet wavelengths
  - HASDM Dynamic Calibration Atmosphere
    - Uses calibration satellites to compute corrections to base atmosphere
    - Provides definitive atmosphere and 3 day predicted corrections
  - Anemomilos Dst Prediction Model
    - Provides 6 day prediction of Dst
    - Empirical relationships driven by x-ray flare observations
  - Solar Indices
    - Provides base inputs for model










# Geomagnetic Correction






# Anemomilos Dst Prediction






# Solar Events


- **Solar events drive the geomagnetic correction to the model**
  - When the Disturbance Storm Time (Dst) measure is above -75 nT, model uses Jacchia's  $a_p$  based temperature correction
  - For Dst less than -75 nT, model applies Dst based equations and integrates over the storm to find temperature correction
- **Anemomilos model currently providing predictive Dst inputs to JSpOC model**
  - Watches for X-ray flares
  - Determines if event is geoeffective
  - Applies template based on magnitude of X-ray flux
  - Does not currently update storm prediction




# Solar Event Timeline







# Solar Event Timeline cont.



--- Prediction  
 --- Data

Notes:  
 Storm not to scale

