



Langley
Research
Center



Improving climate projections using “intelligent” ensembles

Noël C. Baker and Patrick C. Taylor
NASA Postdoctoral Program


Presented at the AGU Joint Assembly in Montréal, Canada
May 5, 2015

CMIP5: group of ~45 models

CMIP5 Model

- BCC-CSM1.1
- BCC-CSM1.1.m
- BNU-ESM
- CanCM4
- CanESM2
- CCSM4
- CESM1-BGC
- CESM1-CAM5
- CESM1-WACCM
- CMCC-CESM
- CMCC-CM
- CMCC-CMS
- CNRM-CM5
- ACCESS1.0
- ACCESS1.3
- CSIRO-Mk3.6.0
- EC-EARTH
- FGOALS-g2
- FGOALS-s2
- FIO-ESM
- GFDL-CM3
- GFDL-ESM2G
- GFDL-ESM2M
- GISS-E2-H
- GISS-E2-H-CC
- GISS-E2-R
- GISS-E2-R-CC
- HadCM3
- HadGEM2-AO
- HadGEM2-CC
- HadGEM2-ES
- INM-CM4
- IPSL-CM5A-LR
- IPSL-CM5A-MR
- IPSL-CM5B-LR
- MIROC4h
- MIROC5
- MIROC-ESM
- MIROC-ESM-CHEM
- MPI-ESM-LR
- MPI-ESM-MR
- MPI-ESM-P
- MRI-CGCM3
- NorESM1-M
- NorESM1-ME

Models are averaged together to make climate predictions




Predicted temperature changes


IPCC AR5 Ch.12

But models can have a large spread in predictions, and individual models can perform very differently from observations

GLOBAL TEMPERATURES: comparing CMIP5 & HadCRUT4



The traditional **Multi-Model Ensemble Approach** uses the model mean to provide an improved “best estimate” forecast




IPCC AR5 Figure SPM.7

The multi-model ensemble generally performs better than individual models

Example: I^2 performance index (Reichler and Kim 2008)


Calculates aggregated model errors relative to NCEP/NCAR reanalyses for multiple climate variables



The multi-model ensemble generally performs better than individual models


Example: I^2 performance index (Reichler and Kim 2008)

Calculates aggregated model errors relative to NCEP/NCAR reanalyses for multiple climate variables




Some models perform better than others:

Can we use knowledge of model performance for a better way to combine model output?



The “intelligent ensemble” approach for creating multi-model ensemble projections



Project goal:
determine future climate state
using observed current climate
and an ensemble of models

$$f(x_{obs}) = \Delta x$$

Observed
climate

Future
climate
state

Previous work has explored model performance and some unequal-weighting metrics

Several examples:

- Use only subsets of models (USGCRP 2009)
- Create mean-state metrics using model skill (Giorgi and Mearns 2002, 2003; Reichler and Kim 2008)
- Constrain model projections using mean-state CERES data (Tett et al. 2013)
- Weight using regression between observed and future trends (Boe et al 2009)
- Apply bias correction for present-day to future trends (Baker and Huang 2012)

“The community would benefit from a larger set of proposed methods and metrics” (Knutti 2010)

This project tests new climate model performance metrics

Radiation budget quantities:

- Top-of-atmosphere (TOA) longwave (LW) and shortwave (SW) radiation fluxes
- Surface LW and SW radiation fluxes
- Surface temperature

New process-oriented metrics:

- $$\frac{\Delta TOA \text{ Radiation flux}}{\Delta \text{ Surface temperature}}$$

Statistical tests:

- F-test for equal variances
- Kolmogorov-Smirnov test for distribution similarity
- Earth Mover's Distance (EMD): test for area of distribution overlap
- Local Variance: test variance of first difference time series (Baker and Taylor 2015)

Model data:

35 CMIP5 models

<http://pcmdi9.llnl.gov/>

- ‘Pre-Industrial Control’ simulations (monthly mean, 100 years) to create metric weights
- ‘RCP 8.5’ future simulations (monthly mean, 2081-2100 minus 2011-2030 to produce 21st-century trends)

Observational datasets:

NASA CERES EBAF-TOA and surface monthly global-mean


<http://ceres.larc.nasa.gov/>

NASA GISS Surface Temperature Analysis (GISTEMP)


<http://data.giss.nasa.gov/gistemp/>

Step 2: Using skill-subset of models, apply “perfect model” approach (Räisänen and Palmer 2001)

Create set of potential “Earths” each with a continuous time series of observations




- 1 OLR all-sky variance test
- ▲ 2 OLR all-sky K-S test
- ▼ 3 OLR all-sky local variance test
- 4 OLR all-sky EMD value
- 5 OLR cloudy-sky variance test
- ▲ 6 OLR cloudy-sky K-S test
- ▼ 7 OLR cloudy-sky local variance test
- 8 OLR cloudy-sky EMD value
- 9 OLR clear-sky variance test
- ▲ 10 OLR clear-sky K-S test
- ▼ 11 OLR clear-sky local variance test
- 12 OLR clear-sky EMD value
- 13 SW all-sky variance test
- ▲ 14 SW all-sky K-S test
- ▼ 15 SW all-sky local variance test
- 16 SW all-sky EMD value
- 17 SW cloudy-sky variance test
- ▲ 18 SW cloudy-sky K-S test
- ▼ 19 SW cloudy-sky local variance test
- 20 SW cloudy-sky EMD value
- 21 SW clear-sky variance test
- ▲ 22 SW clear-sky K-S test
- ▼ 23 SW clear-sky local variance test
- 24 SW clear-sky EMD value
- 25 Surface temperature variance test
- ▲ 26 Surface temperature K-S test
- ▼ 27 Surface temperature local variance test
- 28 Surface temperature EMD value
- 29 OLR/Ts variance test
- ▲ 30 OLR(cloudy-sky)/Ts variance test
- ▼ 31 OLR/Ts K-S test
- 32 OLR(cloudy-sky)/Ts K-S test
- ◆ 33 OLR Ts regression means test
- ▶ 34 OLR(cloudy-sky) Ts regression means test
- 35 SW/Ts variance test
- ▲ 36 SW(cloudy-sky)/Ts variance test
- ▼ 37 SW/Ts K-S test
- 38 SW(cloudy-sky)/Ts K-S test
- ◆ 39 SW Ts regression means test
- ▶ 40 SW(cloudy-sky) Ts regression means test
- 41 Equal-weighted mean



Model mean I^2 performance index


Metric performance and consistency is correlated:

Metrics which best reduce error in future projections behave similarly across model ensemble




Results: new 21st-century projections (surface temperature)

"Intelligent" ensemble mean temperature trend (°C)




Difference between "Intelligent" and Equal-weight ensemble means (°C)




Results: new 21st-century projections (precipitation)

"Intelligent" ensemble mean precipitation trend (cm/year)




Difference between "Intelligent" and Equal-weight ensemble means (cm/year)




Results: new 21st-century projections (surface downward SW radiation)

"Intelligent" ensemble mean surface shortwave radiation trend (W/m²)




Difference between "Intelligent" and Equal-weight ensemble means (W/m²)




Results: new 21st-century projections (TOA LW radiation)

"Intelligent" ensemble mean outgoing longwave radiation trend (W/m²)




Difference between "Intelligent" and Equal-weight ensemble means (W/m²)




Results: new 21st-century projections (TOA SW radiation)

"Intelligent" ensemble mean reflected shortwave radiation trend (W/m^2)




Difference between "Intelligent" and Equal-weight ensemble means (W/m^2)




Results: new 21st-century projections (regional-mean weights)

"Intelligent" ensemble mean temperature trend (°C)




Difference between "Intelligent" and Equal-weight ensemble means (°C)




Results: new 21st-century projections (regional-mean weights)

"Intelligent" ensemble mean precipitation trend (cm/year)




Difference between "Intelligent" and Equal-weight ensemble means (cm/year)




Results: new 21st-century projections (regional-mean weights)


"Intelligent" ensemble mean surface shortwave radiation trend (W/m^2)



Difference between "Intelligent" and Equal-weight ensemble means (W/m^2)




USDA Farm Resource Regions (1° resolution)




Results: new 21st-century projections

"Intelligent" ensemble mean temperature trend (°C)




US mean temperature increase: 3.9 °C




- Basin and Range: 3.9 °C
- Fruitful Rim: 3.4 °C
- Prairie Gateway: 3.8 °C
- Northern Great Plains: 4.1 °C
- Heartland: 4.1 °C
- Northern Crescent: 4.3 °C
- Eastern Uplands: 3.8 °C
- Southern Seaboard: 3.5 °C
- Mississippi Portal: 3.6 °C

Results: new 21st-century projections

"Intelligent" ensemble mean precipitation trend (cm/year)





US mean precipitation increase: 3.4 cm/year




- Basin and Range: 0.6 cm/year
- Fruitful Rim: 0.8 cm/year
- Prairie Gateway: -1.8 cm/year
- Northern Great Plains: 2.7 cm/year
- Heartland: 7.2 cm/year
- Northern Crescent: 9.1 cm/year
- Eastern Uplands: 6.8 cm/year
- Southern Seaboard: 6.8 cm/year
- Mississippi Portal: 5.4 cm/year

Results: new 21st-century projections

"Intelligent" ensemble mean surface shortwave radiation trend (W/m²)



US mean decrease in surface solar radiation: -0.33 Watts/m^2



- Basin and Range: -2.4 Watts/m^2
- Fruitful Rim: -0.5 Watts/m^2
- Prairie Gateway: 0.7 Watts/m^2
- Northern Great Plains: -1.9 Watts/m^2
- Heartland: 0.7 Watts/m^2
- Northern Crescent: -0.1 Watts/m^2
- Eastern Uplands: 2.7 Watts/m^2
- Southern Seaboard: 2.5 Watts/m^2
- Mississippi Portal: 2.6 Watts/m^2