

Matthew S. Johnson
NCTS #25415-17 AGU Fall Meeting 2016
12-16 Dec. 2016
Presentation Date: Thursday, 15 December 2016

Title:

Evaluating A Priori Ozone Profile Information Used in TEMPO Tropospheric Ozone Retrievals

Abstract:

Ozone (O₃) is a greenhouse gas and toxic pollutant which plays a major role in air quality. Typically, monitoring of surface air quality and O₃ mixing ratios is primarily conducted using in situ measurement networks. This is partially due to high-quality information related to air quality being limited from space-borne platforms due to coarse spatial resolution, limited temporal frequency, and minimal sensitivity to lower tropospheric and surface-level O₃. The Tropospheric Emissions: Monitoring of Pollution (TEMPO) satellite is designed to address these limitations of current space-based platforms and to improve our ability to monitor North American air quality. TEMPO will provide hourly data of total column and vertical profiles of O₃ with high spatial resolution to be used as a near-real-time air quality product.

TEMPO O₃ retrievals will apply the Smithsonian Astrophysical Observatory profile algorithm developed based on work from GOME, GOME-2, and OMI. This algorithm uses a priori O₃ profile information from a climatological data-base developed from long-term ozone-sonde measurements (tropopause-based (TB) O₃ climatology). It has been shown that satellite O₃ retrievals are sensitive to a priori O₃ profiles and covariance matrices. During this work we investigate the climatological data to be used in TEMPO algorithms (TB O₃) and simulated data from the NASA GMAO Goddard Earth Observing System (GEOS-5) Forward Processing (FP) near-real-time (NRT) model products. These two data products will be evaluated with ground-based lidar data from the Tropospheric Ozone Lidar Network (TOLNet) at various locations of the US. This study evaluates the TB climatology, GEOS-5 climatology, and 3-hourly GEOS-5 data compared to lower tropospheric observations to demonstrate the accuracy of a priori information to potentially be used in TEMPO O₃ algorithms. Here we present our initial analysis and the theoretical impact on TEMPO retrievals in the lower troposphere.

Authors

- [Matthew S Johnson](#)
 - NASA Ames Research Center
- [John T Sullivan](#)
 - University of Maryland Baltimore County
- [Xiong Liu](#)
 - Harvard-Smithsonian Center for Astrophysics
- [Mike Newchurch](#)
 - University of Alabama in Huntsville
- [Shi Kuang](#)

- Univ. of Alabama in Huntsville
- [Thomas J McGee](#)
 - NASA Goddard Space Flight Center
- [Andrew O'Neil Langford](#)
 - NOAA ESRL
- [Christoph J Senff](#)
 - NOAA Boulder
- [Thierry Leblanc](#)
 - California Institute of Technology
- [Timothy Berkoff](#)
 - NASA Langley Research Center
- [Guillaume Gronoff](#)
 - Science Systems and Applications, Inc. Hampton
- [Gao Chen](#)
 - NASA Langley Research Ctr
- [Kevin B Strawbridge](#)
 - Environment Canada Toronto