

NASA Plant Research for Life Support in Space

Raymond M. Wheeler

*NASA Exploration Research and Technology Directorate
Kennedy Space Center, Florida, USA*

IPSAM, June 2017, Limerick

Human Life Support Requirements:

Inputs

	Daily Rqmt.	(% total mass)
Oxygen	0.83 kg	2.7%
Food	0.62 kg	2.0%
Water (drink and food prep.)	3.56 kg	11.4%
Water (hygiene, flush laundry, dishes)	26.0 kg	83.9%
TOTAL	31.0 kg	


Outputs

	Daily	(% total mass)
Carbon dioxide	1.00 kg	3.2%
Metabolic solids	0.11 kg	0.35%
Water (metabolic / urine)	29.95 kg	96.5%
(hygiene / flush)		12.3%
(laundry / dish)		24.7%
(latent)		55.7%
		3.6%
TOTAL	31.0 kg	


Source: NASA SPP 30262 Space Station ECLSS Architectural Control Document
 Food assumed to be dry except for chemically-bound water.

International Space Station Life Support Systems


Source: Jim Reuter, NASA MSFC



Plants for “Bioregenerative” Life Support



Bioregenerative Life Support Testing Around the World



Crop Considerations for Space

- High yielding and nutritious (CHO, protein, fat, micronutrients)
- High harvest index (edible / total biomass)
- Horticultural requirements
 - planting, watering, harvesting, pollination, propagation
- Environmental requirements
 - lighting, temperature, mineral nutrition, CO₂
- Processing requirements
- Dwarf or low growing types

Some Crops for Life Support

Tibbitts and Alford ^a	Hoff, Howe, and Mitchell ^b	Salisbury and Clark ^c	Russian BIOS-3 Testing ^d
Wheat Soybean Potato Lettuce Sweetpotato Peanut Rice Sugar Beet Pea Taro Winged Bean Broccoli Onion Strawberry	Wheat Potato Soybean Rice Peanut Dry Bean Tomato Carrot Chard Cabbage	Wheat Rice Sweetpotato Broccoli Kale Lettuce Carrot Canola Soybean Peanut Chickpea Lentil Tomato Onion Chili Pepper	Wheat Potato Carrot Radish Beet Nut Sedge Onion Cabbage Tomato Pea Dill Cucumber Salad spp.

^a Tibbitts and Alford (1982); ^b Hoff, Howe, and Mitchell (1982); ^c Salisbury and Clark (1996);

^d Gitelson and Okladnikov (1994)—diet also included supplemental animal protein and sugar.

Crop Selection and Breeding for Space (Utah State University)



Selection of Existing Rice Genotypes



Wheat Breeding for Short Growth and High Harvest Index



'Apogee' Wheat

'Perigee' Wheat




Genetic Engineering Tools



Overexpression of FT flowering gene in plums (USDA researchers) resulted in dwarf growth habit and early flowering

Water and Nutrients for Growing Crops

Recirculating Hydroponics




*Conserve Water & Nutrients
Eliminate Water Stress
Optimize Mineral Nutrition
Facilitate Harvesting*

Root Zone Crops in Nutrient Film Technique (NFT)




Fig. 7


Evapotranspiration from Plant Stand (potato)



Water, Nutrient, and pH Control



High Yields from NASA Sponsored Studies




*Wheat - 3-4 x World Record
Potato - 2 x World Record
Lettuce-Exceeded Commercial
Yield Models*




Bubgee, B.G. and F.B. Salisbury. 1988. Plant Physiol. 88:869-878.

Wheeler, R.M., T.W. Tibbitts, A.H. Fitzpatrick. 1991. Crop Science 31:1209-1213.


Canopy CO₂ Uptake / O₂ Production (20 m² Soybean Stand)




CO₂ Exchange Rates of Soybean Stands



Effect of CO₂ Concentration on Photosynthesis (potato)



Canopy / Stand Ethylene Production



Ethylene in Closed Systems



Epinastic
Wheat Leaves
at ~120 ppb




Epinastic
Potato Leaves
at ~40 ppb

NASA's Biomass Production Chamber (BPC)

Early Vertical Agriculture !

External View - Back



20 m² growing area; 113 m³ vol.; 96 400-W HPS Lamps;
400 m³ min⁻¹ air circulation; two 52-kW chillers

Control Room




Hydroponic System

NASA's Biomass Production Chamber (BPC)




Wheat

(Triticum aestivum)




planting



harvest


Soybean

(Glycine max)



Lettuce


(Lactuca sativa)





Potato

(*Solanum tuberosum*)



Automation Technologies for CEA



ALSARM Robot in NASA
Biomass Production Chamber



Electric Lighting Considerations

<i>Lamp Type</i>	<i>Conversion* Efficiency</i>	<i>Lamp Life* (hrs)</i>	<i>Spectrum</i>
• Incandescent/Tungsten**	5-10%	2000	Intermd.
• Xenon	5-10%	2000	Broad
• Fluorescent***	20%	5,000-20,000	Broad
• Metal Halide	25%	20,000	Broad
• High Pressure Sodium	30-35%	25,000	Intermd.
• Low Pressure Sodium	35%	25,000	Narrow
• Microwave / RF Sulfur	35-40%+	?	Broad
• LEDs (white)	30-40%	50,000 ?	Broad
• LEDs (red and blue)****	>40%	50,000 ?	Narrow

* *Approximate values.*

** *Tungsten halogen lamps have broader spectrum.*

*** *For VHO lamps; lower power lamps with electronic ballasts last up to ~20,000 hrs.*

**** *State-of-Art Blue and Red LEDs most efficient.*

Light Emitting Diodes (LEDs)

Red...photosynthesis

Blue...photomorphogenesis

Green...human vision



*North American Patent for Using LEDs to
Grow Plants Developed with NASA Funding
at University of Wisconsin !*

Solar Collector / Fiber Optics For Plant Lighting




2 m² of collectors on solar tracking drive (NASA Kennedy Space Center, Florida)


Up to 400 W light delivered to chamber
(40-50% of incident light)
Takashi Nakamura, Physical Sciences Inc.




Light and Crop Yield




*Solar Collectors
for Crop Production*



*Buried Plant
Growth Chambers*



University of Arizona Lunar / Mars Greenhouse



Deployable Mars Greenhouse - Low Pressure Systems



Photosynthetic Radiation at Mars Surface over 2 Martian Years *(J. Clawson, Dissertation 2006)*





Figure 11 The daily integrated total, direct, and diffuse PPF versus latitude and Martian Sol for two Mars years. The labeled sols correspond to the start of each season on Mars. For example, sol 150 corresponds to the Northern Autumnal equinox.

Hypobaric Testing with Plants


Testing at:
NASA KSC
Univ. of Guelph
Texas A&M Univ. ⇒
Univ. of Florida





Lettuce, radish, and wheat plants exposed to rapid pressure drop (27 days old)

Phase Change of Water



Some other Benefits of Plants in Space




- Fresh Foods to Augment the Diet
 - Colors*
 - Texture*
 - Flavor*
 - Nutrients*
- Bright Light
- Aromas
- Gardening Activity

Plant Chamber at US South Pole Station






Plants and Human Well-Being




Testing Crops in Human Habitats



Plant Testing on the International Space Station—VEGGIE Plant Chamber



Sequential Development for Space Agriculture



Some Lessons Learned from NASA CEA Research

- 20-25 m² of crops could provide all the O₂ for one person, and 40-50 m² all of the food (dietary energy)
- Better adapted crops are needed—short growth, high harvest index, improved nutrition—Use genetic engineering?
- Lighting is key to sustaining high yields
- CEA systems require large quantities of water (e.g., 5 L m⁻² d⁻¹) and this water must be recycled.
- Up to 90 kg of fertilizer would needed per person per year, emphasizing the need for recycling nutrients.
- Plants can provide psychological benefits to humans—this needs further study.
- The use of plants for life support will likely evolve sequential, starting with small, supplemental food production and expanding for future missions.

Plants and Living in Space

A man in a blue hoodie is crouching in the center of a large, circular, geodesic dome structure. The interior is filled with rows of green plants growing in red soil. The structure has a complex metal framework and translucent walls. A tall, cylindrical apparatus is visible in the background.

As we explore sustainable living for space, we will learn more about sustainable living on Earth

KSC Advanced Life Support Team, Hangar L, KSC 1994



One of our Kennedy Space Center Researchers !



Michelle McKeon-Bennett, 2004, Space Life Science Laboratory, KSC, Florida

Effect of Light on Productivity and Crop Area Requirements

