

A CASE FOR NEBULA SCALE MIXING BETWEEN NON-CARBONACEOUS AND CARBONACEOUS CHONDRITE RESERVOIRS: TESTING THE GRAND TACK MODEL WITH CHROMIUM ISOTOPIC COMPOSITION OF ALMAHATA SITTA STONE 91A

M. E. Sanborn¹, Q.-Z. Yin¹, C. A. Goodrich², M. Zolensky³, and A. M. Fioretti.⁴ ¹Department of Earth and Planetary Sciences, University of California at Davis, Davis, CA 95616 USA (mesanborn@ucdavis.edu), ²Lunar and Planetary Institute, Houston, TX, 77058 USA, ³ARES, NASA JSC, Houston, TX, USA, ⁴CNR, Padova, Italy.

Introduction: There is an increasing number of Cr-O-Ti isotope studies [1-6] that show solar system materials are divided into two main populations, one carbonaceous chondrite (CC)-like and the other is non-carbonaceous (NC)-like, with minimal mixing attributed to a gap opened in the protoplanetary disk due to Jupiter's formation [5,7,8]. The Grand Tack model [9] suggests there should be large-scale mixing between S- and C-type asteroids, an idea supported by our recent work on chondrule $\Delta^{17}\text{O}-\epsilon^{54}\text{Cr}$ isotope systematics [10].

The Almahata Sitta (AhS) meteorite provides a unique opportunity to test the Grand Tack model. The meteorite fell to Earth in October 2008 and has been linked to the asteroid 2008 TC₃ which was discovered just prior to the fall of the AhS stones [11-13]. The AhS meteorite is composed of up to 700 individual pieces with ~140 of those pieces having some geochemical and/or petrologic studies [14]. Almahata Sitta is an anomalous polymict ureilite with other meteorite components, including enstatite, ordinary, and carbonaceous chondrites with an approximate abundance of 70% ureilites and 30% chondrites [14,15]. This observation has led to the suggestion that TC₃ 2008 was a loosely aggregated rubble pile-like asteroid with the non-ureilite sample clasts within the rubble-pile (e.g., [14,15] and references therein). Due to the loosely-aggregated nature of AhS, the object disintegrated during atmospheric entry resulting in the weakly held clasts falling predominantly as individual stones in the AhS collection area. However, recent work [16] has identified one sample of AhS, sample 91A, which may represent two different lithologies co-existing within a single stone. The predominate lithology type in 91A appears to be that of a C2 chondrite based on mineralogy but also contains olivine, pyroxene, and albite that have ureilite-like compositions.

Previous Cr isotope investigations into AhS stones are sparse and what data is available show nearly uniform isotopic composition similar to that of typical ureilites with negative $\epsilon^{54}\text{Cr}$ values [17,18].

Methods: The aliquot for Cr isotopic analysis (24.77 mg) was prepared from a small chip of material from Almahata Sitta 91A. Chromium was separated from the sample matrix using a three-column chromatography procedure [19]. The Cr isotopic composition was measured using a Thermo *Triton Plus* thermal ionization mass spectrometer at the University of California, Davis.

Results and Discussion: The Cr isotopic composition of AhS 91A shows clearly resolved excesses in both ^{53}Cr and ^{54}Cr with an $\epsilon^{53}\text{Cr} = +0.09 \pm 0.05$ and $\epsilon^{54}\text{Cr} = +1.83 \pm 0.08$. This is the highest $\epsilon^{54}\text{Cr}$ value observed thus far in any CC-like reservoir. In fact it is the highest $\epsilon^{54}\text{Cr}$ in any bulk planetary materials in hand. This $\epsilon^{54}\text{Cr}$ is clearly resolved from ureilites that have negative $\epsilon^{54}\text{Cr} \sim -0.90$ [18] as the extreme end member of the NC reservoir in terms of $\epsilon^{54}\text{Cr}$ value. The difference indicates that the AhS 91A fragment originated from a Cr isotopic reservoir distinct from ureilites. The recent work studying the bulk mineralogy inferred that AhS 91A was likely comprised of material similar to C2 chondrites [16]. However, the $\epsilon^{54}\text{Cr}$ of AhS 91A is higher than CI, CM, and CR chondrites and has a $\epsilon^{54}\text{Cr}$ composition higher than any observed in any carbonaceous materials ([4,6] and references therein). This would indicate that the material that comprises AhS 91A is not typical CI-like material and instead is a carbonaceous chondrite material that we have yet to sample in our collections as a separate meteorite. Oxygen isotopes are still pending, but even if oxygen isotopes were similar to CI, CM, or CR chondrites or another carbonaceous chondrite group, the difference in $\epsilon^{54}\text{Cr}$ would indicate a unique source. The observation of a different $\epsilon^{54}\text{Cr}$ in AhS 91A compared to ureilites or the other AhS samples [17] has implications for large scale nebular wide mixing between NC and CC-like reservoirs, as suggested by the Grand Tack model [9]. This is supported by a rubble-pile like nature of the AhS asteroid.

Acknowledgements: We thank M. H. Shaddad and P. Jenniskens for providing the sample of AhS 91A.

References: [1] Yin Q.-Z. et al. (2009) *LPS XL*, A2006. [2] Warren P. H. (2011) *EPSL* 311:93-100. [3] Sanborn M. E. et al. (2014) *Goldschmidt Conf.*, A2171. [4] Sanborn M. E. et al. (2015) *LPS XLVI*, A2259. [5] Williams C. D. et al. (2016) *Goldschmidt Conf.*, A3415. [6] Yin Q.-Z. et al. (2017) *LPS XLVIII*, A1771. [7] Morbidelli A. et al. (2015) *Icarus* 258:418-429. [8] Andrews S. M. et al. (2016) *ApJ* 820:L40. [9] Walsh K. J. et al. (2011) *Nature* 475:206-209. [10] Defouilloy C. (2016) *Goldschmidt Conf.*, A629. [11] Jenniskens P. et al. (2009) *Nature* 458:485-488. [12] Jenniskens P. et al. (2010) *MAPS*. 45:1590-1617. [13] Shaddad M. H. et al. (2010) *MAPS* 45:1557-1589. [14] Horstmann M. and Bischoff A. (2014) *Chemie der Erde* 74:149-183. [15] Goodrich C. A. et al. (2015) *MAPS* 50:782-809. [16] Goodrich C. A. (2017) *80th MetSoc*, this meeting. [17] Qin L. et al. (2010) *MAPS* 45:1771-1777. [18] Yamakawa A. et al. (2010) *ApJ*. 720:150-154. [19] Yamakawa A. et al. (2009) *Anal. Chem.* 81:9787-9794.