
Analysis of an Anemone-Type Magnetic Field in an On-Disk Coronal Hole



M.L. Adams¹, N.K. Panesar², R.L. Moore³, A.F. Tennant¹, R. Woolley⁴

- 1. NASA/Marshall Space Flight Center
- 2. NASA/USRA
- 3. University of Alabama in Huntsville
- 4. Bass Chandler School (Arizona)


Overview

- Introduction -- What is a coronal hole?
 - -- What is an Anemone Event?

- Data
- Analysis
- Summary
- Future Work

What is a Coronal Hole?

SDO AIA 193 Å



Seen
anytime
in solar
cycle,
more
common
in
declining
phase

Sun's field

Darker,


Cooler

Less dense,


open to

interplanetary medium, high speed solar wind > 400km/s


Coronal Hole "Now"



Coronal Hole and Anemone Event




DATA




DATA


Solar Dynamics Observatory's Atmospheric Imaging Assembly



and

Helioseismic and Magnetic Imager






ANALYSIS


HMI line-of-sight component

and Continuum




Before


Emerging Flux




Appearance of Pores 20:06:27



Further Development of Region





Interchange-Reconnection Schematic




SUMMARY

We observed the birth of a bipolar solar active region in a small coronal hole with obvious field-emergence driven interchange magnetic reconnection without obvious jets...








FUTURE WORK

- Finish analysis with 211 Å data



- Identify more events like this one for multi-wavelength study