

GeoDash: Assisting Visual Image Interpretation in Collect Earth Online by Leveraging Big Data on Google Earth Engine


Kel Markert^{1,2}, William Ashmall^{1,3}, Gary Johnson⁴, David Saah^{4,5}, Danilo Mollicone⁶, Alfonso Sánchez-Paus Díaz⁶, Eric Anderson^{1,2}, Africa Flores^{1,2}, Robert Griffin⁷ ¹NASA-SERVIR, Science Coordination Office; ²Earth System Science Center, University of Alabama in Huntsville; ³Science and Technology Institute, University of Alabama in Huntsville; ³Science and Technology Institute, University of Alabama in Huntsville ³Science and Technology Institute, University of Alabama in Huntsville; ³Science and Technology Institute, University of Alabama in Huntsville; ³Science and Technology Institute, University of Alabama in Huntsville; ³Science and Technology Institute, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, University of Alabama in Huntsville; ⁴Spatial Informatics Group; ⁵Geospatial Analysis Lab, ⁵Geospatial Analysis Lab, ⁵Geospatial Analysis Lab, ⁵Geospatial Analysis Lab, ⁵Geospatial Ana Session: IN23C: Browser-Based Mapping, Space-Time Visualization, and Analysis of Big Earth Science Data I; AGU Fall 2017, 12 Dec. 2017; New Orleans, LA

Results

Abstract

Collect Earth Online (CEO) is a free and open online implementation of the FAO Collect Earth system for collaboratively collecting environmental data through the visual interpretation of Earth observation imagery. The primary collection mechanism in CEO is human interpretation of land surface characteristics in imagery served via Web Map Services (WMS). However, interpreters may not have enough contextual information to classify samples by only viewing the imagery served via WMS, be they high resolution or otherwise. To assist in the interpretation and collection processes in CEO, SERVIR, a joint NASA-USAID initiative that brings Earth observations to improve environmental decision making in developing countries, developed the GeoDash system, an embedded and critical component of CEO. GeoDash leverages Google Earth Engine (GEE) by allowing users to set up custom browser-based widgets that pull from GEE's massive public data catalog. These widgets

Methodology



can be quick looks of other satellite imagery, time series graphs of environmental variables, and statistics panels of the same. Users can customize widgets with any of GEE's image collections, such as the historical Landsat collection with data available since the 1970s, select date ranges, image stretch parameters, graph characteristics, and create custom layouts, all on-the-fly to support plot interpretation in CEO. This presentation focuses on the implementation and potential applications, including the back-end links to GEE and the user interface with custom widget building. GeoDash takes large data volumes and condenses them into meaningful, relevant information for interpreters. While designed initially with national and global forest resource assessments in mind, the system will complement disaster assessments, agriculture management, project monitoring and evaluation, and more.

CEO Web

Front End

Back End

FAO Collect Database

GEE

GEE Image Collections

Collect Earth Online promotes consistency in locating, interpreting, and labeling reference data plots for use in classifying and monitoring land cover / land use change. The full functionality of Collect Earth Online including collaborative compilation of reference point databases, using the FAO collect system, is implemented online so there is no need for desktop installation. The GeoDash system is integrated into CEO where collection plots are inputs into the GEE queries for display. The GeoDash system being separate allows for building custom widgets the leverage Earth Engine for additional applications such as forest monitoring.

Conclusions

- CEO facilitates collaboratively collecting reference data within a free and open online system.
- GeoDash is a flexible system that leverages GEE for custom widget creation to access and view petabytes of data in meaningful information.

Acknowledgements

This work is coordinated with SilvaCarbon and the FAO. Support provided through NASA Cooperative Agreement NNM11AA01A. The authors would like to thank the many people that have contributed to the work, especially Dan Irwin (NASA SERVIR), Ashutosh Limaye (NASA SERVIR), Ray French (NASA SERVIR), Karis Tenneson (USFS/GTAC), and many others. A special thanks goes to Google and the Earth Engine Team for developing and providing the Earth Engine platform. A special thanks goes to Nick Clinton (Google) for technical support on Earth Engine. High resolution data used within CEO is provided by DigitalGlobe.

Project Partners







Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Aeronautics and Space Administration (NASA). Any







Fig. 2 Example of the CEO Collection Page for users to assign land cover sub-plot samples.



Fig. 3 Example of the GeoDash Page for users to assess additional information gleaned from Landsat. Index values in maps or time series to be viewed to identify, for example, seasonality.



PROJECT STATS Project	kel.markert@nasa.gov Madison County Coll	lection) 111	ECT NE	EAR	TH	
Plots Assigned Plots Flagged	0 (0%) 0 (0%)		llser	r int	erface	ı		
Plots Completed	0 (0%)						\frown	
Plots Total	250			•	es for t		U	
PLOT NAVIGATION			syste	em [·]	for the			
NEXT PLOT SAVE ASSI	GNMENTS FLAG PLOT	AS BAD	inter	rpre	etation	of		
MAGERY OPTIONS				•	image		Ч	
BingAerial		÷			on of r	-		
SAMPLE VALUE: LULC						eleren	Ce	
UKNOWN	WATER		data	•				
FOREST	URBAN							
CROP		NDVI Imag	ne Collecti	on				
types to	EVI Image Collection EVI Graph EVI2 Image Collection EVI2 Graph NDMI Image Collection NDMI Graph NDWI Image Collection NDWI Graph Statistics Custom widget					×		
		NDVI Map)	ange y	/ou would	like]	
		2015-01-0)1	to	2016-12	-31	ר	
				.0	2010 12			
		Columns 12						
NDWI			С	reate	widget	Cancel		
u ⁿ		סען e plot area t	o zoom in					
om	Area 1.02043032890466777 Elevation N/A - N/A m	04303289046677 ha Elevation		Fig. 4 Example GeoDash widget setup. Users can select from pre-selected indices to display in either a map or time series for a selected time period				

This system enables the collection of more information within the CEO systems by allowing users to view data relevant to identifying land cover types.

The exploration for use of CEO and GeoDash for other application such as identification of surface water is on going and will continued to be pursued.

RCMRD

Contact

adpc

Questions? About Project – kel.markert@nasa.gov About SERVIR – servirglobal.net About CEO – ceo.sig-gis.com About OpenForis – openforis.org

Code repositories github.com/openforis/collect-earth-online - github.com/openforis/gee-gateway

ICIMOD