

# From LIMS to OMPS-LP: limb ozone observations for future reanalyses

K. Wargan<sup>1,2</sup>, N. Kramarova<sup>3</sup>, E. Remsberg<sup>4</sup>, L. Coy<sup>1,2</sup>, L. Harvey<sup>5</sup>, N. Livesey<sup>6</sup> and S. Pawson<sup>2</sup>

<sup>1</sup>) SSAI, Lanham, MD, USA, <sup>2</sup>) NASA GSFC, Code 610.1, Greenbelt, MD, USA, <sup>3</sup>) NASA/GSFC, Code 614, Greenbelt, MD, USA, <sup>4</sup>) NASA Langley Research Center, Hampton, VA, USA, <sup>5</sup>) LASP, Boulder, CO, USA, <sup>6</sup>) JPL, Caltech Pasadena, CA, USA

## Summary of Issues

**Ozone poses a unique set of challenges for atmospheric reanalyses.**

**Chemically:** the distribution is controlled by sunlight, stratospheric transport and chemistry including anthropogenic pollutants that rise between 1960 and 1997, then decline after the Montreal Protocol becomes effective.

**Radiatively:** ozone in the upper troposphere and lower stratosphere is a climate gas; it also impacts the use of infrared radiances to constrain the 3D thermal field.

**Observationally:** It is the most widely observed trace gas, yet the observations are inhomogeneous in space and time, especially when information about vertical profiles is needed.

## Characterizing the Observations in Periods of Ozone Decline and Expected Recovery


WMO-UNEP documents the global ozone decline between about 1980 and 1997; this is also captured in chemistry-climate models. Early signs of the projected 21<sup>st</sup> century ozone recovery, as CFCs decline and the stratosphere cools, are evident in satellite observations.

There is a well-documented series of total and partial column ozone data (SBUV, TOMS) for this period of ozone decline. NASA's research observations provide only "snapshots" of the ozone profiles, in 1978-1979 with LIMS and the 1990s with Aura-MLS. Many non-NASA satellite data are also available.

Challenge is to integrate the model, with chemistry, to the observations and to use the assimilation to produce a steady long-term ozone record.


NASA's EOS-Aura MLS so far spans the period 2004-2017. The OMPS-LP (Limb Profiler) observations will continue that record into the late 2020s and beyond.

Here we show two examples of initial integration of LIMS (historical) and OMPS-LP (going forward) ozone observations into the GEOS Data Assimilation System, building on the setup used to produce the MERRA-2 reanalysis, which uses SBUV, OMI and MLS ozone data.



A juxtaposition of past and future ozone change from the WMO-UNEP (2014) assessment and near-global satellite observations of total-column, partial-column, and profile ozone that can be used in reanalyses.

## Example 1: MLS/OMPS-LP Agreement (2016)



## Example 2: Assimilating LIMS ozone (1978-1979 NH winter)

