


Validation of CYGNSS V2 Level 2 Winds

J. Brent Roberts and Timothy J. Lang

NASA Marshall Space Flight Center





Background

- CYGNSS Version 2 Level 2 wind products are a major overhaul from V1, and include improvements in the Level 1 dataset as well as updated geophysical model functions (GMFs)
- Now two GMFs in CYGNSS L2 data:
 - Limited Fetch (LF) appropriate for tropical cyclones
 - Fully Developed Seas (FD) appropriate everywhere else
- Validation was performed for August-September 2017 data against a subset of available buoys and volunteer observing ships



Daily Summary Example

Version 1

Level 2 Wind Speeds - 20170828




Level 2 Wind Speeds - 20170828



Spurious high winds in V1 have been removed in V2

Version 2 FD




Version 1

CYGNSS by PRN Code
(i.e., individual GPS satellites)



V2 has major reductions in differences between different GPS satellites, suggesting better information on transmitter characteristics (e.g., power)

Version 2 FD





CYGNSS vs. Buoys/Ships (scatterplots)

- Good correlation between FD winds and observations
- CYGNSS high bias appears to be related to influence of outliers
- LF winds are biased high due to lack of tropical cyclones in the validation dataset




Comparison to ECMWF+GDAS and Microwave Imager




- Microwave imager winds (derived from GPM constellation L1C dataset) close to CYGNSS, but slightly weaker.
- Some wind variability between different CYGNSS satellites.

~2-Month Average

CYGNSS:FM-ALL WSPD-FD V2-Beta Average




CYGNSS temporal averages in V2 look relatively smooth and realistic



~2-Month Time Series

Domain (~45N/S) Daily Area-Averaged Estimate



CYGNSS domain average time series shows variable offset; typically $< 0.5 \text{ m s}^{-1}$

Conclusions

- CYGNSS V2 Level 2 wind product is a major improvement over V1.
- FD winds close to $\pm 2 \text{ m s}^{-1}$ mission requirement for low winds. LF winds difficult to validate with such a short dataset with few tropical cyclones.
- Some remaining differences between individual GPS and CYGNSS satellites, suggesting additional work needed to make dataset fully self-consistent.
- Good agreement with microwave imagers, although CYGNSS tends to see somewhat stronger winds.