


# **Linearity Analysis and Efficiency Testing of The Chromospheric Lyman-Alpha Spectro-Polarimeter** (CLASP) Science Cameras for Flight



Marshall Space **Flight Center** 

Salma C. Walker<sup>1</sup>, Laurel Rachmeler<sup>2</sup>, Amy Winebarger<sup>2</sup>, Patrick Champey<sup>2,3</sup>, Christian Bethge<sup>2,4</sup>

<sup>1</sup>California State University, Northridge, <sup>2</sup>NASA Marshall Space Flight Center, <sup>3</sup>The University of Alabama in Huntsville, <sup>4</sup>Universities Space Research Association


**Center for Space Plasma and Aeronomic Research (CSPAR)** 





## Abstract

To unveil the complexity of the solar atmosphere and transition region is fundamentally important, as this is where the forces transition from plasma to magnetic field dominated. Measurements of the field are also needed to elucidate the energy transport from the lower atmospheric regions to the first flight of the international solar sounding rocket mission, CLASP. For the first time, linear polarization measurements in this emission line, high sensitivity is required due to the relatively weak polarization signal compared to the intensity. To achieve this high sensitivity, a low-noise sensor is required with good knowledge of its characterization, including linearity. This work presents further refinement of the current from a photodiode in the light path to the digital response of the detectors. Pre-flight CCD linearity measurements were taken for all three flight cameras and calculations of the linear fits and residuals were performed. However, the previous calculations included a smearing pattern and a digital saturation region on the detectors which were not properly taken into account. The calculations have been adjusted and were repeated for manually chosen sub-regions on the detectors that were found not to be affected. We present a brief overview of the instrument, the calibration data and procedures, and a comparison of the successor mission, CLASP2, which will measure the Magnesium II h & k emission lines between 279.45 nm and 280.35 nm. The new approach will help to better prepare for and to improve the camera characterization for CLASP2.



3-dimensional solar atmosphere. Green lines

SN4 140620: Intensity per Pixel

charge from a saturated pixel can spill over to adjacent regions

(Hasinoff / Google Inc. & MIT) which is known as blooming, a

possible explanation for these smear regions. Though another

likely reason for smearing and saturation among these images

could be a result of not having the readout register properly

cleared prior to sampling images.

represent likely magnetic field lines.

Figure 2: CLASP-2 instrument design. Figure from Narukage et al., 2016

#### **1. Scientific Motivation**

To understand the solar atmosphere, measurement of the magnetic field in the upper chromosphere and transition region is fundamentally important, as this is where the forces transition from plasma to magnetic field dominated. In the photosphere, the magnetic field is moved by the plasma. There, gas force is greater than magnetic force. In the corona, plasma follows the magnetic field, where magnetic force is then greater than gas force. Measurement and interpretation of the strength and structure of the magnetic field in such regions is the intent of this project.

### 2. Introduction


The CLASP instrument measures the polarization in the Hydrogen Lyman-Alpha line, which comes from the chromosphere. The polarization in this ultraviolet line contains information about the chromospheric magnetic field. This polarization signal is very weak; to measure it, it is important to characterize the entire CLASP system, including the camera.


The first measurement of linear polarization in the Hydrogen Lyman-Alpha emission line at 121.60 nm occurred from the successful international solar sounding rocket mission, the Chromospheric Lyman-Alpha Spectro-Polarimeter (CLASP) launched in September of 2015.

## 4. Project Objectives and Description

The focal objective of this project involved characterizing the camera linearity by improving the linearity analysis of CLASP-1 to gain a greater understanding of the cameras for their second flight on on the CLASP-2 payload.

With miniscule intensity variation to begin with, maintaining low noise levels is required to prevent the polarization signal from drowning in noise. Hence for linear polarization measurements in the H Lyman-Alpha emission line, high sensitivity is required due to the relatively low strength polarization signal compared to the intensity. To achieve high sensitivity, a low-noise sensor is required with





## **3. Instrument Assembly**

The CLASP-1 payload was designed to survey ultraviolet spectra formed by hydrogen ions residing in the chromosphere, collectively forging the Hydrogen Lyman-Alpha UV emission line. It consists of a Cassegrain telescope, Slitjaw optics, and a Spectro-polarimeter. Within the Cassegrain telescope resides a continuously rotating half waveplate that allows for selection of the linear polarization direction. Also found within the Cassegrain telescope, the cold mirror permits for only a slim, narrow band to reflect the target wavelength, Hydrogen Lyman-Alpha, then rejects and transmits remaining visible light collected within the telescope.

light

mage

Array

light from sun



Grating

Order

First

Order

Oth,

Continuously Lyman-Alpha Change the Rotating

#### good knowledge of its characterization, including linearity.

We compared the current from a photodiode in the light path to the digital response of the detectors. Pre-flight CCD linearity measurements were taken for all three flight cameras and calculations of the linear fits and residuals were performed.

500

6x10<sup>4</sup>

5x10<sup>4</sup>

4x10<sup>4</sup>


3x10<sup>4</sup>

2x10<sup>4</sup>□

5.0x10<sup>4</sup>

#### Intensity (DN):

360



#### orders. This emulates Zeroth Order the diffraction in the Spectro-Polarimeter First Order (Figure 4). Incident Lyman-Alpha Second Order **Figure 5:** A CLASP Frame-Transfer CCD showing the two distinct regions: image and storage array. The dark area **5. Design and Operation of Science Cameras** is the image region, where light is gathered. The opaque light shield on the storage array blocks photons from reaching the rest of the sensor. The science cameras used contain frame-transfer ccds. The composition of the ccds themselves persist of dual parallel register clocks, two individual components, including an image array and storage array. The image array is a light-sensitive photodiode that collects photons casted onto the ccd. Essentially, the image focusing is done in the image array, where it then gets temporarily held prior to

**4**×10<sup>4</sup>

3x10<sup>4</sup>

**5** 2x10<sup>4</sup>


readout in the storage array. The parallel register clocks allow for charge to shift independently among the two arrays. This operation efficiently enables for continuous frame transfer at rapid succession and in the absence of a shutter. Frame transfer is performed by shifting rows individually in parallel to reach the serial shift register where it is then shifted through output as quantifiable data.

Figure 3: A genera


diffraction, casting its

1st, and 2nd


**SN4** Camera (Left Tap)



CCD Camera



Figures 9-11: Camera No. - Voltage (Brightness). Saturated frame for each CLASP camera. These images show the saturation and smearing pattern in strongly illuminated frames.



We tried a variety of polynomial functions to fit the linearity relationship. We found that as the degree of the polynomial is increased, the maximum percentage residual decreases. In reference to Figure 11, the 1st degree, the max residual of 0.31% completely differs in comparison to the 5th degree, 0.02%. The results show a smooth residual, perhaps not quite a polynomial, but is close enough.

8. References & Acknowledgements Hasinoff / Google Inc., S. W., & MIT. (n.d.). Saturation (imaging). Ishikawa, S. (NAOJ) *The Chromospheric* Lyman-Alpha Spectro-Polarimeter [PDF

Narukage, N. et al, "Chromospheric LAyer SpectroPolarimeter (CLASP2)", Proc. SPIE Space Telescopes and Instrumentation 2016: Ultraviolet to Gamma 990508 (July 2016); 11, doi:10.1117/12.2232245.

The material presented is supported by the National Science Foundation funded Research Experience for Undergraduates program "Solar and Heliospheric Physics at UAH and MSFC" run by the University of Alabama in Huntsville in partnership with NASA - Marshall Space Flight Center through Grant # AGS-1460767

Special thanks to Laurel Rachmeler, Amy Winebarger, Patrick Champey, Christian Bethge, and David Falconer.