


Improved hydrology over peatlands in a global land modeling system

M. Bechtold¹, G. De Lannoy¹, R. Reichle², R. Koster²,
S. Mahanama², D. Roose¹

¹ KU Leuven, Belgium, ² NASA-GMAO, USA



Background: Peatlands Under Pressure



Global Peatland (Histosol) Distribution; Xu et al., 2018

Long list of ‘peatland ecohydrological models’

- Focus: Carbon Cycle (with or without hydrological simulation)
- Water Level simulation challenging

Peat accumulation model	(Hilbert et al., 2000)
PCARS	(Frolking et al., 2002)
McGill Wetland	(St-Hilaire et al., 2010)
Biome-BGC	(Bond-Lamberty et al., 2007)
Wetland-DNDC	(Zhang et al., 2002)
Ecosys	(Dimitrov et al., 2011)
InTEC	(Ju et al., 2006)
BEPS	[Chen et al., 2007, 2005]
DigiBog	(Baird et al. 2012, Morris et al. 2012)
PEATBOG	(Wu et al., 2013)
... and several more	

Integration into continental/global land surface schemes


CLM	(Shi et al., 2015)
CLASS-CTEM	(Wu et al. 2016)
ORCHIDEE-PEAT	(Qui et al., 2017)

+ PEAT-CLSM


CLSM: Catchment Land Surface Model of NASA’s Goddard Earth Observing System Model (GEOS-5)

Motivation: Why CLSM of GEOS-5 ?

- 1) Coupled Ocean-Atmosphere-Land Model: Changed energy balance over peatlands affects atmospheric simulations
- 2) Land Data Assimilation System (→ e.g. SMAP L4 Soil Moisture Product)




Peatlands: Potential to monitor wetness variation with passive and active microwave observations (Kim et al. 2017, Bechtold et al. 2018)



CLSM: Main Characteristics

- High emphasis on efficiency (global appl.) (Koster et al. 2000)
 - Partitioning of land surface into hydrologic catchments
 - Topographic Wetness Index based model
→ subgrid soil moisture + water level variability and runoff
 - Each grid cell modeled with dominant catchment and soil
 - No numerical coupling between grid cell
-
- Peat as soil class (De Lannoy et al. 2014, JAMES)
→ Water levels however mostly still far too deep (~ 2 meter) and dynamics not typical for peatlands



Objective

- Implement typical peatland hydrological characteristics into CLSM
- Maintain simplicity and efficiency of CLSM

Scope narrowed to

- Northern Peatlands
- Degree of groundwater influence highly variable and unknown at global scale → All peatlands treated as rain-fed peatlands

Next:


- Model Modifications
- Validation
- Summary and Outlook

Model Modification #1



Topographic Wetness Index
Distribution from Catchment
Topography


Elevation Distribution
from typical Peatland
Micro-topography





Weston et al. 2015
Example of “hummock and
hollow microtopography”

Model Modification #1: dynamic surface water storage



Mineral land surface (here: no microrelief)



Peatland surface (microrelief)



Model Modification #2: Runoff



Continuous transition
from baseflow to
overland flow

Romanov, 1968,
Ivanov 1975


$$\text{Conductivity: } K_s(z) = \frac{K_{s,z=0}}{(1-z)^m}$$

$$\text{Transmissivity: } T_a(WTD) = \int_{z_{ac}}^{WTD} K_s(z) dz$$

$$\text{Runoff: } r(WTD) = v T_a$$


Model Modification #3: Evapotranspiration

- Evapotranspiration: Water stress coupled to water table depth
- Vegetation classes and evapotranspiration calculation as in CLSM



Validation (water table depth data)


70 monitoring wells
18 peatlands



Bogs


Fens

Temporal statistics




Validation (water table depth data)

Example 1: Bog in NW Germany
Mild winter, high precipitation, $R=0.9$



Example 2: Bog in Belarus
Long freezing period, $R=0.6$



• in situ
— CLSM1
— CLSM2
— CLSM3
— PCLSM


Here:
bias + std
corrected

- Water levels level off smoothly close to surface
- Capability to predict summer anomalies
- Capability to predict snow melt peaks


Validation: Inundation Extent

GIEMS: Global Inundation Extent from Multi-Satellites

1993-2007: monthly, 28km resolution
No calibration/validation over peatlands



Prigent et al., 2007



Summary

- Peatlands have a specific hydrological dynamics
- Simple solutions for global land surface models with significant effects

Outlook

- Validation: Evapotranspiration (Eddy Towers)
- Validation: Inundation (masking non-peatland areas, GIEMS 2.0)
- Data Assimilation using SMOS/SMAP Brightness Temperatures

Acknowledgments


Aurela, M., Braumann, F., Burdun, I., Devito, K., Drösler, M., Flanagan, L.B., Grygoruk, M., Kurbatova, J., Lohila, A., Mäck, U., Mauersberger, R., Munir, T., Price, J., Röhl, M., Sagris, V., Thiele, A., Tiemeyer, B., van der Kamp, G., Zaufit, M., Prigent C.



Validation


- Simulation experiments using different versions of the GEOS-5 Catchment Land Surface Model
- Domain: Northern Hemisphere
- Forcing data: MERRA-2 (corrected precip.)
- No parameter calibration for new model (PCM)
- Comparison with ~ 60 observed multi-year time series (13 clusters) of water table depth (WTD)

Experiment	M2	P	PC	F + Catchments + New Model Structure
Description	Operational Merra-2, only mineral soils	Revised soil input including Peat class (De Lannoy et al. 2015)	Peat class + Refined Topography and Catchment delineation	
Resolution	2/3° x 1/2°	EASEv2 M09	5' x 5'	5' x 5'




Radiative transfer parameters

- Brightness Temperature (passive microwave)



Dielectric constant of soil-water-air mixture
= f (sand, clay, poros, wp, soil moisture...)



Radiative transfer model

