

Soil Moisture Active/Passive (SMAP) L-band Microwave Radiometer Post-Launch Calibration Upgrade

Jinzheng Peng, Jeffrey R. Piepmeier, Emmanuel P. Dinnat, David M. Le Vine,
Rajat Bindlish, Giovanni De Amici, Priscilla N. Mohammed, Liang Hong, Derek Hudson

NASA Goddard Space Flight Center

Sidharth Misra, Simon H. Yueh, Kacie E. Shelton,
Adam P. Freedman, R. Scott Dunbar, Steven K. Chan

NASA Jet Propulsion Laboratory

Thomas Meissner
Remote Sensing Systems

Thomas Jackson
USDA ARS Hydrology and Remote Sensing Lab



Outline


- Current Status
 - Performance
 - Existing problems
- Calibration algorithm Upgrades for Next Release
 - Algorithm
 - Performance
 - Validation
- Conclusion



Current Performance

- Radiometer status is stable
 - L1B_TB Version 3 released in April 2016
 - Satisfy science requirements with margin
- NEDT is stable
 - Daily average include all qualified data



Metric	Allocation	Measured
Ocean Model RMSD (incl. NEDT)	1.4 K	1.2 K
NEDT (land)	1.6 K	1.2 K
NEDT (ocean)	1.1 K	0.95 K
Monthly Drift	0.4 K	+0.1/-0.25







Existing Problems

- Reflector thermal model update
 - Reflector loss reset to baseline value
 - Drift during eclipse season
- Bias over CS
 - Change with SAR on/off





Calibration Algorithm Upgrade



Note: CS: 110° pitch
SCS: 180° pitch




Reflector Calibration



- Excess Emissivity for Reflector/Radome derived by 3 independent groups

	Pol	JPL	GSFC	RSS	Baseline
Reflector	H	1.014	1.016	1.01	1.0026
	V	1.012	1.012	1.01	1.0023
Radome	H	NA	1.0023	NA	1.0003
	V	NA	1.0017	NA	1.0003

- Verified by a half-orbit SCS (180° pitch angle) during eclipse 2017





Tnd, Tref Offset & APC Calibration



- Joint-calibration in general
- Divided into 3 segments
 - Segment 1: Jul 07, 2015 ~
 - Global ocean, CS (110° pitch), SCS (180 ° pitch) during non-eclipse season
 - Constant Tnd & Tref offset
 - Constant APC for all phases
 - Segment 2: 03/31~04/03, 04/14~07/07, 2015
 - Global ocean, CS (110° pitch)
 - Dynamic Tnd & Tref offset
 - Segment 3: Apr 03~13, 2015
 - Global ocean, Amazon
 - Same Amazon emissivity in April 2015
 - Constant Tnd & Tref offset



Calibration Model

- Model

$$C_r \delta T_{ND} + \delta T_{ref} = \frac{1}{L_{RF}} (-\delta T_A + S \delta G)$$

where $C_r = \frac{C_{ant} - C_{ref}}{C_{nd}}$


L_{RF} : loss of the radiometer RF front-end

S : δT_A sensitivity to antenna gain uncertainty

δG : antenna gain uncertainty

δT_A : = measured TA - modeled TA

- Data for Calibration
 - Non-eclipse
 - Global ocean, CS, and SCS






Expected TA Adjustment



- TB changes vs Ver 3 without adjustment
 - Land TB increment: 2.90 K (V), 5.01 K (H)
 - Amazon TB difference (V-H): -0.68 K
- Bias need to be added to expected TAh for external cal targets
 - Three ocean roughness models compared
 - GMF
 - Aquarius
 - 2-scale roughness model
- Nadir looking maneuver performed
 - Bias for H-pol determined






Nadir-Looking Result





- Biases for TAexp_h
 - -0.25 K for CS
 - -0.65 K for GO
- Validation
 - Apply these biases to generate new calibration coefficients for V4
 - Apply new calibration coefficients to TA measurement & simulation
 - δTAs for nadir looking: 0.00 K over ocean and land, respectively.





Calibration Coefficients

- For data release V4





	δG
V-pol	-0.23%
H-pol	-0.46%



Calibration Drift over Global Ocean



- RMSD over global ocean: 0.06 K (V) 0.10 K (H)
- RMSD over CS: < 0.10 K





TA/TB Difference over Amazon



- TAv-TAh: 2.1 K
- TBv-TBh: 1.9 K







Land TB Increment (V4 – V3)



- Tested using T15160 (Version 4, Sep 2015 ~ Aug 2016)
- Compared to R13080 (Version 3)

	Estimated*	Actual	Actual $ lat < 45$
V	2.90 K	2.87 K	3.43 K
H	3.20 K	3.32 K	3.96 K

* Based on using CS, SCS and GO. Additional CS and GO adjustment for H-pol.





Conclusion

- SMAP radiometer are calibrated using upgraded calibration algorithm
 - Biases are added to modeled TA_h for global ocean and CS based on nadir-looking analysis
- Problems in V3 have been solved
 - Cal coefficients have been tested
 - Cal coefficients have been validated by using nadir-looking data.
- Land TB increment (V4 – V3)
 - 2.87 K (V), 3.32 K (H)
- Amazon TA/TB difference (V4)
 - 2.1 K for TA
 - 1.9 K for TB