


# Consideration of Dynamical Balances

Ronald M. Errico


Goddard Earth Sciences Technology and Research Center  
Morgan State University  
Global Modeling and Assimilation Office  
GSFC NASA

A presentation about concepts rather than techniques



# The Need for Balance: An Early Example


A time series of forecast surface pressure at a point



Solid: balance not considered  
Dashed : expected behavior


# Geostrophic Adjustment

0 hour




CONTOUR INTERVAL: 10 M

1 hour




CONTOUR INTERVAL: 4 M → REPRESENTS 0.5 M/S

2 hours




CONTOUR INTERVAL: 4 M → REPRESENTS 0.5 M/S

3 hours



CONTOUR INTERVAL: 4 M → REPRESENTS 0.5 M/S

6 hours



CONTOUR INTERVAL: 4 M → REPRESENTS 0.5 M/S

Figure 6.3 (a) Geostrophic adjustment of initial geopotential perturbation. (b–e) Solutions at 1, 2, 3, and 6 hours. Contoured field is geopotential, and wind arrows indicate speed and direction of windfield. (After Barwell and Bromley, 1988)

## Why does balance matter in data assimilation?

1. Unrealistic initial imbalances will create unrealistic forecasts.
2. Unrealistic imbalances can be accentuated through moist diabatic processes.
3. Large initial imbalances will tend to create less accurate backgrounds.
4. Balance can be exploited to relate  $u$ ,  $v$ ,  $T$ ,  $p_s$  (esp. in extra-tropics).
5. Errors in balanced initial conditions will tend to create balanced background errors, so the error statistics should reflect that; i.e., background errors of  $u$ ,  $v$ ,  $T$ ,  $p_s$  tend to be correlated, esp. in extra-tropics.

## Why is the extra-tropical atmosphere quasi-balanced?

Charney: 1955 *Tellus* (a paraphrase)

The observed extra-tropical, tropospheric motions are dominantly quasi-balanced because:

1. The principal atmospheric forcing is large scale and long period.
2. The quasi-balanced motion must be relatively stable with respect to gravity-wave perturbations. (*by inference; also see Errico 1981*)
3. Dissipation must be sufficient to remove what energy is otherwise leaked into gravity waves. (*added by R. Errico*)

Lorenz 1980 *JAS* Atmospheric dynamics lies on a slow-manifold.

Imposing balance is not just about getting rid of high frequency oscillations

# Filtered Equations

1st-order balance

e.g., quasi-geostrophic equations

2<sup>nd</sup>-order balance

e.g., nonlinear balance equation and  
quasi-geostrophic omega equation

These are too restrictive except for simple examples like the SWE.

# Dynamical Initialization




FIGURE 1.—Schematic representation of iteration methods for initialization with the primitive forecast equations.

Nitta and Hovermale 1969 *MWR*

The filters were not sufficiently selective.

## Normal Modes of the Linearized PE

Linearize P. E. about  $u = 0, v = 0, T = T_r, p_s = p_{sr}, z_s = 0$ :


$$\begin{aligned}\frac{\partial u}{\partial t} &= fv - \frac{\partial \phi}{\partial x} \\ \frac{\partial v}{\partial t} &= -fu - \frac{\partial \phi}{\partial y} \\ \frac{\partial \phi}{\partial t} &= -\tau_p \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right)\end{aligned}$$

Solutions:

$$\begin{pmatrix} u \\ v \\ \phi \end{pmatrix} = \sum_{m,n,k,j} c_{m,n,k,j} \Phi_k(p) \exp(imx + i\lambda_{m,n,k,j}t) \begin{pmatrix} \tilde{u} \\ \tilde{v} \\ \tilde{\phi} \end{pmatrix}_{m,n,k,j}(y)$$




Vertical structures of some normal modes  
(why noise may be most apparent in the stratosphere)




## Example structures of 2 normal modes

A fast gravity -wave like mode




A slow Rossby-wave like mode




# Normal-Mode Initialization

Solid Lines: balance not considered; Dashed Lines : balanced imposed at  $t=0$

*Linear:  $c(t=0) = 0$*



*Nonlinear:  $dc/dt (t=0) = 0$*



Machenhauer 1997 Cont. Atmos. Phys.; Baer and Tribbia 1997 MWR  
Daley 1991; Williamson and Temperton 1981 MWR

## Gravitational modes considered as forced and damped harmonic oscillators

Define  $g(t)$  as the complex amplitude of a gravity-wave like mode at each time  $t$ , and let  $R$  and  $G$  be the sets of Rossby- and gravity-wave like modes.

Then

$$\frac{dg}{dt} = -i\lambda g + N(R) + N(R, G) + N(G) + D(R, G) - \nu g$$

Consider  $N(R) = F(t)$  as the dominant nonlinear term. Approximately then

$$\frac{dg}{dt} = -i\lambda g + F(t) - \nu g$$

Consider  $F(t) = F(0) \exp(-i\mu t)$ . Then

$$g(t) = \left[ g(0) - \frac{F(0)}{i\lambda - i\mu + \nu} \right] \exp(-(i\lambda + \nu)t) + \frac{F}{i\lambda - i\mu + \nu}$$

## QG Theory and NNMI

In the extra-tropics, the NNMI balance condition  $dg/dt = 0$  is equivalent to

- (1) The nonlinear balance equation relating mass and vorticity fields, with some additional small terms;
- (2) The QG-omega equation defining the wind divergence, with some additional small terms;
- (3) Solved with the constraint that a form of linearized potential vorticity is specified;
- (4) And applied only to large vertical but small horizontal scales for which the resonant frequency is large.

The choice of constraint and scale selectivity matter!!

## Drawbacks of Initialization Techniques

“Initialization” was originally applied to an already produced analysis.

It would generally draw the state away from the analysis (and observations), rendering it balanced but no longer “optimal”.

NMMI was applied to some slow as well as fast gravitational modes in some DAS thereby destroying parts of the general circulation

When NNMI was considered a constraint in a variational DAS, the required repeated transforms between fields of  $u, v, T, p_s$  and the mode coefficients were very expensive.

## Balance imposed as a strong or weak constraint

### Strong Constraint

$$\begin{aligned} \text{minimize } J(\mathbf{x}) &= J_B[\mathbf{x} - \mathbf{x}_b] + J_O[\mathbf{y} - \mathbf{H}(\mathbf{x})] \\ \text{given } \mathbf{C}(\mathbf{x}) &= \mathbf{0} \end{aligned}$$



### Weak Constraint

$$\text{minimize } J(\mathbf{x}) = J_B[\mathbf{x} - \mathbf{x}_b] + J_O[\mathbf{y} - \mathbf{H}(\mathbf{x})] + J_C[\mathbf{C}(\mathbf{x})]$$

Daley 1978, Kleist et al. 2009 MWR, Courtier and Talagrand 1990, Tribbia 1982, Fillion and Temperton 1989

# Digital Filter Initialization

$$x_0^I = \sum_{k=-N}^N h_k x_k^u$$



Lynch and Huang 1992 MWR  
Fillion et al 1995 Tellus




# IAU (Incremental Analysis Update)

For  $n = -N, \dots, N - 1$

Bloom et al. 1996 MWR


$$f(t_0 + (n + 1)\Delta t) = f(t_0 + n\Delta t) + \Delta f_M(t_0 + n\Delta t) + \frac{1}{2N} \Delta f_A(t_0)$$




# Localization in Ensemble DA

Example response to a single observation of  $\phi$

Balanced  
Bkg Error  
Covariances



Same, with  
Schur product  
Localization



## Lessons Learned

1. There are many ways to balance models, each with varying degrees of success.
2. Most balance schemes have some undesirable consequences.
3. Balance should not be applied everywhere, at all scales, in the same way, to the same degree.
4. Balance should be considered when performing an analysis.
5. Details matter.

# Non-Dynamical Balances

There are other kinds of balances to consider:

Constraints on relative humidity

Cloud forecasts

Precipitation forecasts

See talk next week by T. Auligne  
Also, refs in Errico et al. 2008 JAS

## Common Misconceptions About Balance

F: Small scales are not balanced.

T: Balance depends on both vertical and horizontal scales.

T: Deep modes are likely balanced even on the mesoscale.

F: Atmospheric fields are on a “slow manifold.”

T: Some atmospheric forcing has short time scales.

T: In realistic models, freely propagating gravity waves are present to some degree.

## Common Misconceptions about Initialization

F: Initialization is inappropriate when gravity waves are important.

T: It is necessary when gravity waves may affect forecasts.

T: It removes waves which are not really there.

T: It is unnecessary when gravity waves are unimportant.

Techniques may come and go,  
but fundamentals remain (almost) forever.

(Unless, of course, they are neglected.)

# Recommended References

**References cited in the lecture or in the papers listed below.**

Concepts about balancing in general and nonlinear normal mode initialization in particular:

Daley, R., 1991: *Atmospheric Data Analysis*, Cambridge University Press. 420pp.

Presentation of some basic NNMI concepts using a simple, periodic f-plane model.

Errico, R. M., 1989: *Theory and application of nonlinear normal mode initialization*. NCAR Technical Note, NCAR/TN--344+IA, 145 pp.

Short presentation of some fundamental issues regarding balance and initialization:

Errico, R.M., 1997: On the removal of gravitational noise in numerical forecasts. *J. Meteor. Soc. Japan*, **75**, 219-227.

Revelation of some peculiar effects of nudging methods for assimilation:

Bao, J.W. and R. M. Errico, 1997: An adjoint analysis of the nudging method for data assimilation. *Mon. Wea. Rev.*, **125**, 1355-1373.

Incorporation of balance within (rather than after) the assimilation problem, including for the mesoscale:

Papers by Luc Fillion, Andrew Lorenc



ADDITIONAL SLIDES

## Consistency between analysis and initialization






FIG. 2. The 500-mb height field on 14 March 1990 (a) as analyzed by NOGAPS, (b) analysis increments, (c) initialization increments, and (d) the sum of analysis and initialization increments. Contour interval is 60 m in (a) and 10 m in (b)–(d). Zero contours are omitted; negative contours are dashed; and labels in (a) are dekameters.

# Global mean squared divergence tendency




Structures of 3 largest scale vertical normal modes

GMAO-GSI 3DVAR




# Harmonic Dial for External $m=4$ Mode, Period=3.7h

Without NNMI



With NNMI





Errico 1997 J Japan Met Soc

# Harmonic Dials from a Climate Simulation

External Mode  $P=3.7h$

Internal Mode  $P=11.6h$



$$\frac{dc_j}{dt} = -i\omega_j c_j + A(r,r) + B(r,g) + C(g,g) + D$$

Balance of  
Modes in a  
Climate Model

(a sophisticated  
scale analysis)

Normalized  
Sizes of Terms

