

Global Assimilation of Loon Stratospheric Balloon Observations

Lawrence Coy¹ Mark Schoeberl² Steven Pawson¹ Salvatore Candido³ and Robert W. Carver³

¹NASA Goddard Space Flight Center, Greenbelt, MD, USA; ²Science and Technology Corp, Columbia, MD, USA; ³Loon, Mountain View, CA, USA

The Problem

Verification of global data assimilation system (DAS) output fields by independent observations insures the quality of these products for scientific studies and can highlight the need for additional observations to improve the quality. This is especially important for middle atmosphere winds where there are few "in situ" observations available to constrain the DAS. Here we use constant pressure balloon derived winds to evaluate the lower stratospheric winds in the NASA GEOS DAS.

Loon Balloons



Goal: launch and maintain a fleet of balloons to provide Internet coverage to users on the ground.
Status: over 25 million km of test flights since the project began.
Flight Duration: up to 190 days in the stratosphere.
Winds: Derived from Loon Balloon GPS determined locations.

<https://loon.co/>


Methodology

NASA GMAO MERRA-2 Data Assimilation System

Two Experiments:


- Control Experiment** (assimilation of standard observation set)
- Loon Experiment** (assimilation of Loon balloon derived winds + standard observation set)

Test Time Period: June-August 2014 (large number of Southern Hemisphere Loon Balloons)



Loon Balloon Distribution (June-August 2014)


Number of Loon Observations (Jun-Aug 2014) per 1°x 250 m grid box




Southern Hemisphere
Loon Balloons 70-50 hPa
(18-21 Km) Altitude
60°S to 10°N Latitude

Balloon Altitudes are adjusted while in flight

Examples of Loon Balloon Coverage



6-Hour Assimilation System Data Window
Loon balloons can fill in wind observations over the oceans.



14 Days Tropical Coverage


Tropical observations provided some of the largest DAS adjustments when assimilating the Loon winds.

This Loon trajectory will be examined in more detail.

Forecast Improvement (o-f) RMS Summary

Assimilated Loon Balloon Winds greatly improves the tropical background forecast


Note: o-f denotes the difference between the observation and the short background forecast



Analysis Improvement (o-a) RMS Example

Analysis winds interpolated by the DAS in space and time to the Loon position

Note: o-a denotes the difference between the observation and the DAS analysis




Large differences occur between the Control Experiment Analysis and Loon Observations that are corrected in the Loon Experiment Analysis. (Black and gray are on top of each other.)


Analysis Differences (RMS) August 2014

Control and Loon Balloon differences found mainly at Loon Balloon Altitudes

Evidence of some influence extending into the troposphere, below the Loon Balloon altitude.




Additional Test Time Period (June-August 2016)



Sample Trajectories (~100-70 hPa Altitude Range) Western Hemisphere Concentration

Confirmation of 2014 Results

O-F RMS
Improvements at all latitudes, more significant improvement in tropics



Conclusions

- Additional wind observations can improve data assimilation products in the lower stratosphere, especially in the tropics
- Current tropical wind analyses in the lower stratosphere can have errors larger than 10 ms⁻¹
- Some of the stratospheric wind analysis improvement may be accompanied by improvements in the upper troposphere wind analysis (further investigation is planned)

Coy, L, M. R. Schoeberl, S. Pawson, S. Candido, & R. Carver, 2018, Global assimilation of X Project Loon stratospheric balloon observations, Submitted to *J. Geophys. Res.*



GMAO Publication List

E-mail: lawrence.coy@nasa.gov

Web: gmao.gsfc.nasa.gov

National Aeronautics and Space Administration

