


## **Increased Dependence on Saccades for Ocular Tracking with Low-dose Alcohol**

Terence Tyson, Nathan Feick, Patrick Cravalho, Tiffany Tran, Erin Flynn-Evans, Leland Stone

Previous studies have shown that certain features of oculomotor performance are impaired at or slightly below the legal limit for driving in most U.S. States (0.08% Blood Alcohol Concentration or BAC). Specifically, alcohol impairs saccadic velocity (Fransson et al., 2010; Roche & King, 2010), and steady-state tracking (Fransson et al., 2010; Moser et al., 1998; Roche & King, 2010) at levels between 0.04% and 0.1% BAC. Here we used a suite of standardized oculometric measures (Liston & Stone, 2014) to examine the effect of ultra-low levels of alcohol (down to 0.01% BAC) on steady-state tracking. Our high-uncertainty tracking task reveals that the smooth pursuit system is highly sensitive to BAC, with impairment extrapolating back to BAC levels at or below 0.01%. BAC generates a dose-dependent increase in reliance on the saccadic system that fully maintains overall steady-state tracking effectiveness at least up to 0.08% BAC, albeit with a significant decrease in smoothness.





# Increased Dependence on Saccades for Ocular Tracking with Low-Dose Alcohol

SJSU SAN JOSÉ STATE UNIVERSITY

Terence L. Tyson<sup>1</sup>, Nathan H. Feick<sup>1,2</sup>, Patrick F. Cravalho<sup>1,2</sup>, Tiffany Tran<sup>1,2</sup>, Erin E. Flynn-Evans<sup>1</sup>, and Leland S. Stone<sup>1</sup>

<sup>1</sup>Human Systems Integration Division, NASA Ames Research Center, <sup>2</sup>San José State University Research Foundation

Email: [terence.l.tyson@nasa.gov](mailto:terence.l.tyson@nasa.gov)


## Introduction

Previous studies have shown that certain features of oculomotor performance are impaired at or slightly below the legal limit for driving in most U.S. States (i.e., 0.08% BAC). Specifically, alcohol impairs saccadic velocity (Fransson et al., 2010; Roche & King, 2010), and steady-state tracking (Fransson et al., 2010; Moser et al., 1998; Roche & King, 2010) at levels between 0.04% and 0.1% BAC. Here we used a suite of standardized oculometric measures (Liston & Stone, 2014) to examine the effect of ultra-low levels of alcohol (down to 0.003% BAC) on steady-state tracking.

## Methods


- 16 healthy participants (8 females, mean age  $\pm$  SD = 25.6  $\pm$  3.1 years) with normal or corrected-to-normal visual acuity.
- 2-day study where, on a given day, subjects consumed one of two possible doses of ethanol (40% ABV Vodka mixed with juice; targeting either 0.06% or 0.02% peak BAC), with 3 pre-dose and 6-9 post-dose ocular tracking tests using a 5 minute Rashbass-like task (Krukowski & Stone, 2005).

Speed & Directional Uncertainty Target Motion Onset Uncertainty




- We then computed 6 largely independent measures of pursuit and saccadic performance in a steady-state analysis interval, 400-700 ms after target motion onset (Liston & Stone, 2014).

## Example Trial




## Pursuit Behavior



Blood Alcohol Concentration (%)

- Steady-state gain and proportion smooth was compromised above 0.01% BAC (linear regression: both  $p < 0.0001$ , and x-intercepts: 0.003% BAC and 0.002% BAC).
- Gain loss causes  $\sim 1.1$  deg of "lost ground" in steady-state for the 0.065% BAC bin.

## Saccade Behavior



Blood Alcohol Concentration (%)

- Saccades became larger and more frequent (linear regression:  $p < 0.0001$  and  $p < 0.003$ ).
- Saccades gained back, on average, 0.9 deg ( $\sim 80\%$ ) of the lost ground.
- Main sequence was disrupted with peak velocity increasingly less sensitive to changes in amplitude with increasing BAC (linear regression:  $p < 0.0002$  and  $p < 0.0001$ ).

## Conclusions


- A high-uncertainty tracking task revealed that the smooth pursuit system is highly sensitive to BAC, with impairment extrapolating back to BAC levels at or below 0.01%.
- This cannot be explained by subjective, non-BAC effects of alcohol consumption.
- The results suggest that ethanol consumption generates a dose-dependent increase in reliance on the saccadic system, which compensates for up to 80% of the lost ground at  $\sim 0.065\%$  BAC, albeit with a significant decrease in smoothness.

## References

1. Fransson, P. A., Modig, F., Patel, M., Gomez, S., & Magnusson, M. (2010). Oculomotor deficits caused by 0.06% and 0.10% blood alcohol concentrations and relationship to subjective perception of drunkenness. *Clin Neurophysiol*, 121(12), 2134-2142. doi:10.1016/j.clinph.2010.05.003
2. Krukowski, A. E., & Stone, L. S. (2005). Expansion of direction space around the cardinal axes revealed by smooth pursuit eye movements. *Neuron*, 45(2), 315-323. doi:10.1016/j.neuron.2005.01.005
3. Liston, D. B., & Stone, L. S. (2014). Oculometric assessment of dynamic visual processing. *J Vis*, 14(14), 12. doi:10.1167/14.14.12
4. Moser, A., Heide, W., & Kömpf, D. (1998). The effect of oral ethanol consumption on eye movements in healthy volunteers. *J Neurol*, 245(8), 542-550.
5. Roche, D. J., & King, A. C. (2010). Alcohol impairment of saccadic and smooth pursuit eye movements: impact of risk factors for alcohol dependence. *Psychopharmacology (Berl)*, 212(1), 33-44. doi:10.1007/s00213-010-1906-8

Supported by the Force Health Protection Program of the Office of Naval Research (SAA 402925)

## Controls



Blood Alcohol Concentration (%)

There was a significant non-BAC effect of initial dose size on subjective drunkenness (different slopes,  $p < 0.03$ ), but non-BAC effects on slope and intercept did not reach significance for any of the 6 objective performance measures ( $p > 0.17$ ). Data from peak velocity metrics not shown.