



# On the Usage of Recalibrated Radiance in Reanalysis Experiments

Mohar Chattopadhyay<sup>1,2</sup>



Will McCarty<sup>1</sup> and Isaac Moradi<sup>1,3</sup>

<sup>1</sup>Global Modelling and Assimilation Office, NASA GSFC

<sup>2</sup>Science System and Application Inc.

<sup>3</sup>ESSIC, University of Maryland

## Motivation



- Sudden changes in background departure of Channel 14 in MERRA-2
  - Introduction of NOAA-17 AMSU-A
  - Introduction of AQUA AMSU-A
  - Data gap of NOAA16 AMSU-A on 19-28 November 2002
- Loss of NOAA-16 AMSU-A reflected in upper stratosphere temperature
  - Mean T shown at 2hPa and between 20-50° S



## Motivation

This effort aims to explore the impact of alternate data-source/inter-calibrated radiance on a similar system by:

- Conducting experiments using inter-calibrated radiance to test the sensitivity of the system
- Study if these new data sets can successfully alleviate sensitivity of temperature in upper stratosphere.



## Data/Experiment Design

Ferraro et al. (2016) :

- channels 1, 2, 3 and 15 which have 23, 31, 50, 89 GHz frequency
- Corrections: Viewing angle, location dependent bias and many more

Zou and Wang (2013):

- Simultaneous Nadir Overpass (SNO) CH6 from NOAA-18 and CH(4-7) and CH(8-14) from NOAA-15 as reference
- Integrated Microwave Inter-Calibration Approach (IMICA) method to obtain a long-term data product
- Correction derived from the global ocean differences between Aqua and the reference satellite.

Data Source	Reference	Description
<b>M2</b>	N/A	Antenna Temperature bufr stream used in MERRA-2
<b>FERRARO</b>	Ferraro et al. 2016	Inter-calibrated AMSU-A measurements for Channels 1-3, 15
<b>Z&amp;W</b>	Zou and Wang 2013	Inter-calibrated AMSU-A measurements for Channels 3-14
<b>NOAA</b>	NOAA KLM User's Guide, 1999	Antenna Temperature acquired from Z&W based on NOAA CLASS



## Experiment Design

Assimilation (3DVar) and the model configurations are similar to MERRA-2 specification

Horizontal Resolution C90  
~100km (MERRA-2 C180  
~50km)

72 vertical levels to 0.1 hPa


Study period: 2 July –  
31 December 2002

Experiment	AMSU-A Data Source		
	CH: 1-3, 15	CH: 4-13	CH: 14
<b>CTL</b>	M2	M2	M2
<b>EXP_TANT</b>	NOAA	NOAA	NOAA
<b>EXP_IC</b>	FERRARO	Z&W	Z&W
<b>EXP_IC14</b>	NOAA	NOAA	Z&W
<b>MERRA-2</b>	M2	M2	M2

## Climatology : MERRA-2 and CTL


(a)

MERRA-2 Climatology




(b)

MERRA-2-CTL Climatology




- (a) Zonal mean temperature and zonal mean zonal wind for MERRA-2 from 2 July to 31 December 2002 is shown
- (b) Minimal differences between MERRA-2 and CTL
- This confirms that results from experiments at C90 can be translated to MERRA-2 at C180

# Observation statistics: AMSU-A Channel 13




- AMSU-A channel 13 is bias corrected
- CTL bias correction values vary between instruments
- Implies difference in AMSU-A radiance between instruments in CTL
- Similar bias correction values in MERRA-2
- Values of bias correction are more consistent across sensors in EXP\_TANT

# Observation Statistics : AMSU-A Channel 14



- AMSU-A Channel 14 is not bias corrected
- Shows discontinuities in analysis and background departure with the change in observing system in CTL
- EXP\_TINT has a smoother analysis and background departure response compared to CTL
- EXP\_IC and EXP\_IC14 show response similar to EXP\_TINT
- Issues with AMSU-A data used in MERRA-2

## Upper Troposphere Temperature: July to December 2002




Time-series: Zonal Mean Temp at  
2hPa/mean of 50° S and 20° S Lat

- A bias of average 4 K is noted between CTL and the other experiments
- CTL shows a drop in upper stratospheric temperature with absence of NOAA-16 in assimilation
- EXP\_TANT and EXP\_IC14 have very similar temperature profile than EXP\_IC due to corrections applied to CH14 in Z&W radiance

## Upper Stratosphere Temperature : August 2004

Mean Temperature at 2 hPa between 20- 50°S



- EXP\_TANT, EXP\_IC and EXP\_IC14 simulated in 2004 to test stability
- MLS temperature assimilation from 5hPa and above commenced on 13 August 2004
- Negligible stratosphere temperature difference between experiments after MLS temperature assimilation
- MLS temperature acts as an anchor in upper stratosphere



## Conclusion

Established that MERRA-2 data stream needs to be reprocessed for future reanalyses.

- The MERRA-2 AMSU-A data prior to 2005 was the same as MERRA

At minimum, the early AMSU-A record needs to be reprocessed to be consistent with the base NOAA antenna temperature data record

Using NOAA data-stream for AMSU-A will partially alleviate the upper stratosphere temperature sensitivity, but more considerations on the recalibrated radiances are underway

Upper stratosphere temperature achieves stability with the introduction of MLS temperature assimilation

- MLS temperatures anchor to the upper stratospheric temperature
- Prior to MLS, this is primarily done by assimilation CH14 without bias correction, but that data record was questionable in previous GMAO reanalyses
- MLS has no follow-on, and it has yet to be studied how much GNSSRO acts to anchor the radiances in the context of the upper stratosphere