


Assessing the Impact of Urbanization Using Remote Sensing On A Global Scale, Past Present And Future Directions




Jeffrey C. Luvali and Glynn Hulley
NASA Marshall Space Flight Center
JPL

Global Urbanization- A Sense of Scale

- The 21st century is the first “urban century”
- In 2000, approximately 3 billion people (40% of global population) resided in urban areas
- The United Nations estimates that by 2025, 60% of the world’s population will live in cities
- As a consequence, the number of “megacities” – those cities with populations of 10 million or more – will increase to 100 by 2025



A dark blue silhouette of a city skyline with various skyscrapers and buildings, positioned behind the text.

Surface Radiation Budget

$$Q^* = (K_{in} + K_{out}) + (L_{in} + L_{out})$$

Q^* = Net Radiation

K_{in} = Incoming Solar

K_{out} = Reflected Solar

L_{in} = Incoming Longwave

L_{out} = Emitted Longwave

A dark blue silhouette of a city skyline with various skyscrapers and buildings of different heights, set against a lighter blue background.

Surface Energy Budget

$$Q^* = H + LE + G$$

H = Sensible Heat Flux

LE = Latent Heat Flux

G = Storage (maybe + or -)

- 
- A bright sun is visible in the center of the slide, partially obscured by the text. The sky is a hazy orange color, and the sun itself is a bright yellow-white circle with a soft glow.
- **European heat wave caused 35,000 deaths 2003**
 - **Over 15,000 likely dead in Russian 2010 heat wave; Asian monsoon floods kill hundreds more**
 - **Heat wave death toll in NYC rises to 8 NYDN 7/23/13**
 - **UK Heat wave death toll: Up to 760 killed and total may double as temperatures above 30° c continue 7/18/13**
 - **Chicago July 1995 more than 700 died**

ATLANTA 25
→



Climate Impacts of Land-Cover and Land-Use Changes in Tropical Islands under Conditions of Global Climate Change

DANIEL E. COMARAZAMY

NOAA/CREST Center, City College of New York, New York, New York

JORGE E. GONZÁLEZ

NOAA/Cooperative Remote Sensing Science and Technology Center (CREST), and Department of Mechanical Engineering, City College of New York, New York, New York

JEFFREY C. LUVALL AND DOUGLAS L. RICKMAN

Global Hydrology and Climate Center, NASA Marshall Space Flight Center, Huntsville, Alabama

ROBERT D. BORNSTEIN

Department of Meteorology and Climate, San Jose State University, San Jose, California

(Manuscript received 7 February 2012, in final form 6 September 2012)

Quantification and mitigation of long-term impacts of urbanization and climate change in the tropical coastal city of San Juan, Puerto Rico

Daniel E. Comarazamy¹, Jorge E. González^{2*} and Jeffrey C. Luvall³

¹The NOAA-CREST Center, The City College of New York, New York, NY, USA

²The NOAA-CREST Center and Department of Mechanical Engineering, The City College of New York, New York, NY, USA; ³Global Hydrology and Climate Center, NASA Marshall Space Flight Center, Huntsville, AL, USA

JOURNAL OF GEOPHYSICAL RESEARCH
Atmospheres
AN AGU JOURNAL



Research Article

Combined impacts of land cover changes and large-scale forcing on Southern California summer daily maximum temperatures

Pedro Sequera , Jorge E. González, Kyle McDonald, Robert Bornstein, Daniel Comarazamy

First published: 21 September 2015 Full publication history

DOI: 10.1002/2015JD023536 View/save citation

Cited by: 0 articles Check for new citations



Funding Information







Urban Remote Sensing and Air Quality Models

**Volatile Organic Compounds
+ Nitrogen Oxides
+ Sunlight**

→ Ozone



- Air pollution remains a National issue.
- Temperature increases the ozone levels.
- Urban heat island has major effect on temperature and height of mixing layer.
- Measurement program is defining land use patterns and relationship to heat production.
- Remote sensing data are being used to improve air quality modeling.



NASA's Project Objectives



- ▲ To use high spatial resolution thermal infrared and visible data obtained from aircraft to measure, map, and model the surface energy budget characteristics of surfaces typical of the urban landscape for three US cities.
- ▲ Provide these data to EPA for evaluation of the overall "fabric" of the cities in relation to the urban heat island and air quality modeling.
- ▲ Transfer NASA technology and research to the public.

NASA's Project Atlanta
~ 1996 - 2001

EPA/NASA Urban Heat Island Pilot Project
~ 1997- 2000



NASA EPSCoR San Juan, Puerto Rico UHI
2004

Urban Heat Island Mitigation Strategies



- ▲ Albedo Modification
 - Lighter colored roofs and pavements
 - New materials/coatings
- ▲ Plant trees and increase green space
 - Shade buildings, rooftops, parking lots and roads
 - Cool the air through transpiration
- ▲ Rooftop gardens
 - Keep roofs cool by shading and/or transpiration
 - storm water reduction




Baton Rouge
Albedo - May 11, 1998



Baton Rouge
Temperature - May 11, 1998




Baton Rouge
Scatter Plots of Albedo vs Temperature




Baton Rouge Scatter Plots Albedo vs Temperature


Industrial
(refinery)




Bayou
(Forest)





CBD



Residential




Whole
Mosaic





Epidemiologic Triangle of Disease (Vector-borne Diseases)

A multi-factorial relationship between hosts, agents, vectors and




ECOsystem Spaceborne Thermal Radiometer Experiment on Space Station


Dr. Simon J. Hook, JPL, Principal Investigator

Science Objectives

- Identify **critical thresholds of water use and water stress** in key climate-sensitive biomes
- Detect the timing, location, and predictive factors leading to plant **water uptake decline** and/or cessation over the **diurnal cycle**
- Measure **agricultural water consumptive use** over the contiguous United States (CONUS) at spatiotemporal scales applicable to improve drought estimation accuracy



When stomata close, CO₂ uptake and evapotranspiration are halted and plants risk starvation, overheating and death.



Water stress is quantified by the Evaporative Stress Index, which relies on evapotranspiration measurements.

ECOSTRESS will provide critical insight into **plant-water dynamics** and how **ecosystems change with climate** via **high spatiotemporal** resolution thermal infrared radiometer measurements of evapotranspiration from the International Space Station (ISS).



HyspIRI Objectives and Approach



Key Science and Science Applications

Climate: Ecosystem biochemistry, condition & feedback; spectral albedo; carbon/dust on snow/ice; biomass burning; evapotranspiration

Ecosystems: *Global* biodiversity, plant functional types, physiological condition, and biochemistry including agricultural lands

Fires: Fuel status; fire frequency, severity, emissions, and patterns of recovery *globally*

Coral reef and coastal habitats: *Global* composition and status


Volcanoes: Eruptions, emissions, regional and *global* impact

Geology and resources: *Global* distributions of surface mineral resources and improved understanding of geology and related hazards

Applications: Disasters, EcoForecasting, Water, Health/AQ

Mission Urgency

The HyspIRI science and applications objectives are critical today and uniquely addressed by the combined imaging spectroscopy, thermal infrared measurements, and IPM direct broadcast.



Measurement




Imaging Spectrometer (VSWIR)

- 380 to 2500nm in ≤ 10 nm bands
- 60 m spatial sampling*
- 19 days revisit*
- Global land and shallow water

Thermal Infrared (TIR):

- 8 bands between 4-12 μ m
- 60 m spatial sampling
- 5 days revisit; day/night
- Global land and shallow water

IPM-Low Latency data subsets



Mission Concept Status

Level 1 Measurement Requirements: Vetted by community and stable

Payload: VSWIR Imaging Spectrometer, TIR Multi-spectral Radiometer, and Intelligent Payload Module (IPM)

Full Mission original option: Mature

Separate Small Mission option: Pegasus-based solutions identified and studied

***SLI Support:** HyspIRI VSWIR evolving to 30m at 185km swath

ECOSTRESS TIR: Selected EVI for ISS

VSWIR Dyson Option: Technology/Science ISS Demonstration

Summary: The HyspIRI mission measurement requirements and baseline instruments approach are mature and stable with good heritage, low risk and modest cost. Now exploring a range of instrument and data options to save cost, per guidance letter.

HyspIRI TQ4. Urbanization/Human Health

- **How does urbanization affect the local, regional and global environment? Can we characterize this effect to help mitigate its impact on human health and welfare?**
- **How do changes in land cover and land use affect surface energy balance and the sustainability and productivity of natural and human ecosystems?**
- **What are the dynamics, magnitude, and spatial form of the urban heat island effect (UHI), how does it change from city to city, what are its temporal, diurnal, and nocturnal characteristics, and what are the regional impacts of the UHI on biophysical, climatic, and environmental processes?**

- **Human Health - heat mortality, vector borne diseases**
- **Heat and Air Quality**
- **Urban Heat Island (UHI)**
- **Land Cover/Land Use change**
- **Regional climate impacts**

References

- Bornstein, R., and Q. Lin (2000), Urban heat islands and summertime convective thunderstorm in Atlanta: Three cases studies, *Atmos. Environ.*, 34, 507–516.
- D. Comarazamy, J.E. González, J. Luvall, D. Rickman, and R. Bornstein. 2013. Climate impacts of land cover and land use changes in tropical islands under conditions of global climate change. *J. of Climate*, doi.org/10.1175/JCLI-D-12-00087.1
- A. Dominguez, J. Kleissl, J. C. Luvall, D. L. Rickman. High-resolution urban thermal sharpener (HUTS), *Remote Sensing of Environment* 115 (2011), pp. 1772-1780.
- DOUIRA>2.0.CO;2. Tso, C. P. (1995), A survey of urban heat island studies in two tropical cities, *Atmos. Environ.*, 30, 507–519.
- González, J. E., J. C. Luvall, D. Rickman, D. E. Comarazamy, A. J. Picón, E. W. Harmsen, H. Parsiani, N. Ramírez, R. Vázquez, R. Williams, R. B. Waide, and C. A. Tepley, 2005: Urban heat islands developing in coastal tropical cities. *EOS Transactions, AGU*, 86, 42, pp. 397 & 403.
- Jauregui, E., and E. Romales (1996), Urban effects of convective precipitation in Mexico City, *Atmos. Environ.*, 30, 3383–3389.
- Lo, C. P., D.A. Quattrochi, and J. C. Luvall (1997), Applications of high-resolution thermal infrared remote sensing and GIS to assess the urban heat island effect, *Int. J. Remote Sens.*, 18(2), 287–204.
- Luvall, J. C., D. Rickman, D. Quattrochi, and M. Estes (2005), *Aircraft based remotely sensed albedo and surface temperatures for three US cities*, paper presented at Cool Roofing: Cutting Through the Glare Roofing Symposium, Roof Consult. Inst. Found., Atlanta, Ga., 12–13 May.
- Luvall, J.C., Quattrochi, D.A., Rickman, D.L., Estes Jr., M.G., 2015. Urban Heat Islands. In: Gerald R. North (editor-in-chief), John Pyle and Fuqing Zhang (editors). *Encyclopedia of Atmospheric Sciences*, 2nd edition, Vol 1, pp. 310–318. ISBN: 9780123822253.
- Shepherd, J. M., and S. J. Burian (2003), Detection of urban-induced rainfall anomalies in a major coastal city, *Earth Interact.*, 7(4), doi:10.1175/1087-3562(2003)007<0001:
- United Nations Population Fund (1999), *The state of world population 1999*, 76 pp., New York. (Available at <http://www.unfpa.org/swp/1999/index.htm>)