

An Ocean-atmosphere Simulation for Studying Air-sea Interactions

**Ehud STROBACH^{1,2}, Andrea MOLOD², Atanas TRAYANOV^{2,3}, William PUTMAN²,
Gael FORGET⁴, Jean-Michel CAMPIN⁴, Chris HILL⁴, Dimitris MENEMENLIS⁵,
Patrick HEIMBACH⁶**

¹University of Maryland, United States, ²NASA / GMAO, United States, ³Science Systems and Applications, Inc., ⁴Massachusetts Institute of Technology, United States, ⁵Jet Propulsion Laboratory, California Institute of Technology, United States, ⁶University of Texas at Austin, United States



Overall motivation of the research program

- Couple GEOS atmospheric model and MITgcm ocean model.
- Perform ocean analysis using the MITgcm 4D-var data assimilation capability.
- **Develop a prototype ocean-ice-atmosphere weakly coupled data assimilation system by exploiting and leveraging GEOS and MITgcm data assimilation capabilities.**

Applications

- Recent sea ice and ice sheet changes.
- Sub-seasonal to decadal climate predictions.
- **Mesoscale air-sea interactions.**
- Observation System Simulation Experiments (OSSEs).

Current state of project

- GEOS-MIT model is now running with overall realistic results.
- Issues:
 - Too much net heat flux to the ocean (cloud forcing).
 - “The double ITCZ problem”
 - Too much accumulation of sea-ice in some regions (e.g. the Beaufort Sea).
- Tuning is about to commence using Green’s function method (Menemenlis *et al.*, 2005).



Air sea interactions in the high resolution GEOS-MIT






Current objectives of this study

- Develop a high resolution coupled ocean-atmosphere run for studying air sea interactions and simulating an observation system.
- Investigate the ability of the coupled model to capture the strong observed positive correlations between SST and wind stress/speed.
- Compare near-surface diagnostics of the fully coupled ocean-atmosphere set-up to equivalent atmosphere-only simulations.


Background: observed SST/wind stress anomaly correlations

Agulhas Return Current

QuikSCAT and AMSR SST




ECMWF and RTG SST




Gulf Stream

QuikSCAT and AMSR SST



ECMWF and RTG SST



1 DegC contours

“Satellite observations have revealed a remarkably strong positive correlation between sea surface temperature (SST) and surface winds on oceanic mesoscales of 10–1000 km.”

Chelton *et al.*, *Oceanography* (2010)


... correlation between SST and surface wind stress, is realistically captured only when the ocean component is eddy resolving.”

Bryan *et al.*, *J. Clim.* (2010)

Two-month averages (January–February 2008) of **spatially high-pass-filtered sea surface temperature (SST)** overlaid as contours on **spatially high-pass-filtered wind stress**.

Methods - models

- Atmosphere – GEOS:
 - Horizontal grid type – Cubed sphere, $1/8^\circ \times 1/8^\circ$
 - Vertical grid type – hybrid sigma-pressure, 72 levels
- Ocean – MITgcm
 - Horizontal grid type – Lat-Lon-Cap, $1/12^\circ \times 1/12^\circ$
 - Vertical grid type – z^* rescaled height vertical coordinate, 90 levels



Cubed sphere grid (left) and Lat-Lon-Cap (right)

Methods - experimental setup

1) Atmosphere Only – GEOS (AGCM)

- Feb, 9 – Apr 9, 2012
- Forcing: SST and ice fraction from an equivalent ocean-only experiment
- Initial conditions: MERRA-2


2) Coupled – GEOS-MITgcm (AOGCM)

- Feb, 9 – Apr 9, 2012
- Ocean initial conditions: from an equivalent ocean-only experiment
- Atmospheric initial conditions: MERRA-2 (same as the run 1)






Ocean surface current




Lagged correlation between daily SST $\left(\frac{\Delta SST}{\Delta t}\right)$ and wind speed $\left(\frac{\Delta WS}{\Delta t}\right)$

GEOS-MITgcm




Correlation between daily SST $\left(\frac{\Delta SST}{\Delta t}\right)$ and wind speed $\left(\frac{\Delta WS}{\Delta t}\right)$

GEOS



Possible mechanism



Conclusions

- First analysis of the ~10km coupled GEOS-MITgcm model reproduces realistic synoptic and mesoscale patterns.
- The coupled model shows positive correlations between SST and wind speed/stress, and the relation is slightly closer to observational estimates compared to previous simulations.
- The fact that the atmosphere-only experiment can reproduce the positive correlation suggests that the atmosphere responds to the ocean.
- Daily time series suggest a three-four-day cycle induced by air-sea feedbacks.



Next steps/future work

- Model tuning using green's function method.
- Increasing horizontal resolution (~1km).
- Recent sea ice and ice sheet changes.
- Initialized sub-seasonal to decadal prediction system.
- Observation System Simulation Experiments (OSSE).



Computational Issues – Doubling the Resolution

1/16°X1/16° Atmosphere, 1/24°X1/24° ocean:

- Initialize/finalize - ~2 hours to initialize, ~1 hour finalize
- Node memory – using only 20 out of CPUs per 128GB node
- Pre/post processing (1 3D field ~21GB, ~0.5TB for restart file)
- Time stepping: ~1 time step per 15 sec
- SYNCIO/IOSERVER: parallel I/O