

An AI Based System for Objective Tropical Cyclone Intensity Estimation

Jeffrey Miller¹, Manil Maskey^{1,2}, Rahul Ramachandran^{1,2}, Iksha Gurung¹, Brian Freitag¹,
Drew Bollinger³, Ricardo Mestre³, Daniel da Silva³, Andrew Molthan², Christopher Hain², Dan Cecil²

1 – NASA IMPACT, 2 – NASA Marshall Space Flight Center, 3 – Development Seed



development **SEED**



Outline

- Motivation
- Introduction/Background
- Data/Methodology
- Model Evaluation
- Deploying model in production
- Challenges and Lessons Learned
- Conclusions/Future Work

Motivation

15 UTC 10 Oct 17 NHC advisory on Tropical Storm Ophelia:

*“Dvorak intensity estimates range from T2.3/**33 kt** from UW-CIMSS to T3.0/**45 kt** from TAFB to T4.0/**65 kt** from SAB. For now, the initial intensity will remain at **45 kt**, which is an average of the scatterometer winds and all of the other available intensity estimates.”*

1. Can we objectively estimate wind speed from satellite images?
2. Can we estimate more frequently?


Intensity Estimation Current Approach

- The Dvorak technique

- Vernon Dvorak (1970s)
- Satellite-based method
- Based on cloud presentation in IR images
- A T-number is assigned based on the visual pattern of clouds



- Problems with current approach:

- Subjective
- Lack of generalizability
- Inconsistency
- Requires domain expertise




Data

- GOES IR imagery
 - Atlantic and Eastern Pacific Basins
 - 103,600 total images
 - 2000 - 2017
 - 2017 used for testing (5410)
 - GOES08 – GOES15
 - 5 degrees around center of storm
- Labeled with HURDAT2 reanalysis data



Methodology


- Performance struggled when binning the wind speeds into discrete classes
- Used a custom network architecture that outputs an actual wind speed estimate rather than a class (linear model)
 - 4 convolutional layers
 - 4 dense layers
 - 1 output
- Model was built using TensorFlow



Model Evaluation

Test set (2017 storms):

- RMSE: 10.39 Kts
- MAE: 7.58 Kts



North Atlantic

Piñeros et al. (2011): 14.7kt


Ritchie et al. (2012): 12.9kt

North Pacific


Ritchie et al. (2014): 14.3kt

Performance on individual storms

Atlantic:




East Pacific:



"The cloud pattern now only supports an intensity of about 45 kt, but **Dvorak intensity estimates are constrained to higher values by the rules of the technique**. Given the small size of Otis, it seems possible that this is a rare case where **the intensity is dropping faster than the Dvorak technique allows.**"

- Tropical Storm Otis Discussion Number 28

Model Deployment



Coordinated Effort



ML Researchers

- Transform ideas into models
- Training data
- Monitor



Marshall Space
Flight Center

Domain experts

- Evaluation
- Performance baselines
- Science use case



End-user stakeholders


- Production Requirements




ML/System developers

- Design
- Quick prototype
- Deploy to production
- Scale
- Log

Portal Workflows





Challenges and Lessons Learned

- Acquiring consistent large scale training data
 - NRL/CLASS
 - HURDAT/Aircraft recon
- Interpreting why a model is performing poorly
 - AI black box
- Properly versioning training data, models, and algorithms can be difficult
- Complexity with evolving platforms and infrastructure

Conclusions/Future Work

- We developed a deep learning model to objectively estimate hurricane wind speed
 - 10.39 kts RMSE
 - Works in both basins
- Deployed model in production environment
 - <http://hurricane.dsig.net/>
 - Utilizes *Cumulus* to execute workflows
- Other satellite imagery deep learning projects
 - Event detection
 - Phenomena detection
 - Smoke detection

Questions?

jjm0022@uah.edu

<http://hurricane.dsig.net>



development **SEED**

