

AQUEOUS ALTERATION

Affiliation

Yoko Kebukawa

Faculty of Engineering, Yokohama National University

79-5 Tokiwadai, Hodogaya-ku, Yokohama 240-8501, Japan

Email: kebukawa@ynu.ac.jp

Michael E. Zolensky

ARES, NASA Johnson Space Center

2101 NASA Parkway, Houston, TX 77058, USA

Email: michael.e.zolensky@nasa.gov

Synonyms

[If possible; choose the appropriate heading; only true synonyms will be considered. Please enter your text here ...]

Chemical Formula (*optional*)

[Please enter your text here ...]

Acronyms (*optional*)

[Please enter your text here ...]

Definition

[A short dictionary type description (<100 words) Please enter your text here ...]

A secondary process whereby liquid water modified the nature of anhydrous primary nebular components. In general, this process induced decomposition and changed the structures and compositions of primary minerals and formed secondary minerals in their place. This process occurred in the early history of meteorite parent bodies that contained water ice. The most effective heat source is considered to be the decay of short-lived radioactive nuclide such as Aluminium-26, but other causes have been suggested. The degree of aqueous alteration of meteorites is categorized as type 1 (most altered) to type 3 (least altered), according to mineralogy and petrology.

Keywords

[Significant terms which give a short overview of the following text. Please enter your text here ...]

Meteorites, Asteroids, Water, Minerals, Organic matter

History (*optional*)

[Please enter your text here ...]

Overview

[Extended discussion <600 words. Please enter your text here ...]

Some meteorites provide evidence that liquid water was widespread and very active during the early history of some meteorite parent bodies-asteroids and comets. Characteristic secondary minerals are hydrous minerals such as serpentines, smectites, micas, and tochilinite, as well as sulfides, oxides, phosphates and carbonates (e.g., Brearley 2006). Hydrous phases are observed spectroscopically on C-complex asteroids (C-, G-, F- and B-type asteroids) that are plausibly to the source of some carbonaceous chondrites (e.g., Vilas and Gaffey 1989). The degree of aqueous alteration is classified into petrologic types 1 to 3, based on the abundances of chondrule glass and metal, sulfide compositions, matrix mineralogy and quantity, carbon and water content, and stable isotopic compositions (Van Schmus and Wood 1967). Type 1 is most aqueously altered and type 3 is least altered. CI, CM and CR carbonaceous chondrites are extensively aqueously altered. CI group contains type 1, the CM group contains type 1 and 2, and the CR group contains type 1 to 3. CM chondrites have been further assigned to subtype, least altered CM2.6 to most altered CM2.0 (=CM1) (Rubin et al. 2007). There are no type 1 and 2 meteorites in other groups of carbonaceous chondrites (e.g., CV and CO) and ordinary chondrites, however some samples of all of these latter groups show some degree of aqueous alteration (e.g., Grossman et al. 2000). In many cases the effects of aqueous alteration have been obscured by subsequent thermal metamorphism (Tonui et al. 2014).

The most widely accepted location of aqueous alteration is meteorite parent bodies. There are also possibilities for limited alteration with water ice or vapour in the nebula (e.g., Ciesla et al. 2003). After accretion of planetesimals, water ice melted due to either heating by decay of short-lived radioactive nuclides, impacts, or solar heating, triggering aqueous alteration. Isotopic chronologies based on Mn-Cr isotope measurements of carbonates revealed that aqueous alteration occurred as long as $4563.4 \pm 0.4 / -0.5$ Myr ago (~ 4.8 Myr after calcium-aluminium-rich inclusion (CAI) formation) for CM chondrites (Fujiya et al. 2012). However, these dates are now being reassessed.

Aqueous alteration models have been constructed based on mineralogic assemblages, stable isotopes, and thermal evolution (e.g., Clayton and Mayeda 1999; Grimm and Mccween 1989; Zolensky et al. 1989). Depending on the heating mechanism, thermal evolution largely depends on the canonical values of ^{26}Al , porosity, water/rock ratio and size of body, and orbital histories. Conditions of aqueous alteration varied for the different meteorite groups. Temperatures have been estimated as $\sim 20\text{-}150$ °C for CI, $0\text{-}80$ °C for CM2, and $\sim 50\text{-}150$ °C for CR (Brearley 2006 and references there in). Estimation of water/rock ratios are 1.1-1.2 with $\text{pH} = 7\text{-}10$ and $f\text{O}_2 > 10^{-55}\text{-}10^{-70}$ for CI, and water/rock ratio was 0.3-0.6 with $\text{pH} = 7\text{-}12$ and $f\text{O}_2 < 10^{-85}$ for CM2 (Zolensky et al. 1993; Zolensky et al. 1989). Again, these estimates depend on model assumptions. During the aqueous alteration, formation and evolution of organic matter would be expected. For example, during early stages of aqueous alteration, macromolecular organic matter and amino acids can be produced from simple molecules such as formaldehyde and ammonia (e.g., Cody et al. 2011; Kebukawa et al. 2017). Further alteration modifies the compositions of organic matter, e.g., decreases H,N,O contents and aliphatics and oxygen-bearing functional groups (e.g., Alexander et al. 2014; Alexander et al. 2007). Among amino acids, ω -amino acids dominate in heavily-altered chondrites in contrast to α -amino acids which dominate in the less-altered ones, probably due to lower stabilities of α -amino acids and/or formation of ω -amino acids by Fischer-Tropsch-type (FTT) reaction (e.g., Elsila et al. 2016). Larger L-isovaline enrichments in the aqueously-altered compared to less-altered meteorites suggest that aqueous alteration promoted L-isovaline excesses on meteorite parent bodies (Glavin and Dworkin 2009).

Aqueous alteration provided abundant water in the form of hydrated minerals (e.g., Alexander et al. 2012), as well as a diverse suite of organic compounds formed in parent bodies in addition to pre-accretional organic matter (e.g., Schmitt-Kopplin et al. 2010) to the early Earth.

Cross-References

[Add Cross-references to related entries within this reference work; the current table of contents is available for download from the Meteor website.

Please enter cross-references here]

→Meteorites

→Asteroid

→Comets

Further Reading

Brearley AJ (2006) The action of water. In: Laurretta DS, McSween JHY (eds) *Meteorites and the Early Solar System II*. University of Arizona Press, Tucson, pp 587-624

References

[A maximum of 20 references. Please enter your text here...]

- Alexander CM, Bowden R, Fogel ML, Howard KT, Herd CD, Nittler LR (2012) The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets *Science* 337:721-723 doi:10.1126/science.1223474
- Alexander CMOD et al. (2014) Elemental, isotopic, and structural changes in Tagish Lake insoluble organic matter produced by parent body processes *Meteoritics & Planetary Science* 49:503-525 doi:10.1111/maps.12282
- Alexander CMOD, Fogel M, Yabuta H, Cody GD (2007) The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter *Geochimica et Cosmochimica Acta* 71:4380-4403 doi:10.1016/j.gca.2007.06.052
- Brearley AJ (2006) The action of water. In: Laurretta DS, McSween JHY (eds) *Meteorites and the Early Solar System II*. University of Arizona Press, Tucson, pp 587-624
- Ciesla FJ, Laurretta DS, Cohen BA, Hood LL (2003) A nebular origin for chondritic fine-grained phyllosilicates *Science* 299:549-552
- Clayton RN, Mayeda TK (1999) Oxygen isotope studies of carbonaceous chondrites *Geochimica et Cosmochimica Acta* 63:2089-2104
- Cody GD, Heying E, Alexander CMO, Nittler LR, Kilcoyne ALD, Sandford SA, Stroud RM (2011) Establishing a molecular relationship between chondritic and cometary organic solids *Proceedings of the National Academy of Sciences of the United States of America* 108:19171-19176 doi:10.1073/pnas.1015913108
- Elsila JE, Aponte JC, Blackmond DG, Burton AS, Dworkin JP, Glavin DP (2016) Meteoritic amino acids: Diversity in compositions reflects parent body histories *ACS Central Science* 2:370-379 doi:10.1021/acscentsci.6b00074
- Fujiya W, Sugiura N, Hotta H, Ichimura K, Sano Y (2012) Evidence for the late formation of hydrous asteroids from young meteoritic carbonates *Nature Communications* 3 doi:10.1038/ncomms1635
- Glavin DP, Dworkin JP (2009) Enrichment of the amino acid L-isovaline by aqueous alteration on CI and CM meteorite parent bodies *Proceedings of the National Academy of Sciences of the United States of America* 106:5487-5492 doi:10.1073/pnas.0811618106

- Grimm RE, Mccween HY (1989) Water and the thermal evolution of carbonaceous chondrite parent bodies *Icarus* 82:244-280
- Grossman JN, ALEXANDER CMD, Wang J, Brearley AJ (2000) Bleached chondrules: Evidence for widespread aqueous processes on the parent asteroids of ordinary chondrites *Meteoritics & Planetary Science* 35:467-486
- Kebukawa Y, Chan QHS, Tachibana S, Kobayashi K, Zolensky ME (2017) One-pot synthesis of amino acid precursors with insoluble organic matter in planetesimals with aqueous activity *Science Advances* 3:e1602093 doi:10.1126/sciadv.1602093
- Rubin AE, Trigo-Rodriguez JM, Huber H, Wasson JT (2007) Progressive aqueous alteration of CM carbonaceous chondrites *Geochimica et Cosmochimica Acta* 71:2361-2382 doi:10.1016/j.gca.2007.02.008
- Schmitt-Kopplin P et al. (2010) High molecular diversity of extraterrestrial organic matter in Murchison meteorite revealed 40 years after its fall *Proceedings of the National Academy of Sciences of the United States of America* 107:2763-2768 doi:10.1073/pnas.0912157107
- Tonui E, Zolensky M, Hiroi T, Nakamura T, Lipschutz ME, Wang M-S, Okudaira K (2014) Petrographic, chemical and spectroscopic evidence for thermal metamorphism in carbonaceous chondrites I: CI and CM chondrites *Geochimica et Cosmochimica Acta* 126:284-306 doi:10.1016/j.gca.2013.10.053
- Van Schmus W, Wood JA (1967) A chemical-petrologic classification for the chondritic meteorites *Geochimica et Cosmochimica Acta* 31:747-765
- Vilas F, Gaffey MJ (1989) Phyllosilicate Absorption Features in Main-Belt and Outer-Belt Asteroid Reflectance Spectra *Science* 246:790-792 doi:10.1126/science.246.4931.790
- Zolensky M, Barrett R, Browning L (1993) Mineralogy and composition of matrix and chondrule rims in carbonaceous chondrites *Geochimica et Cosmochimica Acta* 57:3123-3148
- Zolensky ME, Bourcier WL, Gooding JL (1989) Aqueous alteration on the hydrous asteroids - Results of EQ3/6 computer-simulations *Icarus* 78:411-425 doi:10.1016/0019-1035(89)90188-7