

# Astronomy




Abell 1758  
Galaxy Cluster  
~3.2 billion ly

From <http://chandra.harvard.edu>

A Presentation for Arrow Head District Merit Badge Day

Mitzi Adams, Dr. Dennis Gallagher, Dr. Michael Zanetti  
NASA/MSFC  
March 9, 2019

# Putting it into Context \*Astronomical\* Scales



## Time, Distance Size

How big is a million, a billion, 13.8 billion ?

Count numbers, consider each number as one second.

Count to one million -- 11.6 days


Count to one billion -- Multiply 11.6 days by 1000 = 32 years

Count to 13.8 billion --> 439 years


# Optical Telescopes

Reflector  
Newtonian


Mirrors



Schmidt-Cassegrain




Refractor  
First Used by Galileo to do Astronomy




Lenses

Astronomical



# Chandra X-ray Optics



The alignment of the mirrors from one end of the mirror assembly to the other (2.7 meters or 9 feet) is accurate to 1.3 micrometers (50 millionths of an inch) or about one fiftieth the width of a human hair!

# Constellations and Bright Stars

# Polaris (α UMi) - HIP 11767

Magnitude: **1.95** (B-V: 0.63)  
Absolute Magnitude: -3.66  
RA/DE (J2000): 2h31m50.9s/+89°15'51.4"  
RA/DE (of date): 2h55m52s/+89°20'41"  
Hour angle/DE: 3h13m33s/+89°20'41"  
Az/Alt: +359°24'04"/+35°08'37"  
Spectral Type: F7:lb-IIv\_\*SB  
Distance: 431.42 Light Years  
Parallax: 0.00756

Meridian

Ursa Major

Ursa Minor

Cepheus

Draco

Lacerta



# Ecliptic, Equator, and Meridian





Image from Wikipedia: Attribution-ShareAlike 3.0 Unported (CC BY-SA 3.0)



# Regulus ( $\alpha$ Leo) - HIP 49669


Magnitude: **1.35** (B-V: -0.09)  
Absolute Magnitude: -0.53  
RA/DE (J2000): 10h08m22.1s/+11°58'02.9"  
RA/DE (of date): 10h09m24s/+11°52'23"  
Hour angle/DE: 20h00m4s/+11°52'23"  
Az/Alt: +97°22'14"/+31°18'10"  
Spectral Type: B7V  
Distance: 77.49 Light Years  
Parallax: 0.04209"





# Alnitak ( $\zeta$ Ori) - HIP 26727 A

Magnitude: **1.85** (B-V: -0.09)  
Absolute Magnitude: -5.15  
RA/DE (J2000): 5h40m45.5s/-1°56'33.2"  
RA/DE (of date): 5h41m44s/-1°56'02"  
Hour angle/DE: 0h27m43s/-1°56'02"  
Az/Alt: +191°30'08"/+52°47'01"  
Spectral Type: O9.5Ib\_SB  
Distance: 817.43 Light Years  
Parallax: 0.00399"



S


# Capella ( $\alpha$ Aur) - HIP 24608

Magnitude: **0.05** (B-V: 0.79)  
Absolute Magnitude: -0.51  
RA/DE (J2000): 5h16m41.6s/+45°59'48.0"  
RA/DE (of date): 5h18m7s/+46°00'59"  
Hour angle/DE: 0h37m41s/+46°00'59"  
Az/Alt: +330°33'13"/+76°37'42"  
Spectral Type: M1: comp  
Distance: 42.20 Light Years  
Parallax: 0.07729"



# Alrescha ( $\alpha$ Psc) - HIP 9487 A

Magnitude: **4.10** (B-V: -0.03)  
Absolute Magnitude: 0.95  
RA/DE (J2000): 2h02m2.9s/+2°45'49.3"  
RA/DE (of date): 2h03m3s/+2°51'19"  
Hour angle/DE: 4h06m25s/+2°51'19"  
Az/Alt: +255°21'33"/+24°45'39"  
Spectral Type: A2  
Distance: 139.09 Light Years  
Parallax: 0.02345"



Aldebaran  
Taurus

Mars

Moon

W

# Great Nebula in Andromeda (M 31 - NGC 224)

Type: **Galaxy**  
Magnitude: **3.50**  
RA/DE (J2000): 0h42m42.0s/+41°16'00.0"  
RA/DE (of date): 0h43m45s/+41°22'17"  
Hour angle/DE: 5h25m42s/+41°22'17"  
Az/Alt: +302°52'31"/+27°55'35"  
Size: +2°58'00"

Uranus

Aries

Cassiopeia


Andromeda

Pisces


# A Beautiful Binary Star System Albireo

430 ly away  
Look for Cygnus in summer

Albireo A is a binary, the yellowish color comes from a star with spectral type, K2, temperature ~ 4000 K  
Albireo B is blue, type B8, temperature ~13,000K



# Our Dynamic Sun: A Star at the Center of the Solar System




Planet  
Nine?


**What is a Star?**

# What is a Star?

A star is an astrophysical body that produces its own light by thermonuclear reactions in its core.



Betelgeuse: A red giant star, about 600 ly away, 3500 K, 1,180  $R_{\odot}$ , 7.7  $M_{\odot}$ .



To produce energy, hydrogen converts to Helium




Rigel: A blue-white star, about 770 ly away, 11,000 K, 80  $R_{\odot}$ , 20  $M_{\odot}$ .



# Stars Classified According to Color (Temperature)

Brighter



Cooler

OBAFGKM

Overseas Broadcast A Flash, Godzilla Kills Mothra

# Layers of the Sun

## The Convection Zone

Energy continues to move toward the surface through convection currents of heated and cooled gas in the convection zone.

## The Corona

The ionized elements within the corona glow in the x-ray and extreme ultraviolet wavelengths. NASA instruments can image the Sun's corona at these higher energies since the photosphere is quite dim in these wavelengths.

## The Radiative Zone

Energy moves slowly outward—taking more than 170,000 years to radiate through the layer of the Sun known as the radiative zone.

## Sun's Core


Energy is generated by thermonuclear reactions creating extreme temperatures deep within the Sun's core.

## Coronal Streamers

The outward-flowing plasma of the corona is shaped by magnetic field lines into tapered forms called coronal streamers, which extend millions of miles into space.

## The Chromosphere

The relatively thin layer of the Sun called the chromosphere is sculpted by magnetic field lines that restrain the electrically charged solar plasma. Occasionally larger plasma features—called prominences—form and extend far into the very tenuous and hot corona, sometimes ejecting material away from the Sun.






# Sunspots

Sunspots are dark (and cooler) regions on the surface of the Sun. They have a darker inner region (the Umbra) surrounded by a lighter ring (the Penumbra).



Sunspots usually appear in groups that form over hours or days and last for days or weeks.


The earliest sunspot observations (c. 1609) indicated that the Sun rotates once in about 27 days.



# Sunspots

## Examples





# 23 Full Cycles

Heinrich Schawbe discovered (1844) there was a cycle of sunspot number.

The average cycle lasts about 11 years, but ranges from 9 to 14.

The average maximum number is about 100, but ranges from 50 to 200.

# The Corona and the Solar Cycle




28 Sep 1991

27 Mar 1992

26 Jan 1993

04 Nov 1993

20 Sep 1994



12 Nov 1966

07 Mar 1970

20 Jun 1973


11 Mar 1985

18 Mar 1988



[SMM Coronagraph]

# Solar Eruptions

Solar Flares and Coronal Mass Ejections (CMEs)



This combo of SDO and Soho C2 shows X2-flare and a halo CME





Three distinct CMEs: First to right, second from north pole, third from far side of Sun. All three eruptions happened within hours of each other.

Animations!

c2\_halloween\_2003.mpg, c3\_halloween\_2003.mpg, X2\_C2\_combo\_best.mpg

## Sunspots

Sunspots are comparatively cool areas at up to 7,700° F and show the location of strong magnetic fields protruding through what we would see as the Sun's surface. Large, complex sunspot groups are generally the source of significant space weather.




## Coronal Mass Ejections (CMEs)

Large portions of the corona, or outer atmosphere of the Sun, can be explosively blown into space, sending billions of tons of plasma, or superheated gas, Earth's direction. These CMEs have their own magnetic field and can slam into and interact with Earth's magnetic field, resulting in geomagnetic storms. The fastest of these CMEs can reach Earth in under a day, with the slowest taking 4 or 5 days to reach Earth.


## Solar Wind

The solar wind is a constant outflow of electrons and protons from the Sun, always present and buffeting Earth's magnetic field. The background solar wind flows at approximately one million miles per hour!



## Solar Flares

Reconnection of the magnetic fields on the surface of the Sun drive the biggest explosions in our solar system. These solar flares release immense amounts of energy and result in electromagnetic emissions spanning the spectrum from gamma rays to radio waves. Traveling at the speed of light, these emissions make the 93 million mile trip to Earth in just 8 minutes.



# Space Weather


Space weather refers to the variable conditions on the Sun and in the space environment that can influence the performance and reliability of space-based and ground-based technological systems, as well as endanger life or health. Just like weather on Earth, space weather has its seasons, with solar activity rising and falling over an approximate 11 year cycle.

## Sun's Magnetic Field

Strong and ever-changing magnetic fields drive the life of the Sun and underlie sunspots. These strong magnetic fields are the energy source for space weather and their twisting, shearing, and reconnection lead to solar flares.

## Solar Radiation Storms

Charged particles, including electrons and protons, can be accelerated by coronal mass ejections and solar flares. These particles bounce and gyrate their way through space, roughly following the magnetic field lines and ultimately bombarding Earth from every direction. The fastest of these particles can affect Earth tens of minutes after a solar flare.




## Geomagnetic Storms

A geomagnetic storm is a temporary disturbance of Earth's magnetic field typically associated with enhancements in the solar wind. These storms are created when the solar wind and its magnetic field interacts with Earth's magnetic field. The primary source of geomagnetic storms is CMEs which stretch the magnetosphere on the nightside causing it to release energy through magnetic reconnection. Disturbances in the ionosphere (a region of Earth's upper atmosphere) are usually associated with geomagnetic storms.

## Earth's Magnetic Field

Earth's magnetic field, largely like that of a bar magnet, gives the Earth some protection from the effects of the Sun. Earth's magnetic field is constantly compressed on the day side and stretched on the night side by the ever-present solar wind. During geomagnetic storms, the disturbances to Earth's magnetic field can become extreme. In addition to some buffering by the atmosphere, this field also offers some shielding from the charged particles of a radiation storm.





# Aurorae




Seen mostly at high latitudes, aurorae are produced when Earth's magnetosphere is disturbed.

Plasma from the magnetosphere precipitates into the upper atmosphere.

Reds are from oxygen  
Greens are from lower in atmosphere.





Energetic Electrons

Solar Flare Protons

Damage to Spacecraft Electronics

Ionospheric Currents

GPS Signal Scintillation

Radiation Effects on Avionics

Geomagnetically Induced Current in Power Systems

Induced Effects in Submarine Cables

Telluric Currents in Pipelines

# What Is an Eclipse?


An eclipse happens when one object blocks light from falling onto another object. The shadow of the eclipsed object falls onto the other object.



©2004 F. Espenak

www.MrEclipse.com

## LUNAR ECLIPSE GEOMETRY



www.MrEclipse.com

©2000 F. Espenak

Images Used With Permission

# Phases of the Moon



Image Credit: NASA/Bill Dunford


Third Quarter

# Solar Eclipses



www.MrEclipse.com

©1999 F. Espenak



Images Used With Permission

# What You Can See



Zophia Edwards wide-angle view, from Jay Pasachoff's Eclipse 2013 page

Image Used With Permission

# The Corona and Prominences





Rob Lucas, with Jay Pasachoff's 2013 Eclipse Expedition

# How to Safely Observe An Eclipse

No Special Rules for Lunar Eclipses


For Solar Eclipses:

Projection  
Special Telescope Filters  
Eclipse Glasses  
Number 14 Welder's Glass






# Use a Kitchen Colander For Partial Phases



# Eclipse Glasses and Welder's Glass



Number 14 or Higher!

# Von Braun Astronomical Society in Monte Sano State Park Observatories and Planetarium



Planetarium Program each Saturday night at 7:30 p.m.  
Telescope Observing after, weather permitting