


Investigating diurnal changes in the normal albedo of the lunar surface at 1064 nm: A new analysis with the Lunar Orbiter Laser Altimeter



Ariel N. Deutsch¹, Gregory A. Neumann², James W. Head¹, Paul G. Lucey³


¹Department of Earth, Environmental and Planetary Sciences, Brown University, Providence RI 02912 (ariel_deutsch@brown.edu)

²NASA Goddard Space Flight Center, Greenbelt, MD 20771

³Hawaii Institute of Geophysics and Planetology, University of Hawaii at Manoa, Honolulu, HI 96822


Introduction

The thermal environment of the lunar surface is extreme. At the equator, temperatures drop ~300 K between local noon and night [1].



• Laboratory studies demonstrate that minerals common to the lunar surface (e.g., pyroxene, olivine) show spectral changes with respect to temperature in near infrared wavelengths [2–4].

• Over temperature changes equivalent to the lunar thermal environment ($\Delta T \approx 300\text{K}$), the reflectance of pure pyroxene samples can vary by a factor of two [4].



Motivating question


How does the surface reflectance of the Moon as measured from orbit by LOLA change during extreme temperature fluctuations experienced by the surface over the course of a lunar day?

Methods

Here we analyze the LOLA data [5] for differences in mean normal albedo during the cycle of the lunar day.

Two groups are selected to represent maximum and minimum surface temperatures:

- Mid-day: 11:00–13:00
- Morning/Evening: 06:00–07:00; 16:00–17:00




We target $1^\circ \times 1^\circ$ regions of interest (ROIs) within the mare and highlands between 65°S and 65°N , latitudes between which temperature fluctuations are greatest.

To date, our analysis includes 65 ROIs located within the maria and 383 ROIs located within the highlands (*below*).

Results

MARE


- 78% of ROIs in the mare display an inverse relationship between normal albedo and temperature (*right*).
- The overall measured change in normal albedo is relatively low, on the order of a few % change.



Caption.

HIGHLANDS

- There is no statistically significant temperature-dependent reflectance change detected.
- Only ~49% of ROIs show a decrease in normal albedo during local mid-day (*left*).




ROIs analyzed in the
■ Mare ■ Highlands

Basemap is LOLA hillshade in Robinson projection.

Temperature-dependent albedo change

- The albedo change measured in the maria is consistent with laboratory studies [2–4].
- Previous laboratory measurements of returned lunar soils revealed a change in relative reflectance with temperature of ~1% or less per 100 K near-IR wavelengths [4].

The normal albedo of regions within the maria tends to decrease during local mid-day, when temperatures are highest.



No temperature-dependent albedo change

- It is possible that we do not detect a clear temperature-dependent albedo change in the highlands due to a variety of factors:

1. Surfaces that are low in iron will show a weaker change because iron is responsible for the temperature-dependent absorption near 1064 nm [4].
2. The reflectance may be affected by grain size effects, where particularly fine-grained regions have a decreased reflectance in comparison to a region of similar composition with larger grains [6,7].
3. Mature soils show less contrast due to the attenuating effect of submicroscopic iron that has accumulated through time [8,9].

Conclusions

- Our statistical analysis, incorporating over 200,000 individual LOLA shots, suggests that temperature variations have a measurable effect on the normal albedo of the surface at 1064 nm wavelength in the maria, and this may be due to temperature-induced spectral changes.
- The diurnal differences are only on the order of a few % change in normal albedo, indicating that temperature changes do not have a large effect on LOLA measurements.
- An ability to understand how the lunar surface varies with temperature will provide important constraints for future remote sensing observations of the Moon [e.g., 10,11].
- Such observations can help constrain the relative abundance of particular minerals (here, pyroxene) that exhibit a change in spectral reflectance with temperature independent of spectroscopic methods.

References

- [1] Williams J.-P. et al. (2017) Icarus, 283, 300–325. [2] Singer R.B. and Roush T.L. (1985) JGR, 90, 12434–12444. [3] Roush T.L. and Singer R.B. (1986) JGR, 91, 10301–10308. [4] Hinrichs J.L. and Lucey P.G. (2002) Icarus, 155, 169–180. [5] Lemelin M. et al. (2016) Icarus, 273, 315–328. [6] Hapke, B. (1993). Cambridge University Press, NY. [7] Clark R.N. and Roush T.L. (1984). JGR, 89, 6329–6340. [8] Hapke B. (2001) JGR, 106, 10039–10074. [9] Pieters C.M. (2000) MAPS, 35, 1101–1107. [10] Cohen B.A. et al. (2015) LEAG, Abstract #2008. [11] Lucey P.G. et al. (2017) LEAG, Abstract #5048.