

Investigating Tropical Cyclone Size and Integrated Kinetic Energy using CYGNSS and Other Datasets

Patrick Duran¹ and Dan Cecil²

1. University of Alabama in Huntsville

2. NASA MSFC

patrick.duran@uah.edu

CYGNSS Science Team Meeting

6 June 2019

Ann Arbor, MI

Integrated Kinetic Energy (IKE)


- A tropical cyclone (TC) intensity metric first proposed by Powell and Reinhold (2007):

$$IKE = \int_V \frac{1}{2} \rho U^2 dV$$

- Accounts for both maximum wind speed and the spatial extent of the surface wind field.
- Can be a better measure of destructive potential than maximum wind speed – particularly for large TCs.


The value of IKE

Hurricane Camille (1969)



Camille was stronger
in terms of V_{max} .


Hurricane Katrina (2005)



H*Wind analyses from
NOAA/AOML
Hurricane Research Division

The value of IKE

Hurricane Camille (1969)




Camille was stronger in terms of V_{max} .

Katrina's larger wind field made it much more destructive.

Damage (2017 dollars)
Camille: \$9.8 billion
Katrina: \$160 billion


H*Wind analyses from
NOAA/AOML
Hurricane Research Division

Hurricane Katrina (2005)



The value of IKE

Hurricane Camille (1969)



Camille was stronger
in terms of V_{max} .


Katrina's larger wind field
made it much more
destructive.

IKE (Powell & Reinhold)

Camille: 63 Terajoules
Katrina: 122 Terajoules


H*Wind analyses from
NOAA/AOML
Hurricane Research Division

Hurricane Katrina (2005)




The value of IKE

Hurricane Camille (1969)



Camille was stronger
in terms of V_{max} .

Hurricane Katrina (2005)



IKE provided a better representation of Katrina's destructive potential than V_{max}

IKE (Powell & Reinhold)

Camille: 63 Terajoules
Katrina: 122 Terajoules

H*Wind analyses from
NOAA/AOML
Hurricane Research Division

IKE Computation

- Assume integration over a 1-m depth:

$$IKE = \frac{\rho_0}{2} \int_0^{2\pi} \int_0^R u(\theta, r)^2 r dr d\theta$$

- Requires knowledge of the velocity at every (θ, r) .
 - Multiple methods possible:
 - Use a data assimilation scheme (e.g. H*WIND) or model analysis.
 - Fit observations to a parametric wind profile (e.g. Morris and Ruf).
 - Piecewise polynomial interpolation (e.g. tension splines).
 - Azimuthally average observations to get a radial profile of velocity.

IKE Computation






- Assume integration over a 1-m depth:

$$IKE = \frac{\rho_0}{2} \int_0^{2\pi} \int_0^R u(\theta, r)^2 r dr d\theta$$

- Requires knowledge of the velocity at every (θ, r) .
 - Multiple methods possible:
 - Use a data assimilation scheme (e.g. H*WIND) or model analysis.
 - Fit observations to a parametric wind profile (e.g. Morris and Ruf).
 - Piecewise polynomial interpolation (e.g. tension splines).
 - Azimuthally average observations to get a radial profile of velocity.






Constructing the radial wind profile

1. Start with an estimate of the radial wind structure using operational wind radii from the *Extended Best Track Dataset*.



Constructing the radial wind profile


1. Start with an estimate of the radial wind structure using operational wind radii from the *Extended Best Track Dataset*.



34 kt


Constructing the radial wind profile

- Gather all observations collected within 3 hours and 500 km of the best-track storm center from **CYGNSS**, **SFMR**, **ASCAT**, and **SMAP**.
 - CYGNSS v2.1**: NBRCS wind retrievals using only the YSLF GMF. All winds with “uncertainty” $> 3.5 \text{ m s}^{-1}$ filtered out.




Constructing the radial wind profile

Removing all observations with “uncertainty” (standard deviation of error) $> 3.5 \text{ m s}^{-1}$ eliminates unrealistically large wind speeds without removing too many good observations.



Constructing the radial wind profile

2. Gather all observations collected within 3 hours and 500 km of the best-track storm center from **CYGNSS**, **SFMR**, **ASCAT**, and **SMAP**.
- **SFMR**: All wind retrievals that did not have any QC flag flipped.
 - **ASCAT**: All wind retrievals that did not have the product monitoring, KNMI, or variational QC flags flipped.




Constructing the radial wind profile

2. Gather all observations collected within 3 hours and 500 km of the best-track storm center from **CYGNSS**, **SFMR**, **ASCAT**, and **SMAP**.


SMAP winds from
Remote Sensing
Systems
(Meissner et al. 2017)

Hurricane Irma
9/5/2017
11 UTC



Constructing the radial wind profile

3. Transform observation locations into a storm-centered polar coordinate system, split up by quadrant, and azimuthally average.




Black Dots:
Individual
Wind
Observations

**Orange dotted
lines:**
Initial guess
wind profiles
from best track.

Blue Lines:
Azimuthally
averaged
wind profiles.

Computing IKE

4. Integrate kinetic energy in each quadrant, using only azimuthally averaged winds greater than 34 kt, and sum them to get total IKE.




Black Dots:
Individual
Wind
Observations

**Orange dotted
lines:**
Initial guess
wind profiles
from best track.

Blue Lines:
Azimuthally
averaged
wind profiles.

Computing IKE

4. Integrate kinetic energy in each quadrant, using only azimuthally averaged winds greater than 34 kt, and sum them to get total IKE.





Black Dots:
Individual
Wind
Observations

**Orange dotted
lines:**
Initial guess
wind profiles
from best track.

Blue Lines:
Azimuthally
averaged
wind profiles.

Potential Applications


- Is there a time of day when TCs exhibit higher IKE?



In an idealized simulation,
IKE maximizes at 9 AM and
minimizes at 9 PM.

Potential Applications

- Is there a time of day when TCs exhibit higher IKE?




In an idealized simulation, **IKE maximizes at 9 AM** and **minimizes at 9 PM**.

Related to a **radial expansion** of the TC wind field overnight and through the morning, and a contraction during the afternoon, into the evening.

Potential Applications

- Is there a time of day when TCs exhibit higher IKE?







Can we verify this with observations?

In an idealized simulation, **IKE maximizes at 9 AM** and **minimizes at 9 PM**.


Related to a **radial expansion** of the TC wind field overnight and through the morning, and a contraction during the afternoon, into the evening.

Hurricane Irma – September 2-3, 2017




Hurricane Irma – September 2-4, 2017


CYGNSS winds ($m s^{-1}$) within 500km and within 3 h prior to IF




CYGNSS winds ($m s^{-1}$) within 500km and within 3 h prior to I




CYGNSS winds ($m s^{-1}$) within 500km and within 3 h prior to IF




CYGNSS winds ($m s^{-1}$) within 500km and within 3 h prior to IF




ASCAT winds ($m s^{-1}$) within 500km and within 3 h prior to IRI




ASCAT winds ($m s^{-1}$) within 500km and within 3 h prior to IRI



ASCAT winds ($m s^{-1}$) within 500km and within 3 h prior to IRI




ASCAT winds ($m s^{-1}$) within 500km and within 3 h prior to IRI




Hurricane Irma – September 2-4, 2017


CYGNSS winds (m s^{-1}) within 500km and within 3 h prior to IF




CYGNSS winds (m s^{-1}) within 500km and within 3 h prior to I




CYGNSS winds (m s^{-1}) within 500km and within 3 h prior to IF




CYGNSS winds (m s^{-1}) within 500km and within 3 h prior to IF




ASCAT winds (m s^{-1}) within 500km and within 3 h prior to IRL




ASCAT winds (m s^{-1}) within 500km and within 3 h prior to IRI



ASCAT winds (m s^{-1}) within 500km and within 3 h prior to IRI



ASCAT winds (m s^{-1}) within 500km and within 3 h prior to IRL




Future Directions


- Use CYGNSS, ASCAT, and SMAP to construct climatologies of wind radii and IKE in TCs across the globe.
 - Stratify by TC intensity, vertical wind shear, ocean basin, etc.
- Investigate the diurnal cycle of TC wind radii and IKE using the climatologies as a reference point.
- Refine algorithm to combine observations from different platforms and construct radial wind profiles.
 - Weight observations by observation platform and/or average the wind speeds from each platform first, then take total average.
 - Account for differences in horizontal resolution.
 - Take maximum value in each radial bin instead of average.

Extra Slides

CYGNSS L3 WIND SPEED : 20170904 (1800Z-0000Z)
 AL11 [IRMA] : VMAX 115 KTS





UNIVERSITY OF MESA
 ROSENSTIEL
 SCHOOL OF MARINE &
 ATMOSPHERIC SCIENCE




- Retrievals where std dev of wind speed error ≥ 3.0 m/s denoted by hollow squares.
- v2.1. Young Seas with Limited Fetch (YSLF) algorithm
- Storm marker placed at beginning of 6-hour window.

CYGNSS winds (m s^{-1}) within 300km and 3 h



IKE History – Hurricane Irma (2017)

- Compute IKE every hour, using 6 hours of observations (all observations within 3 hours before or after best track time).




IKE History – Hurricane Irma (2017)

IRMA | 08/31/2017 00:00 UTC | V_{max} 28.3 m s⁻¹ (55 kt) | RMW 28 km


CYGNSS winds (m s⁻¹) within 500km and 3 h

CYGNSS




SFMR winds (m s⁻¹) within 500km and 3 h

SFMR




ASCAT winds (m s⁻¹) within 500km and 3 h

ASCAT




Total IKE (TJ)

Total IKE




Observations (dots) and azi-averaged wind (line) for NW quadrant

$IKE_{34} = 0.452$ TJ




Observations (dots) and azi-averaged wind (line) for NE quadrant

$IKE_{34} = 0$ TJ




Observations (dots) and azi-averaged wind (line) for SW quadrant

$IKE_{34} = 0.109$ TJ




Observations (dots) and azi-averaged wind (line) for SE quadrant

$IKE_{34} = 0.823$ TJ




IRMA | 09/03/2017 19:00 UTC | V_{max} 51.4 m s⁻¹ (100 kt) | RMW 28 km


CYGNSS winds (m s⁻¹) within 500km and 3 h








SFMR winds (m s⁻¹) within 500km and 3 h



ASCAT winds (m s⁻¹) within 500km and 3 h




Total IKE (TJ)




IRMA | 09/03/2017 20:00 UTC | V_{max} 51.4 m s⁻¹ (100 kt) | RMW 28 km


CYGNSS winds (m s⁻¹) within 500km and 3 h








SFMR winds (m s⁻¹) within 500km and 3 h



ASCAT winds (m s⁻¹) within 500km and 3 h




Total IKE (TJ)




IRMA | 09/03/2017 21:00 UTC | V_{max} 51.4 m s⁻¹ (100 kt) | RMW 28 km


CYGNSS winds (m s⁻¹) within 500km and 3 h








SFMR winds (m s⁻¹) within 500km and 3 h



ASCAT winds (m s⁻¹) within 500km and 3 h




Total IKE (TJ)




Where CYGNSS Adds Value

- When aircraft reconnaissance is unavailable (e.g. far from land).



Where CYGNSS Could be Improved



Where CYGNSS Could be Improved









IRMA | 08/31/20

CYGNSS winds ($m s^{-1}$) with

CYGNSS



CYGNSS rarely depicts (with confidence) the very high wind speeds that should be present in the inner core, *other than producing streaks of high wind speed that appear dubious*, and are filtered out.

TJ




Other Causes of Large IKE Fluctuations

- Presence of land in the averaging radii precludes observations from all platforms currently in the dataset.




IKE History – Hurricane Irma (2017)



- Observations typically produce smaller IKE estimate than best track wind radii.
 - A good thing.
 - Best track wind radii are the *maximum extent* of the winds in a given quadrant.

IKE History – Hurricane Irma (2017)



- Observations typically produce smaller IKE estimate than best track wind radii.
 - A good thing.
 - Best track wind radii are the *maximum extent* of the winds in a given quadrant.
- Sharp drops in IKE can occur when observations become available.
 - Sometimes good; sometimes not.
- Large temporal fluctuations are related to availability of observations, and are typically unphysical.

Extra Details on Best Track Radial Wind Profile

- Use RMW and V_{\max} from best track.
 - Assume that RMW is valid in quadrant with largest r_{34} , and scale the RMW by r_{34} in all of the other quadrants (i.e., a quadrant with a smaller r_{34} has a smaller RMW).
 - V_{\max} is the same in each quadrant, *unless* there is no corresponding wind radius (e.g., if $V_{\max} = 60$ kt, but there is no 50-kt wind radius defined in a quadrant, it does not make sense for v_{\max} to be 60 kt in that quadrant).
 - In this case, define V_{\max} in that quadrant to be 5 kt less than the lowest missing wind radius in that quadrant.
 - In the above example, V_{\max} would be 45 kt.