



Assimilation of Microwave Cloudy Radiances into NASA GEOS Model Using a Novel Bayesian Monte Carlo Technique

Isaac Moradi^{1,2} Frank Evans³,

William McCarty¹, Marangelly Cordero-Fuentes¹

1. GMAO/GSFC/NASA, 2. ESSIC, University of Maryland,
3. U. of Colorado

17th JCSDA Technical Review Meeting and Science Workshop,
May 29 - 31, 2019
Washington, DC, USA

May 29, 2019

This work is supported by the NASA Data for Operation and Assessment program, grant# NNX17AE89G. Many thanks to NDOA Program Manager, Dr. Tsengdar Lee.





Outline

Definitions

Motivation of the work

Bayesian Monte Carlo Integration (BMCI) technique

Implementation into NASA GEOS

Results

All-weather radiative transfer calculations

Cost function for 3D-Var Data Assimilation:

$$J(\vec{x}) = \overbrace{\frac{1}{2}(\vec{x} - \vec{x}_b)^T \vec{B}^{-1}(\vec{x} - \vec{x}_b)}^{J_b} + \overbrace{\frac{1}{2}(H(\vec{x}) - \vec{y})^T \vec{R}^{-1}(H(\vec{x}) - \vec{y})}^{J_o}$$

Relation between the observations (y) and the forward operator (H) can be expressed as: $y = H(\vec{x}, \vec{p}_b, \vec{p}_s) + \epsilon$

\vec{x} state vector, \vec{p}_b parameters such as shape and size distribution of hydrometers, \vec{p}_s indicates the scattering parameters (e.g., phase function)

$$\frac{dl_\nu}{dx} = -(\alpha_\nu + S_\nu)l_\nu + \alpha_\nu B_\nu(T) + S_\nu J_\nu$$

$$J_\nu = \int p_\nu(\Omega) l_\nu d\Omega$$



Limitations of direct assimilation of cloudy radiances

Inaccuracy in the first-guess: the models do not provide a close first guess for cloud parameters or clouds are often displaced.



Limitations of direct assimilation of cloudy radiances

Inaccuracy in the first-guess: the models do not provide a close first guess for cloud parameters or clouds are often displaced.


Lack of required RT inputs: $\vec{\rho}_s$ neither provided by the model nor fully measurable thus estimated from limited in-situ/aircraft measurements.

Limitations of direct assimilation of cloudy radiances

Inaccuracy in the first-guess: the models do not provide a close first guess for cloud parameters or clouds are often displaced.

Lack of required RT inputs: $\vec{\rho}_s$ neither provided by the model nor fully measurable thus estimated from limited in-situ/aircraft measurements.

Non-linearity in the forward model: \bar{x} is the mean value of the model variables within grid-box and because H is non-linear: $\overline{H(\vec{x})} \neq H(\bar{\vec{x}})$.



Limitations of direct assimilation of cloudy radiances

Inaccuracy in the first-guess: the models do not provide a close first guess for cloud parameters or clouds are often displaced.

Lack of required RT inputs: \vec{p}_s neither provided by the model nor fully measurable thus estimated from limited in-situ/aircraft measurements.

Non-linearity in the forward model: \bar{x} is the mean value of the model variables within grid-box and because H is non-linear: $\overline{H(\vec{x})} \neq H(\bar{\vec{x}})$.

Simplified RT models: Operational RT models that use a simplified RT framework, such as spherical hydrometeors, which is not appropriate at higher microwave frequencies where ice scattering is important.

Limitations of direct assimilation of cloudy radiances

- Inaccuracy in the first-guess:** the models do not provide a close first guess for cloud parameters or clouds are often displaced.
- Lack of required RT inputs:** \vec{p}_s neither provided by the model nor fully measurable thus estimated from limited in-situ/aircraft measurements.
- Non-linearity in the forward model:** \bar{x} is the mean value of the model variables within grid-box and because H is non-linear: $\overline{H(\vec{x})} \neq H(\bar{\vec{x}})$.
- Simplified RT models:** Operational RT models that use a simplified RT framework, such as spherical hydrometeors, which is not appropriate at higher microwave frequencies where ice scattering is important.
- Assuming Gaussian Errors:** DA systems assume Gaussian error statistics, examined using the departures, but in the case of cloudy radiances the departures are likely to be non-Gaussian.

The BMCI technique

The BMCI technique can be summarized in three steps:

- ▶ generation of a retrieval database of atmospheric state and cloud variables using a-priori information. The database should also include extreme cases as the extrapolation is not allowed.

The BMCI technique

The BMCI technique can be summarized in three steps:

- ▶ generation of a retrieval database of atmospheric state and cloud variables using a-priori information. The database should also include extreme cases as the extrapolation is not allowed.
- ▶ the atmospheric state and cloud variables are fed into the RT model to generate the synthetic observations. In addition to the state variables such as temperature, water vapor, and cloud profiles, cloud microphysics and parameterization such as particles' shape and size distribution are also utilized as input.



The BMCI technique

The BMCI technique can be summarized in three steps:

- ▶ generation of a retrieval database of atmospheric state and cloud variables using a-priori information. The database should also include extreme cases as the extrapolation is not allowed.
- ▶ the atmospheric state and cloud variables are fed into the RT model to generate the synthetic observations. In addition to the state variables such as temperature, water vapor, and cloud profiles, cloud microphysics and parameterization such as particles' shape and size distribution are also utilized as input.
- ▶ real measurements along with the generated database are given to the retrieval package, then the retrieval package will select the cases which are close to the real measurements and integrate them according to the Bayes' theorem to give the estimate of the mean and uncertainty of the state and cloud variables.

Beam filling

Beam filling was calculated as the difference between the brightness temperatures weighted according to an elliptical Gaussian beam pattern and T_b s calculated using the average profiles. The profiles were generated with 5km resolution using stochastic statistics derived from GPM DPR and central profiles IWP and rain rate.







Top: SkinTemp (left), IWP (right), Bottom: Rain WP (left), Surface Wind Speed (right)

Correlated observation errors

Retrieved Uncertainty Correlation Matrices




SST Analysis






SST Analysis


Mean
obs minus forecast



Std
obs minus forecast



Analysis Intensity Error



Hurricane Irma

TCVital Obs

GEOS0per_C360C

TBATMSTBGMI_C360C

TBATMSTBGMI_thin_C360C

GBATMSGBGMI_C360C


GBATMSGBGMI_thin_C360C

NoSatellite_C360C

TOnly_C360C

CONVonly_C360C

Forecast Intensity Error



GEOS0per_C360C

GBATMSGBGMI_C360C

TBATMSTBGMI_C360C


TBATMSTBGMI_thin_C360C

NoSatellite_C360C

GBATMSGBGMI_thin_C360C


CONVonly_C360C

Forecast Track Error




Irma Track in GEOS-5 Forecast

NoSatellite_C360C




Irma Track in GEOS-5 Forecast

GEOSOper_C360C




Irma Track in GEOS-5 Forecast


GBATMSGBGMI_thin_C360C



Wind speed (km/h) profiles




GEOSoper_C360C




GBATMSGBGMI_thin_C360C

Temperature anomaly



GEOSoper_C360C



GBATMSGBGMI_thin_C360C

Conclusions

- ▶ Conventional data assimilation schemes cannot properly assimilate satellite radiances in the rainband of tropical cyclones due to inaccuracy in RT scattering parameters as well as inaccuracy in the first guess provided by NWP models
- ▶ A new technique is proposed that does not depend on the minimization of the cost function.
- ▶ Preliminary results from BMCI technique are encouraging but require extensive validation, though validation itself is challenging
- ▶ These retrieved profiles are valuable for both analyzing the structure of the hurricanes as well as to provide more accurate initial conditions for the NWP models

**Thank you for
your attention!**

