

# GAMMA RAYS IN THE MODERN MULTI-MESSENGER ASTRONOMY ERA

C. Michelle Hui  
NASA/MSFC

Michigan Technological University  
Physics Colloquium  
Sep 9, 2019




# OUTLINE

- Gamma-ray Bursts and Gravitational Waves
  - binary neutron star merger
    - GRB 170817A / GW170817
    - other similar GRBs
- The Gamma-ray Burst Monitor on the Fermi Gamma-ray Space Telescope
  - gamma-ray transients
- Future instruments



# GAMMA RAY BURSTS

Jet collides with  
ambient medium  
(external shock wave)





# GAMMA RAY BURSTS

Jet collides with  
ambient medium  
(external shock wave)

Collapse of a massive star or merger of two compact objects.



Collimated relativistic outflow.

Prompt keV-MeV emission, afterglow in other wavelengths.

Detected ~ once per day, distributed all over the sky.

Colliding shells emit  
low-energy gamma rays  
(internal shock wave)

Slower  
Faster  
shell



Afterglow




# BINARY NEUTRON STAR MERGERS






2017-08-17

Fermi



LIGO






# 2017-08-17

```

////////////////////
TITLE:      GCN/FERMI NOTICE
NOTICE_DATE: Thu 17 Aug 17 12:41:20 UT
NOTICE_TYPE: Fermi-GBM Alert
RECORD_NUM: 1
TRIGGER_NUM: 524666471
GRB_DATE:   17982 TJD; 229 DOY; 17/08/17
GRB_TIME:   45666.47 SOD {12:41:06.47} UT
TRIGGER_SIGNIF: 4.8 [sigma]
TRIGGER_DUR: 0.256 [sec]
E_RANGE:    3-4 [chan] 47-291 [keV]
ALGORITHM:  8
DETECTORS:  0,1,1, 0,0,1, 0,0,0, 0,0,0, 0,0,
LC_URL:     http://heasarc.gsfc.nasa.gov/FTP/fermi/data/gbm/triggers/2017/
bn170817529/quicklook/glg_lc_medres34_bn170817529.gif
COMMENTS:   Fermi-GBM Trigger Alert.
COMMENTS:   This trigger occurred at longitude,latitude = 321.53,3.90 [deg].
COMMENTS:   The LC_URL file will not be created until ~15 min after the trigger.
  
```



GBM Alert

First On-board  
GBM  
Localization

LIGO Report of  
coincident GW/  
GRB

Joint LIGO/  
Virgo sky map

+16 s


+27 s

+45 min

+5 hour




2017-08-17



Reports of a blue optical transient near an elliptical galaxy NGC 4993 at ~40 Mpc (Abbot et al. 2017).

Discovery credit goes to Smartt et al. (2017) who observed the region with the 1m Swope telescope at Las Campanas Observatory



*Swift* observations reveal bright UV source, but no evidence of X-ray emission (Evans et al. 2017)

*NuStar* observations show no X-ray emission (Evans et al. 2017)

+12 hours

+13 hours

+14 hours




2017-08-17




*Chandra* observations  
show no X-ray emission  
(Fong et al. 2017)

Hubble  
observations  
reveal a reddening  
source  
(Adams et al. 2017)



*Chandra*  
observations  
reveal first evidence  
of delayed X-ray  
emission  
(Troja et al. 2017)



Radio counterpart  
reported by VLA  
(Mooley et al. 2017)

+2 days


+5 days

+9 days

+16.4 days



# GRB 170817A / GW170817



Binary neutron star merger and short gamma-ray burst association confirmed!





- GRB 170817A detected by GBM 1.7s after GW170817, a BNS merger event
  - extensive electromagnetic followup resulting in detection of a kilonova.
  - two components:
    - initial GRB spike — best fit Comptonized model with  $E_{\text{peak}} 185 \text{ keV}$**
    - weak thermal tail — blackbody  $kT \sim 10 \text{ keV}$**
- joint science:
  - tightest constraint on speed of gravity: gravitational waves and gamma rays travelled 130 million light years and arrived within 2 seconds  $\rightarrow$  consistent with speed of light within  $1e^{-15}$
  - constraints on neutron star equation of state
  - open questions: merger and jet geometry, intrinsic properties, population characteristics

Abbot et al. 2017, ApJ, 848, L13



# GRB 170817A

GRB 170817A appears to be a typical short gamma-ray burst.






Goldstein et al. 2017, ApJ, 848, L14




# GRB 170817A

GRB 170817A is the closest GRB ever detected but also the least luminous.





# GRB 170817A



We observed outside the jet of a classical sGRB

Pros:


- Can naturally explain the lower energetics
- Thermal emission could be from the GRB photosphere or the cocoon

Cons:

- Highly unlikely to observe the jet from the side due to relativistic beaming
- The on-axis  $E_{pk}$  would be on the high end of the observed GBM catalog distribution
- Expect bright afterglow in X-ray after  $\sim 1$  day



# GRB 170817A



We observed the less energetic region of a structure jet where the Lorentz factor decreases with viewing angle

Pros:


- Could produce arbitrary  $E_{pk}$  and  $E_{iso}$  values
- GW-EM delay is on the order of T90
- Thermal emission could be from the GRB photosphere or the cocoon

Cons:

- Not entirely clear how such wings are generated or what their Lorentz profiles look like
- On-axis  $E_{iso}$  would still need to be relatively low



# GRB 170817A



Hard emission from mildly-relativistic shock breakout and thermal emission from cocoon

Pros:

- Can naturally explain the lower energetics
- Could naturally explain both hard and thermal components




Cons:

- Cannot explain very high  $E_{pk}$  values
- Difficult to explain fast variability
- Should overproduce look alike sGRBs



# GRB 170817A

Alexander et al. 2018



We believe we observed GRB 170817 off-axis



- The off-axis jet is expected to be moving slower and therefore produce weaker gamma-ray emission
- The observed rise and peak of X-ray and radio emission favors the structured jet interpretation



Are there other Gamma-ray Bursts  
similar to GRB 170817A?




# GRB 150101B



- Very hard initial pulse with  $E_{pk} = 1280 \pm 590$  keV followed by a soft thermal tail with  $kT \sim 10$  keV
- Unlike GRB 170817, 150101B was not under luminous and can be modeled as an on-axis burst
- Suggests that the soft tail is common, but generally undetectable in more distant events
- Thermal tail can be explained as GRB photosphere, but degeneracy with the cocoon model still exists



# SIMILAR GRBS IN GBM DATA




Von Kienlin et al 2019

GRB 170817A-like hard spike followed by a softer thermal tail

- ~10 similar short GRBs found
- Most likely, all of these SGRBs are relatively nearby
- Longer softer bursts like GRB 170817A may be off-axis
- Shorter harder bursts, like GRB 150101B may be more on-axis
- More coincident SGRB/GW detections are needed to confirm!




# FERMI GAMMA-RAY SPACE TELESCOPE



12 NaI detectors  
(8keV—1MeV)



2 BGO detectors  
(200keV—40MeV)





# ALL SKY COVERAGE




GBM instantaneous field of view: ~70% of the sky  
~87% uptime (off during South Atlantic Anomaly)





# ALL SKY COVERAGE

GBM instantaneous field of view: ~70% of the sky  
~87% uptime (off during South Atlantic Anomaly)






# GBM TRIGGERS

6608 triggers in 10 years


2438 GRBs



1177 Solar Flares



280 Magnetars



905 TGFs




726 Others (pulsars and binaries)

1092 particles



# REAL TIME ALERTS

<https://fermi.gsfc.nasa.gov/ssc/data/access/gbm/>



GCN: The Gamma-ray Coordinates Network


Notices by Fermi-GBM:

FERMI_GBM_ALERT	~10s	triggered time, lightcurves
FERMI_GBM_FLT_POSITION	~30s	flight location, classification, lightcurves
FERMI_GBM_GND_POSITION	~45s	ground location, lightcurves, map
FERMI_GBM_FINAL_POSITION	minutes — hour	final position, lightcurves, map (healpix)
Circular	few hours	temporal and spectral analyses, or misclassification report




# GROUND SEARCH PIPELINES

- Continuous Time Tagged Events (CTTE) enabled 2012
    - 2 $\mu$ s, 128 energy channels
- 1.Untargeted search for subthreshold GRB candidate events
  - 2.Targeted search using input event time and optional skymap



Ideal Scenario	Bright GBM	Bright LIGO
GW150914 Scenario	Sub-threshold GBM	Bright LIGO
Typical more distant short GRB	Bright GBM	Sub-threshold LIGO
Both Sources Faint	Sub-threshold GBM	Sub-threshold LIGO







# UNTARGETED SEARCH

**Extends the onboard trigger algorithms, with improved background model.**

- Looks for signals in 2 NaI detectors with  $2.5\sigma$  and  $1.25\sigma$  excess above background.
- The 2 signal detectors must have valid geometry for a point source.
- **18 timescales: 64ms to 32s.**
  - Only candidates  $<2.8s$  are reported at the moment.
- **4 energy ranges** optimized for short GRBs.
  - 27—539 keV; 50—539 keV; 102—539 keV; 102—985 keV

- From April 2017 to now, **64/month**, excluding Oct/Nov 2017
- Found additional burst-like transients from magnetars and X-ray binaries
- **GRB170817A: could dim x0.5 and still recover by untargeted search.**




- 318 short, hard candidates found in 46 months in previous study → ~80 per year.





# UNTARGETED SEARCH

- GCN notice type Fermi-GBM SubThreshold now available.  
[https://gcn.gsfc.nasa.gov/fermi\\_gbm\\_subthreshold.html](https://gcn.gsfc.nasa.gov/fermi_gbm_subthreshold.html)
- Time delay for notice range from 0.5 to 6 hours, due to telemetry schedule.
- List of candidates from older data (2013 and on) are available.  
[http://gammaray.nsstc.nasa.gov/gbm/science/sgrb\\_search.html](http://gammaray.nsstc.nasa.gov/gbm/science/sgrb_search.html)
- Available with the GCN notice:
  - Localization FITS file
  - Contour sky map
  - Lightcurve



## GRB 170921C [Zhang et al. GCN 21919]


- Insight-HXMT  $12\sigma$  detection coincident with Fermi-GBM subthreshold transient 527647422.






# GROUND SEARCH PIPELINES

- Continuous Time Tagged Events (CTTE) enabled 2012
    - 2 $\mu$ s, 128 energy channels
- 1.Untargeted search for subthreshold GRB candidate events
- 2.Targeted search using input event time and optional skymap





Ideal Scenario	Bright GBM	Bright LIGO
GW150914 Scenario	Sub-threshold GBM	Bright LIGO
Typical more distant short GRB	Bright GBM	Sub-threshold LIGO
Both Sources Faint	Sub-threshold GBM	Sub-threshold LIGO





# TARGETED SEARCH

## Coherent search over GBM detectors



background-subtracted counts

product over independent observations (detectors/energy channels)

likelihood including signal model

$$P(d_i|H_1) = \prod_i \frac{1}{\sqrt{2\pi}\sigma_{d_i}} \exp\left(-\frac{(\tilde{d}_i - r_i s)^2}{2\sigma_{d_i}^2}\right)$$

likelihood from noise only

$$P(d_i|H_0) = \prod_i \frac{1}{\sqrt{2\pi}\sigma_{n_i}} \exp\left(-\frac{\tilde{d}_i^2}{2\sigma_{n_i}^2}\right)$$

response

source amplitude


- **Targeted** search in the Continuous Time Tagged Events (CTTE) data. (Blackburn et al. 2015, Goldstein et al. arXiv:1612:02395)
  - Looks for coherent signals in all detectors given an input time and optional skymap.
  - Calculate likelihood ratio of source and background.
  - Search +/- 30 seconds of input event time.
  - Sliding timescales from 0.256s to 8s (capable down to 0.064s) with a factor of 4 phase shift.
  - 3 source spectral templates using Band function: soft, normal, and hard.




# TARGETED SEARCH

- Testing with a control sample: 42 short GRBS detected by Swift BAT also in GBM FOV (2008 Aug 4 — 2017 Aug 4)
  - 31 detected by both instruments
  - 11 only by Swift
    - intrinsically dim and/or poor viewing geometry by GBM

Swift GRB did not trigger GBM



Kocevski et al. 2018





40/42 detected by the targeted search at  $>3\sigma$   
(likelihood ratio  $>9$ )



# TARGETED SEARCH

- Testing with a control sample: 42 short GRBS detected by Swift BAT also in GBM FOV (2008 Aug 4 — 2017 Aug 4)
  - 31 detected by both instruments
  - 11 only by Swift
    - intrinsically dim and/or poor viewing geometry by GBM

Swift GRB did not trigger GBM



GRB 170817 can dim by 60% and still discoverable by this search

-> increases the volume of the Universe in which GRB 170817 could be detected by factor of 5

Kocevski et al. 2018




# GBM-LIGO PARTNERSHIP




- GBM-LIGO MoU allows for a unique data sharing agreement
- GBM provides sub-threshold GRBs in low-latency for GW follow-up
- LIGO provides “sub-threshold” GW candidates below EM Follow-up threshold
- In low-latency for autonomous targeted searches with GBM
- GBM detections would provide increased confidence in weak GW detections, effectively increasing the volume of the Universe accessible to LIGO/Virgo




# GBM-LIGO PARTNERSHIP




Hanford



GBM + Hanford



Hanford + GBM + Livingston




- GW duty cycle ~70-75% (Abbott et al. 2018c)
  - 3 (2) GW detectors operating 34 – 42% (78 – 84% ) of the time
  - GBM will often constrain single interferometer localizations
- For GRB 170817A, GBM+HL map (~60 sq. deg) could have been produced ~1 hr after GW trigger



# INSTRUMENTS


## Current Instruments

- Konus-WIND (launched 1994)
- INTEGRAL (launched 2002)
- Swift (launched 2004)
- Fermi (launched 2008)
- MAXI (launched 2009)
- AstroSAT (launched 2015)
- CALET (launched 2015)
- HXMT (launched 2017)
- [IPN only missions - Messenger, Odyssey]



## Future Missions

- SVOM (launch 2021)
- ISS-TAO (pending down-select in 2019, launch 2022)
- Einstein Probe (launch 2023)
- TAP (pending US Decadal Survey, launch ~2028)
- AMEGO (pending US Decadal Survey, launch ~2028)
- Nimble (NASA SMEX concept, launch ~2025)
- THESEUS (pending down-select in 2021, launch 2032)
- Others?






# CUBESATS / SMALLSATS


## BurstCube

- 6U CubeSat with 4 CsI crystals + Si photomultipliers
- 70% of Fermi-GBM effective area at 100keV
- ISS deployment with 1 year mission lifetime
- Currently funded for development



## MoonBEAM

- 12U CubeSat with scintillating crystals + Si photomultipliers
- Cislunar orbit to provide improved localization via time of flight
- 1-2 year mission lifetime



## Others

- Constellations of 3U CubeSats: CAMELOT, HERMES





# SUMMARY

- GW170817 / GRB 170817A is one of the best observed transient and highlights the science impact of multimessenger observations.
- Many open questions remain, with increased GW interferometer sensitivity, there will be more joint detections with GBM, enabling deeper population studies of SGRBs:
  - Additional distance measures which yield source energetics
  - Constrain jet structure and opening angle distribution
  - Cocoon emission from SGRBs
  - Causes of precursor and extended emission
  - Rates of SGRBs in the universe with implications for source evolution
- Fermi GBM is currently the most prolific short GRB detector
  - Subthreshold searches are crucial to increasing GBM sensitivity and the detection horizon to weak events like GRB 170817A
- Looking forward to future multimessenger discoveries:
  - Neutron star — Blackhole merger, neutrinos, Fast Radio Bursts!?