

Impacts of Degradation on Water, Energy, and Carbon Cycling of the Amazon Tropical Forests

Marcos Longo¹, Sassan Saatchi^{2,3}, Michael Keller^{2,4,5}, Kevin Bowman², António Ferraz^{2,3}, Paul R. Moorcroft⁶, Douglas C Morton⁷, Damien Bonal⁸, Paulo Brando^{9,10,11}, Benoît Burban¹², Géraldine Derroire¹³, Maiza N dos-Santos⁵, Victoria Meyer², Scott Saleska¹⁴, Susan Trumbore¹⁵, Grégoire Vincent¹⁶

¹NASA Postdoctoral Program Fellow, Jet Propulsion Laboratory, California Institute of Technology, Pasadena CA, United States

²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, United States

³Institute of Environment and Sustainability, University of California, Los Angeles, CA, United States

⁴International Institute of Tropical Forestry, USDA Forest Service, Rio Piedras, Puerto Rico

⁵Embrapa Informática Agropecuária, Campinas, SP, Brazil

⁶Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, United States

⁷NASA Goddard Space Flight Center, Greenbelt, MD, United States

⁸Université de Lorraine, INRAE, AgroParisTech, UMR Silva, F-54000 Nancy, France

⁹Department of Earth System Science, University of California, Irvine, CA, United States

¹⁰Woods Hole Research Center, Woods Hole, MA, United States

¹¹Instituto de Pesquisa Ambiental da Amazônia, Brasília, DF, Brazil

¹²Institut National de Recherche en Agriculture, Alimentation et Environnement (INRAE), UMR 0745

EcoFoG, Campus Agronomique, Kourou 97379, France

¹³Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD),

UMR EcoFoG (Agroparistech, CNRS, INRAE, Université des Antilles, Université de Guyane), Campus

Agronomique, Kourou 97379, France

¹⁴University of Arizona, Tucson, AZ, United States

¹⁵Max-Planck-Institut für Biochemie, Jena, Germany

¹⁶AMAP, Univ Montpellier, IRD, CIRAD, CNRS, INRAE, Montpellier, 34000 France

Key Points:

- Airborne lidar can be used to inform degradation-driven changes in structure to vegetation models

Corresponding author: Marcos Longo, mdplongo@gmail.com

- 31 • Forest degradation typically depletes evapotranspiration and productivity and in-
32 creases flammability
- 33 • Extreme droughts reduce functional differences between degraded and intact trop-
34 ical forests

Abstract

Selective logging, fragmentation, and understory fires directly degrade forest structure and composition. However, studies addressing the effects of forest degradation on carbon, water, and energy cycles are scarce. Here, we integrate field observations and high-resolution remote sensing from airborne lidar to provide realistic initial conditions to the Ecosystem Demography Model (ED-2.2) and investigate how disturbances from forest degradation affect gross primary production (GPP), evapotranspiration (ET), and sensible heat flux (H). We used forest structural information retrieved from airborne lidar samples (13,500 ha) and calibrated with 817 inventory plots (0.25 ha) across precipitation and degradation gradients in the Eastern Amazon as initial conditions to ED-2.2 model. Our results show that the magnitude and seasonality of fluxes were modulated by changes in forest structure caused by degradation. During the dry season and under typical conditions, severely degraded forests (biomass loss $\geq 66\%$) experienced water-stress with declines in ET (up to 34%) and GPP (up to 35%), and increases of H (up to 43%) and daily mean ground temperatures (up to 6.5°C) relative to intact forests. In contrast, the relative impact of forest degradation on energy, water, and carbon cycles markedly diminishes under extreme, multi-year droughts, as a consequence of severe stress experienced by intact forests. Our results highlight that the water and energy cycles in the Amazon are not only driven by climate and deforestation, but also the past disturbance and changes of forest structure from degradation, suggesting a much broader influence of human land use activities on the tropical ecosystems.

Plain Language Summary

In the Amazon, timber extraction and forest fires ignited by people are the chief causes of damages that we call forest degradation. Degradation is as widespread as deforestation, and changes how forests behave. Degraded forests may pump less water to the atmosphere and absorb less carbon dioxide from the atmosphere. To understand the differences in behavior between degraded and intact forests, we used high-resolution scanning laser data collected from aircraft flights over regions in the Amazon where we knew if and when forests were degraded. Then, we provided these data to a computer program that calculates the exchange of water and carbon between the forest and the atmosphere. We found that, during the dry season, degraded forests are 6.5°C warmer, pump 1/3 less water (i.e., 400,000 L ha⁻¹ month⁻¹), absorb 1/3 less carbon (i.e., 1 tonC ha⁻¹ month⁻¹),

and show higher fire risk than intact forests. To our surprise, when the Amazon is hit by severe droughts, intact forests start to behave like degraded forests, because all forests run out of water and become hot. Our results are important because they show that forest degradation caused by people can have large impacts on dry-season climate and favor more fire, especially during typical, non-drought years.

1 Introduction

Tropical forests account for 25–40% of total carbon stocks in terrestrial ecosystems (Sabine et al., 2004; Meister et al., 2012), but their maintenance and functioning have been weakened by climate and land-use change. As a result, tropical forests may shift to net sources of carbon to the atmosphere, with residence time of carbon in forests declining by 50% (Davidson et al., 2012; Grace et al., 2014; Lewis et al., 2015; Erb et al., 2016). Land use and land cover changes contribute to nearly 15% of total annual carbon emissions (Harris et al., 2012; Friedlingstein et al., 2019). However, most studies assessing the effects of land use change on tropical forest stocks and fluxes have focused on the effects of deforestation (e.g., Harris et al., 2012; Achard et al., 2014). Logging, understory fires and forest fragmentation — collectively known as *forest degradation* (Hosonuma et al., 2012) — could play a comparable role in the forest’s energy, water, and carbon cycle and induce locally warmer and drier conditions that could be detrimental to their functioning (Grossiord et al., 2020; Sullivan et al., 2020), but these effects remain poorly quantified.

Significant fractions of the remaining tropical forests are located within 1 km from the forest’s edge (Haddad et al., 2015; Lewis et al., 2015) and thus are probably degraded (Asner et al., 2006; Morton et al., 2013; Pütz et al., 2014; Tyukavina et al., 2016; Potapov et al., 2017). The area impacted by forest degradation in the Amazon each year is highly uncertain, but likely comparable to deforestation (Asner et al., 2006; Morton et al., 2013; Tyukavina et al., 2017). Total carbon losses attributable to degradation may be similar or exceed deforestation-related losses in tropical forests (Berenguer et al., 2014; Pearson et al., 2017; Baccini et al., 2017; Aragão et al., 2018; Erb et al., 2018), and degradation may even dominate the carbon losses in indigenous lands and protected areas (Walker et al., 2020). At the local scale, carbon stocks in degraded forests are extremely variable. Lightly disturbed forests (e.g., reduced-impact logging) store as much carbon as intact forests, while forests impacted by severe or multiple disturbances may lose a significant

fraction or nearly all of their original carbon stocks (Berenguer et al., 2014; Alamgir et al., 2016; Longo et al., 2016; Rappaport et al., 2018; Ferraz et al., 2018). Transitions between lightly and heavily degraded forests may be non-linear and abrupt (Brando et al., 2014). Unquestionably, estimates of fluxes from forest degradation and regeneration are more uncertain than emissions from deforestation (Aragão et al., 2014; Morton, 2016; Bustamante et al., 2016), because their impacts on forests are more subtle than deforestation and thus more difficult to detect and quantify with traditional remote sensing techniques.

Selective logging and fires also modify the forest structure, composition and functioning. For example, selective logging in the tropics generally targets large trees (diameter at breast height, $\text{DBH} \geq 40\text{--}60\text{ cm}$) from a few marketable species (e.g., Feldpausch et al., 2005; Blanc et al., 2009; Pinagé et al., 2019), but the other logging structures such as skid trails and log decks kill or damage mostly small trees ($\text{DBH} < 20\text{ cm}$) (Feldpausch et al., 2005). Likewise, fire mortality decreases with tree size and the bark thickness (e.g., Brando et al., 2012; Pellegrini et al., 2016), although areas disturbed by recurrent fires also show significant losses of large trees (Barlow et al., 2003; Martins et al., 2012; Brando, Silvério, et al., 2019; Silvério et al., 2019). Consequently, degradation creates more open canopies and thinner understory (e.g., d'Oliveira et al., 2012; Pinagé et al., 2019; Silvério et al., 2019) and increased abundance of grasses and fast-growing, low wood-density tree species (Barlow et al., 2016; Both et al., 2019; Brando, Silvério, et al., 2019).

Previous studies indicate an increase in dry-season length in parts of the Amazon where both deforestation and forest degradation are pervasive (e.g., Fu et al., 2013; Sena et al., 2018), and that the onset of the wet season is modulated by forest transpiration (J. S. Wright et al., 2017). Temperature and vapor pressure deficit (VPD), important drivers of evapotranspiration (ET), were found by Kapos (1989) to be significantly higher near forest edges. Likewise, Jucker et al. (2018) installed a network of micrometeorological measurements across a study area in Sabah, Malaysia, that included intact forests, a broad range of degraded forests and oil-palm plantations, and found that forest structure, along with topographic features, explained most of the variance in understory temperature. Yet, only a few studies on experimental sites quantified the magnitude, seasonality, and interannual variability of water, and energy cycles in degraded forests. For example, S. D. Miller et al. (2011) analyzed the impact of reduced-impact, low-intensity selective logging in the Amazon using eddy covariance towers and found only minor im-

pacts of logging on sensible and latent heat fluxes. Recently, Brando, Silvério, et al. (2019) compared eddy covariance data from two towers at an experimental fire site in the Amazon forest, and found declining differences in gross primary productivity and small differences in evapotranspiration between the control and burned area between 4 and 8 years after the last burn.

Field inventory plots are fundamental to sample the structure and species composition of tropical forests, but they also have important limitations to characterize the heterogeneity of degraded landscapes. First, the number of plots required to characterize stands increase with heterogeneity, often reaching impractical numbers (Marvin et al., 2014). In addition, most tropical forest degradation occurs in private landholdings and privately managed logging concessions, where limited access by researchers may create sampling bias towards well-managed areas, which generally experience less intensive degradation. However, airborne laser scanning (airborne lidar) can circumvent these limitations over large areas with sub-meter resolution. Airborne lidar data have been used successfully to quantify structural characteristics of the canopy such as height and leaf area distribution (Hunter et al., 2013; Vincent et al., 2017; Shao et al., 2019). Moreover, these data have also been used to quantify changes in canopy structure and carbon stocks at local to regional scale that experienced multiple levels of degradation (e.g., Asner et al., 2010; Longo et al., 2016; Ferraz et al., 2018; Meyer et al., 2019).

Numerical models can be used to understand the links between changes in forest structure, light and water availability for different local plant communities, and the overall impact on energy, water, and carbon fluxes between forests and the atmosphere. In the past, *big-leaf* models have been modified to account for the long-term impacts of selectively logged tropical forests on the carbon cycle of tropical forests (e.g., Huang et al., 2008; Huang & Asner, 2010). However, big-leaf models generally do not represent the mechanisms that control access and availability of light and water in complex and heterogeneous forest structures (D. Purves & Pacala, 2008; Fisher et al., 2018) (but see Braghieri et al., 2019). Individual-based models can represent the changes in the population structure and micro-environments due to degradation (R. Fischer et al., 2016; Maréchaux & Chave, 2017), but the complexity and computational burden of these simulations often limit their application to single sites. Cohort-based models, such as the Ecosystem Demography (ED-2.2) model (Medvigy et al., 2009; Longo, Knox, Medvigy, et al., 2019), strike a balance between these end-members because they can efficiently represent the

horizontal and vertical heterogeneity of forests. However, to represent the impact of heterogeneity in the energy, water, and carbon cycles, it is critical that these models are informed with realistic initial conditions that capture the landscape variability and they accurately represent the complex interactions between climate and the micro-environment variability. Previous studies using a variety of cohort-based models have demonstrated that cohort-based models can realistically reproduce the micro-environment heterogeneity and the long-term dynamics of ecosystems, compared to both individual-based models (Moorcroft et al., 2001; Strigul et al., 2008) and observations (D. W. Purves et al., 2008; Longo, Knox, Levine, et al., 2019; Koven et al., 2019).

In this study, we use airborne lidar data to quantify forest structure variability across the Amazon in order to provide critical initial conditions for ecosystem demography models. We also investigate the role of forest degradation on the Amazon forest productivity, flammability, as well as the degradation impacts on the water and energy cycles. Specifically, we seek to answer the following questions:

1. What are the relationships between degradation metrics (e.g. biomass loss) and changes in carbon, water, and energy fluxes, and how does it vary across seasons and regions with different rainfall regimes?
2. How do droughts affect the relationships between degradation and ecosystem functioning?
3. Does forest degradation make Amazon forests more susceptible to fires? If so, which parts of the Amazon experience the largest flammability response to degradation?

To this end, we integrate field inventory plots with high-resolution airborne lidar data over five study regions in the Eastern Amazon along a precipitation gradient and with a broad range of anthropogenic disturbance histories, to provide initial conditions to ED-2.2 that realistically represent the structural diversity of degraded forests. While limited to specific regions in the Amazon where detailed degradation information exists, our goal is to provide a framework that can be extended to larger scales, including biome- and pantropical scales.

2 Materials and Methods

2.1 Study regions

We selected five study regions across a gradient of disturbance and climate conditions where ground and airborne lidar are available to study the forest function (Figure 1; Table 1). Three of these sites include eddy covariance tower measurement of energy, water, and carbon dioxide fluxes for comparison with the model simulations, and have been the focus of several ecological studies in the past. Additional details on the disturbance history of each region are available at Text S1.

1. *Paracou, French Guiana (GYF)* is a field station where a logging experiment was conducted between 1987 and 1988 that includes intact forest controls and three selective logging treatments: timber extraction using conventional logging techniques, timber extraction and canopy thinning, and timber and fuelwood extraction followed by canopy thinning (Gourlet-Fleury et al., 2004). The eddy covariance tower at the site is located in the undisturbed forest and has been operational since 2004 (Guyafflux; Bonal et al., 2008).
2. *Belterra, Brazil (BTE)*. Over the past 100 years, this region experienced cycles of economic growth and recession that created a complex landscapes dominated by deforestation, degradation and second-growth. The Tapajós National Forest is this region, and has areas of intact forests and selectively logged forests using reduced-impact techniques (VanWey et al., 2007; Pyle et al., 2008; Lei et al., 2018). An eddy covariance tower known as Km 67 overlaps with one of the surveyed sites and has data for 2001–2005, and 2008–2011 (Hayek et al., 2018).
3. The *Paragominas, Brazil (PRG)* region used to be within the largest timber production area in Brazil and has undergone selective logging since the 1970s (Veríssimo et al., 1992). Since the 1990s, the economy has shifted towards agriculture, introducing large-scale deforestation such that nearly half of the original forest cover has been lost, and most of the remaining areas have been logged (Pinto et al., 2009).
4. *Feliz Natal, Brazil (FZN)* is located at the southern fringe of the Amazon in a mosaic landscape of soybean fields, grazing lands, and logged forests. This region regularly experiences severe dry seasons and frequent understory fires (Morton et al., 2013; Rappaport et al., 2018).




Figure 1. Location of the five study regions within the Amazon biome region, along with land classification as of 2013. Intact forest and intact forest loss were obtained from Potapov et al. (2017); open and deforested areas were obtained from PRODES-INPE (2018) (Brazil) and areas with tree cover below 20% according to Hansen et al. (2013) (other countries); wetlands and water bodies in the Amazon River Basin were from Hess et al. (2015) and savannas and mangroves were obtained from Olson et al. (2001).

5. *Tanguro, Brazil (TAN)* is located in an experimental fire study area within a larger landscape covered by intact forests and forests that were disturbed with low-intensity understory fires (one, three, and six times) between 2004 and 2010 (Balch et al., 2008; Brando et al., 2014). The surveyed region also includes two eddy covariance towers that have been operating since 2014 both at the intact and burned forests (Brando, Silvério, et al., 2019).

These five study regions were sampled at multiple sites by small-footprint, multiple-return airborne lidar. The lidar data provided both the terrain elevation at high spatial resolution (1-m) and detailed information about the vertical structure of forests from a uniform point cloud density to meet a minimum return density of 4 returns per m^2 over 99.5% of the area (Leitold et al., 2015). Living trees of diameter at breast height DBH $\geq$ 10 cm were either botanically identified (experimental plots in GYF) or identified from field characteristics by local parataxonomists. To characterize the disturbance history,

Table 1. Overview of the study regions, including mean annual precipitation (MAP) and dry-season length (DSL).

Region (Code)	Coordinates	MAP ^a [mm]	DSL ^b [mo]	Lidar [ha]	Inventory [ha]	Disturbances ^c
Paracou (GYF)	5.28°N; 52.91°W	3040	2(0)	963	79.8	INT, CL1, LTH
Belterra (BTE)	3.09°S; 54.95°W	1890	5(1)	4057	16.7	INT, RIL, BN1, BN2, BN3
Paragominas (PRG)	3.15°S; 47.61°W	1850	6(2)	3217	35.6	INT, RIL, CL1, BN1, LB1, BN2, BN3
Feliz Natal (FZN)	12.14°S; 54.68°W	1940	5(4)	4210	14.0	INT, CL1, CL2, BN1, LB1, BN2, BN3
Tanguro (TAN)	13.08°S; 52.41°W	1800	5(4)	1006	22.9	INT, BN1, BN3, BN6

^a Source for mean annual precipitation (MAP) data: GYF – Gourlet-Fleury et al. (2004); other regions – nearest site available at INMET (2019).

^b Dry-season length (DSL): number of months with precipitation below 100 mm; numbers in parentheses indicate number of severely dry months (precipitation below 30 mm).

^c Disturbance history classes: INT – intact; RIL – reduced-impact logging; CL x – conventional logging (x times); LTH – conventional logging and thinning; LB1 – conventional logging and burned (once); BN x – burned x times.

we used either published information from the experimental regions GYF (Gourlet-Fleury et al., 2004; Bonal et al., 2008; Wagner et al., 2013) and TAN (Brando et al., 2012, 2014), or the disturbance history analysis from (Longo et al., 2016), which was based on a visual interpretation of the Normalized Burn Ratio (NBR) of cloud-free Landsat images since 1984, and complemented with information from logging companies for the reduced-impact logging sites (e.g., Pinagé et al., 2019). Details on site-specific data used in this study are available in Text S2 and previous work (Longo et al., 2016; Vincent et al., 2017; Brando, Silvério, et al., 2019), and were obtained through the Paracou Experimental Station and the Sustainable Landscapes Brazil data servers (Paracou Portal, 2016; Sustainable Landscapes Brazil, 2019; dos-Santos et al., 2019).

2.2 Overview of the modeling framework

In this study, we used the Ecosystem Demography model, version 2.2 (ED-2.2) (Moorcroft et al., 2001; Medvigy et al., 2009; Longo, Knox, Medvigy, et al., 2019) to simulate the impacts of forest structure on energy, water, and carbon cycles. For any point of interest, the ED-2.2 model simulates the forest structure and functional diversity across a landscape, and simulates the energy, water, and carbon budgets for multiple canopy environments, which represent the forest heterogeneity (Longo, Knox, Medvigy, et al., 2019). ED-2.2 has been successfully evaluated and used in both short-term and long-term studies in the Amazon forest (Powell et al., 2013; Zhang et al., 2015; Levine et al., 2016; Longo, Knox, Levine, et al., 2019). In ED-2.2, the horizontal and vertical heterogeneities of forests are represented through a hierarchical structure. Each area with the same climate (e.g.,

footprint of an eddy covariance tower or a grid cell in a gridded meteorological driver) is called a *polygon*. Each polygon is subdivided into *patches*, which represent collections of forest gaps within a polygon that share a similar age since last disturbance and same disturbance type (although not necessarily contiguous in space). Patches are further subdivided into *cohorts*, which are collections of individual plants that have similar size and similar functional group. Importantly, because ED-2.2 incorporates the horizontal heterogeneity of the plant community structure and composition, the model can efficiently incorporate and simulate the dynamics of degraded forests.

Most of the ED-2.2 modules used in this study have been previously described in Longo, Knox, Medvigy, et al. (2019). The main changes used in this study include (1) a modified height-diameter allometry based on the Jucker et al. (2017) approach and locally collected field data that can be used consistently by the initialization and model; (2) an improved allocation to living and structural tissues, which is now based on more recent allometric equations (Chave et al., 2014; Falster et al., 2016) and datasets (Falster et al., 2015); (3) a revised photosynthesis solver, which now accounts for the maximum electron transport ratio and the maximum triose-phosphate utilization (von Caemmerer, 2000; Oleson et al., 2013; Lombardozzi et al., 2018); (4) updated values of traits that are used to define trade-offs in tropical plant functional types in ED-2.2 (wood density and leaf turnover rate), and updated the trade-off relationships of traits that directly or indirectly influence gross primary productivity and light- and water-use efficiency (specific leaf area and leaf carbon:nitrogen ratio, maximum carboxylation rate, maximum electron transport ratio and maximum triose-phosphate utilization), using multiple studies and trait databases, including GLOPNET, TRY, and NGEE-Tropics (I. J. Wright et al., 2004; Santiago & Wright, 2007; Chave et al., 2009; Kattge et al., 2009, 2011, 2020; Baraloto et al., 2010; Powers & Tiffin, 2010; Gu et al., 2016; Bahar et al., 2017; Norby et al., 2017). These changes are described in Text S3. Moreover, we used an approach developed by X. Xu (unpublished) and based on Lloyd et al. (2010) to account for light-dependent plasticity of three leaf traits (specific leaf area, leaf turnover rate, and carboxylation capacity), and calibrated using existing data (Lloyd et al., 2010; Russo & Kitajima, 2016; Keenan & Niinemets, 2016).

To obtain initial conditions for ED-2.2 from airborne lidar, we devised a multi-step approach that links airborne lidar data with ecosystem properties (Figure 2). Here we provide a summary of the initialization procedure; the technical details of this approach

are described in Text S4. For step 1, we split all collected point cloud data into 50×50 m columns, simulated waveforms from the discrete returns (Blair & Hofton, 1999; Popescu et al., 2011; Hancock et al., 2019) to obtain unscaled leaf area density profiles based on the vertical distribution of returns (e.g., MacArthur & Horn, 1969; Ni-Meister et al., 2001; Stark et al., 2012; Antonarakis et al., 2014; Tang & Dubayah, 2017), and assigned the relative proportion of each plant functional type provided by one of the 769 training plots that had the most similar vertical structure; the similarity was based on the profile comparison that yielded the smallest Kolmogorov-Smirnov statistic. The vertical profile was split into cohort layers centered around local maxima or saddle points, using a modified procedure based on function `peaks` (package `RSEIS`, Lees, 2017) of the R statistical software (R Core Team, 2019). For step 2, we used a collection of 817 forest inventory plots (0.16–0.26 ha) that were also surveyed by airborne lidar, which included plots from all study regions as well additional sites available from Sustainable Landscapes Brazil (SLB) and used in a previous study (ancillary SLB sites, Figure 1; Longo et al., 2016); we developed statistical models based on subset selection of regression (A. J. Miller, 1984) and heteroskedastic distribution of residuals (Mascaro et al., 2011) to estimate plot-level properties (aboveground biomass, basal area, stem number density, leaf area index) from point cloud metrics and field estimates, following the approach by Longo et al. (2016). For step 3, we sought to obtain a plot-specific scaling factor to the leaf area density profile that produced the best agreement between the four estimated plot-level properties from step 1 and the plot-level properties obtained by integrating the vertical distribution from step 2, by minimizing the sum of relative square differences of the four properties. For step 4, we analyze the scaling factor distribution for all plots for which we could test the approach, and define a unique and global scaling factor, based on the median scaling factor, that is used to correct all predicted profiles.

Once we obtained the initial conditions for each 50×50 m column, we grouped individual columns based the disturbance history (degradation level) and the study region (Table 1). We used the following broad categories for disturbance history: intact (INT), reduced-impact logging (RIL), conventional logging (CL x , where x is the number of logging disturbances), conventional logging and thinning (LTH), logged and burned once (LB1) and burned (BN x , where x is the number of burns). Importantly, we did not perform any averaging or sampling of the individual columns before providing them to ED-




Figure 2. Schematic representation of the method to obtain initial conditions for ED-2 from airborne lidar. Each light box represents one step in the procedure. The results of each step are highlighted with a red border. Dark blue arrows are stages that require individual-based allometric equations, and light blue arrows are stages that require a light extinction model.

2.2; instead, we provided all columns to the model, so the initial conditions characterize the observed distribution of forest structures that exist within each group.

2.3 Assessment of the modeling framework

We evaluated three characteristics to assess the ability of model framework to represent the forest structure heterogeneity caused by degradation, and to represent components of the energy, water, and carbon cycle. First, we quantified the ability of the airborne lidar initialization to capture the differences in forest structure caused by degradation. Second, we assessed whether the model can realistically represent fluxes and storage of water, energy and carbon across different regions. Third, we compared the model sensitivity to degradation-driven effects on fluxes and storage with independent observations.

To evaluate the airborne lidar initialization, we used a cross-validation approach in which we replicated the procedure described above (Section 2.2) 2000 times, using a

hierarchical bootstrap approach. We first sampled regions (with replacement), to ensure that some regions would be entirely excluded from the replicate, then we sampled plots (also with replacement), to ensure that the replicate had the same number of plots as the original training data set. We then predicted the structure of all plots in the excluded regions, using iterations that did not have any plot in the training data set; to make this number consistent across regions, we used the smallest number of iterations that met this criterion across all regions ($n = 612$). Finally, for each region, we compared the average forest structure from all cross-validation replicates that excluded the region from the training stage. Because estimates of forest properties have larger uncertainties in smaller plots (Chave et al., 2004; Meyer et al., 2013; Mauya et al., 2015), we only evaluated the method when a disturbance class within a region had at least 20 plots.

To verify the model’s ability to realistically represent the regional variability of fluxes and storage, we carried out ED-2.2 simulations initialized with airborne lidar for the intact forests regions where eddy covariance tower and forest inventory plots co-located with airborne lidar were available (GYF and BTE). Region TAN had two eddy-covariance towers, one within the footprint of the burned forests and a second in intact forest (Brando, Silvério, et al., 2019), which allowed us to contrast the model’s predicted impacts of degradation on fluxes and biophysical properties with the pair of tower measurements.

2.4 Model configuration and analyses

Our main focus is to understand the role of degradation-driven changes in forest structure in altering both the state and the fluxes of energy, water, and carbon, both under typical and extreme climate. To account for regional differences in climate and to sample a broad range of interannual variability, we used time series of meteorological drivers pooled from gridded reanalyses (one set of time series per region). For most meteorological variables required by ED-2.2 (pressure, temperature, humidity, incoming short-wave and longwave radiation, and winds), we used $0.625^\circ \times 0.5^\circ$, hourly averages (1980–2016) from the version 2 of the Modern-Era Retrospective Analysis for Research and Applications (MERRA-2, Gelaro et al., 2017). MERRA-2 precipitation is known to have significant negative biases in the tropics (Beck et al., 2019); therefore we used the $0.1^\circ \times 0.1^\circ$, 3-hourly precipitation rates from the version 2 of the Multi-Source Weighted Ensemble Precipitation product (MSWEP-2, Beck et al., 2019). To ensure that the only difference between simulations in the same study region was the distribution of forest structures,

we imposed the same edaphic conditions: free-drainage soils with 8 m deep, and nearly equal fractions of sand (32%), silt (34%), and clay (34%). To avoid confounding effects from post-disturbance mortality and recovery, all simulations were carried out without enabling dynamic vegetation, such that the differences in forest structure would remain the same for the entire time series, and all differences between simulations in the same region could be attributable to well-characterized differences in forest structure. However, disabling dynamic vegetation also precluded us from investigating the effects of climate-driven changes in the canopy structure on the energy, water, and carbon cycle, and thus potentially increasing biases in our estimates of fluxes following extreme events such as droughts.

To investigate the role of degradation on fire risk, we built on the original fire model from ED-1 (Moorcroft et al., 2001) to determine when fire-prone conditions would occur in each patch. The flammable area α_F (% yr⁻¹) is calculated from the fire disturbance rate λ_F (yr⁻¹):

$$\alpha_F = 100 [1 - \exp(-\lambda_F \Delta t)], \quad (1)$$

$$\lambda_F = \begin{cases} I C_{\text{Fuel}} & , \text{ if } \left[\frac{1}{|z_F|} \int_{z_F}^0 \vartheta(z) dz \right] < (1-f) \vartheta_{\text{WP}} + f \vartheta_{\text{Fc}} \\ 0 & , \text{ otherwise} \end{cases} . \quad (2)$$

where $\Delta t = 1$ yr; $I = 0.5 \text{ m}^2 \text{ kgC yr}^{-1}$ is a fire intensity parameter; $z_F = 30$ cm is the depth of the soil layer used to estimate dryness; ϑ (m³ m⁻³) is the soil moisture; ϑ_{WP} is the permanent wilting point and ϑ_{Fc} is the field capacity, both defined as in Longo, Knox, Medvigy, et al. (2019); and $f = 0.02$ is a phenomenological parameter that defines dry conditions. The values of I and f were selected based on the results from a previous model evaluation using ED-2.2 (Longo, Knox, Levine, et al., 2019). Because understory fires are the dominant type of fire in the Amazon (A. Alencar et al., 2006; Morton et al., 2013), we considered fuels to be comprised by above-ground litter, above-ground coarse woody debris, and above-ground biomass from grasses and seedlings (trees with height < 2 m); canopy trees were not considered to be fuels. The fire parameterization, although simple, has been previously demonstrated to capture the general features of fire regime across tropical South America (Longo, Knox, Levine, et al., 2019).

3 Results

3.1 Evaluation of the model initialization and simulated seasonal dynamics

The ED-2.2 model initialization approach from airborne lidar (Figure 3) captured the main differences in forest structure and composition, both across study regions and along degradation gradients. To illustrate the initialization, we focus on the basal area distribution obtained from cross-validation at disturbance histories within study regions that had at least 20 plots (Figure 3). At sites GYF, PRG, and TAN, the airborne lidar initialization predicted the total basal area with absolute biases ranging from 3% (GYF) to 13% (TAN), and root mean square error of the order of 18–27% (Figures 3c, 3f and 3i). The largest absolute discrepancies occurred for intermediate-sized trees ($20 \leq \text{DBH} < 40 \text{ cm}$) at GYF and PRG, where the airborne lidar initialization underestimated basal area by 2.9 and $4.3 \text{ cm}^2 \text{ m}^{-2}$, respectively (Figures 3c and 2f). The largest overestimation of airborne lidar was observed among larger trees ($60 \leq \text{DBH} < 100 \text{ cm}$) in intact forests at GYF ($2.4 \text{ cm}^2 \text{ m}^{-2}$; Figure 3c). The size distribution of most degraded forests were well characterized (Figures 3a-b, 3d-e and 3g); the largest deviations from inventory were observed in logged and burned forests in PRG, where airborne lidar underestimated total basal area by $3.0 \text{ cm}^2 \text{ m}^{-2}$ (Figure 3d). Likewise, the initialization algorithm represented the higher relative abundance of early successional plants in the most degraded sites, and the dominance of mid- and late-successional plants at intact forests at GYF and PRG (Figure S1), and realistically represented the leaf area distribution across regions and degradation levels (Figure S2).

ED-2.2 simulations using forest inventory and airborne lidar as initial conditions were compared with eddy covariance tower estimates of all sites (Figures 4 and S4-S9, and Table S1). Gross primary productivity (GPP) generally showed small biases relative to tower estimates (-0.046 to $+0.394 \text{ kgC m}^{-2} \text{ yr}^{-1}$), and relatively small errors (less than observed variability) at all sites, regardless of the initial conditions (Figure 4; Table S1). While the GPP seasonality was correctly represented at GYF, the model did not capture the late wet-season decrease and early dry-season increase of GPP at BTE, and it showed a delayed dry-season decline GPP at TAN compared to tower estimates (Figure S4). Net ecosystem productivity (NEP), on the other hand, showed significant biases, large errors, and relatively small correlation with tower estimates (Figure 4; Ta-




Figure 3. Assessment of basal area distribution as a function of diameter at breast height (DBH) for different study regions and degradation levels. Grey points are obtained from forest inventory plots, and blue points are obtained from the airborne lidar initialization (Figure 2) using a 612-fold regional cross-validation (i.e. excluding all plots from region in the calibration stage). Bands around points correspond to the standard deviation either across all plots in the same category (inventory) or across all plots and replicates (lidar). Sites: GYF – Paracou, PRG – Paragominas, FZN – Feliz Natal, TAN – Tanguaro. Disturbance classes: BNx – Burned twice or more, CL1 – conventional logging (once), LB1 – logged and burned once, LTH – logged and thinned, RIL – reduced-impact logging, INT – intact. Additional comparisons are shown in the Supporting Information: basal area as functions of plant functional type (Figure S1); leaf area index profiles as functions of height (Figure S2); comparisons for Belterra (BTE-RIL) (Figure S3).

ble S1), which were driven by excessive seasonality of heterotrophic respiration (Figure S5). Because the initial carbon stocks in necromass pools are uncertain, and the results on magnitude and seasonality of ecosystem respiration (and consequently NEP) are inconsistent with tower estimates, we will not discuss the simulation results in terms of respiration and NEP.

Water fluxes also showed small biases relative to the observed variability at GYF, TNF and TAN (Burned), regardless of the initialization (-0.01 to $+0.54$ mm day $^{-1}$; Figures 4a and 4c; Table S1); biases at TAN (Intact) were larger (0.69 – 0.82 mm day $^{-1}$). With the exception of TAN (Burned), the correlation between ED-2.2 and tower was high at daily averages (Figures 4b and 4d; Table S1). At TAN (Burned), the poorer agreement with tower estimates was caused by the model predicting a similar seasonality of water flux at both control and burned forests, whereas towers suggest an increase in water flux during the earlier part of the dry season (Figure S6). ED-2.2 predictions of sensible heat flux had high correlation with observations at all sites (Figures 4b and 4d; Table S1), although sensible heat flux shows significant biases at BTE, and dampened seasonality at GYF and TAN (Burned) (Figures 4a and 4c; Table S1; Figure S6). Outgoing shortwave radiation correctly captured the seasonality at the wettest sites, but it did not capture the sharp dry-season increase at TAN (Figure S8), which may be associated with dry-season leaf senescence and shedding that was likely underestimated by ED-2.2. In addition, ED-2.2 simulations overestimated outgoing longwave radiation at all sites except at TAN (Burned) using inventory initialization (Figure S9). Nonetheless, the seasonality and the intra-seasonal variation of outgoing longwave radiation were correctly captured by ED-2.2, resulting in generally high correlation and small standard deviation of residuals at most sites (Figure 4; Table S1).

3.2 Degradation effects on seasonality of fluxes

From ED-2.2, we found that forest degradation can have substantial impacts on the ecosystem function such as evapotranspiration (ET) or ground temperature in severely or recently degraded forests, and in parts of the Amazon with a longer dry season. At GYF, the airborne lidar survey sampled only intact forests and areas that were logged 25 years prior to the data acquisition: consequently, the average water vapor flux and ground temperature were nearly indistinguishable across degraded and intact forests (Figures 5a, S10a). At the equatorial sites, degradation effects were small during the wet sea-




Figure 4. Summary of ED-2.2 model assessment using eddy covariance towers as benchmarks, using simulations initialized with forest inventory and airborne lidar. (a,c) Bias-variance diagram and (b,d) Taylor diagram of multiple daily-averaged fluxes of carbon, energy, and water for Paracou (GYF), Belterra (BTE) and Tanguro (TAN, control and burned), for simulations initialized with (a,b) forest inventory plots and (c,d) airborne lidar. In the bias-variance diagram, bias (x axis), standard deviation of residuals (y axis) and root mean square error (concentric arcs) are normalized by the standard deviation of observations, as is the standard deviation of models in the Taylor diagram. In both diagrams, $\odot$ corresponds to the perfect model prediction. In all plots, we only compare daily averages of days with no measurement gaps. Comparisons of the seasonal cycle for all variables included in the diagrams are available at Figures S4-S9.

son but showed marked reduction in ET (2.1–6.7% in BTE and 4.3–31.8% in PRG) and increase in daytime temperature (0.4–0.9°C in BTE and 1.0–6.0°C in PRG) during the dry season, with the largest changes relative to intact forests found at burned areas (Figures 5b, 5c, S10b,c). At the southern (driest) sites, the seasonal changes were even more pronounced: at both FZN and TAN, ET decreased by 21–25% early in the dry season (Jun) at the most severely burned forests, whereas ET in intact forests peaked in the middle of the dry season (Jul–Aug; Figures 5d and 5e). Similarly, burned forests were warmer year-round than intact forests at the southern sites (FZN and TAN), with minimum warming during the wet season (Dec–Mar; 0.5–0.8°C), and maximum warming occurring at the peak of the dry season (Jul–Aug; 1.0–6.5°C; Figures S10d and S10e).

Importantly, the ED-2.2 results in Figures 5 and S10 emerge from the different distribution of forest structures associated with degradation histories. ED-2.2 accounts for the diversity of forest structures within each disturbance history by means of patches; each patch represents a different forest structure found within any disturbance regime, and patch area is proportional to the probability of finding such forest structure (Longo, Knox, Medvigy, et al., 2019). For example, the ground temperature is consistently warmer at the low biomass patches, but the differences between the lowest and highest patch temperatures are as low as 1°C at GYF (Figure 6a) and less than 4°C during the wet season even at the southern regions (Figures 6d and 6e). In contrast, differences along biomass gradients exceed 9°C during the dry season at all regions except GYF (Figure 6).

Likewise, when all simulated patches are considered, we observe strong coherence between biomass and gross primary productivity (GPP) across all regions and throughout the year (Figures 7 and S11). However, the effect of local communities on GPP is seasonal: differences in typical GPP between low-biomass and high-biomass patches do not exceed $1.1 \text{ kgC m}^{-2} \text{ yr}^{-1}$ during the wettest months (Figures 7a–7c), whereas the range of GPP reaches $0.7 \text{ kgC m}^{-2} \text{ yr}^{-1}$ at the short dry-season at GYF and exceeds $2.0 \text{ kgC m}^{-2} \text{ yr}^{-1}$ during the dry season at the most degraded and driest sites (Figures 7e and 7f). Similar effects were observed in evapotranspiration, where differences along biomass are the strongest during the dry season (Figure S12).




Figure 5. Monthly mean evapotranspiration (ET) as a function of region and degradation. Monthly averages correspond to the 1980–2016 period, simulated by ED-2.2 for (a) Paracou (GYF), (b) Belterra (BTE), (c) Paragominas (PRG), (d) Feliz Natal (FZN), and (e) Tanguro (TAN), aggregated by degradation history within each region (lines). Grey rectangles in the background correspond to the average dry season.

3.3 Degradation impacts on forest flammability

The impact of forest degradation on ecosystem functioning showed important year-to-year variability, and differences between intact and degraded forests were generally




Figure 6. Monthly mean daytime ground temperature as a function of region and local (patch) aboveground biomass. Monthly averages correspond to the 1980–2016 period, simulated by ED-2.2 for (a) Paracou (GYF), (b) Belterra (BTE), (c) Paragominas (PRG), (d) Feliz Natal (FZN), and (e) Tanguro (TAN), and the y axis corresponds to the aboveground biomass for each patch, linearly interpolated for visualization. White areas are outside the range of biomass of each region and thus excluded.

490 larger during typical years than during extreme droughts. For this section, we calculate
 491 the monthly water deficit based on the difference between potential evapotranspiration




Figure 7. Variability of gross primary productivity (GPP) as a function of local (patch) aboveground biomass (AGB). Scatter plot of AGB (x axis) and GPP (y axis) at sites (a,d) Paracou (GYF), (b,e) Paragominas (PRG), (c,f) Feliz Natal (FZN), for (a-c) the peak of wet season — May (GYF), March (PRG), and February (FZN) — and (d-f) peak of dry season — October (GYF and PRG), and August (FZN). Each point represents the 1980–2016 average GPP of each patch solved by ED-2.2; point shapes correspond to the disturbance history, and point colors represent the time between the last disturbance (undetermined for intact forests) and lidar data acquisition. Curves correspond to non-linear least squares fits of the most parsimonious function, defined from Bayesian Information Criterion (Schwarz, 1978), between shifted exponential or shifted Weibull functions. Only fits that produced $R^2_{adj} > 0.5$ were included.

(calculated following Priestley & Taylor, 1972) and rainfall, and relate the 12-month running averages of multiple response variables with the maximum cumulative water deficit over the previous 12 months, and define drought length as the number of consecutive months in water deficit exceeds 20 mm. Using region PRG as an example, as the region has the broadest range of recent disturbances and maximum cumulative water deficit, we found that, during typical rainfall periods, evapotranspiration in logged forests and burned forests were 3–6% and 11–22% lower than intact forests, respectively (Figure 8a); this difference was significantly reduced or even reversed during severe droughts, when evapotranspiration of degraded forests were up to 4% higher than in intact forests (Figure 8a). De-

501 graded forests have a lower proportion of shade-tolerant, late-successional trees, and typ-
 502 ical stomatal conductance is higher by 19–34% in burned forests and by 5–13% in logged
 503 forests (Figure 8b). This result indicates that the reduced typical evapotranspiration re-
 504 sults from degraded forests having lower leaf area index relative to intact forests, as lo-
 505 cal leaf area index is related to local aboveground biomass (Figure S13). In addition, ex-
 506 treme droughts did not substantially reduce the differences in stomatal conductance be-
 507 tween degraded and intact forests (Figure 8b). While evapotranspiration was generally
 508 lower in degraded forests, total evaporation (from ground and canopy intercepted wa-
 509 ter) was higher in most degraded forests, with burned forests experiencing 3–26% more
 510 evaporation in typical years and 0–14% during severe droughts (Figure 8c). The com-
 511 bination of higher evaporation and relatively shorter canopy (shallower roots) in degraded
 512 forests were typically translated into slightly drier near-surface soils (Figure 8d): dur-
 513 ing typical years, soil water availability at the top 30 cm layers was 1.2–12% lower in burned
 514 forests than intact forests, whereas the differences were more modest in logged forests
 515 (0.2–3%) and even reversed during extreme droughts (Figure 8d). Carbon and energy
 516 fluxes showed similar behavior. Gross primary productivity in intact forests steadily de-
 517 creased with increased drought severity, and the depletion of productivity caused by degra-
 518 dation is most marked during typical years but is reduced during severe droughts (Fig-
 519 ure S14a). While ground temperature is always higher in degraded forests (Figure S14b),
 520 differences in sensible heat fluxes and outgoing longwave radiation also diminish during
 521 extreme drought conditions (Figure S14c,d).

522 Degraded forests show drier near-surface soils (Figure 8d) and warmer surface tem-
 523 peratures (Figure S14) than intact forests for most years, yet the interannual variabil-
 524 ity of climate also modulates the differences in water, carbon, and energy cycles between
 525 degraded and intact forests (Figures 8 and S14). Therefore, both degradation and cli-
 526 mate may influence the flammability of forests. The average flammable area predicted
 527 by ED-2.2 (Section 2.4) shows large variation across regions, ranging from nearly zero
 528 at GYF forests (the wettest region) to over 25% yr⁻¹ at some of the forests in TAN (the
 529 driest region) (Figure 9a). Within each region (i.e. under the same prescribed climate),
 530 the model generally predicted higher flammability for the shortest forests (< 10 m), al-
 531 though predictions also indicate large within-region variability of flammable area for forests
 532 with intermediate canopy height (10–25 m) (Figure 9a). For most forests, flammable con-
 533 ditions were predicted mostly during moderate or severe droughts, regardless of the degra-




Figure 8. Response of the water cycle components across a forest degradation gradient and drought severity in Paragominas (PRG). Selected components: (a) Total water vapor flux, (b) stomatal conductance, averaged by leaf area, (c) evaporation, and (d) soil available water (i.e. in excess of permanent wilting point) of the top 30 cm. Points correspond to the median value of 12-month running averages, aggregated into 40 quantiles along the range of maximum cumulative water deficit (MCWD). Bands around the points correspond to the 95% range within each MCWD bin. Top panels are the absolute value for intact forests, and bottom panels are the absolute difference between degraded and intact forests. Background shades denote the MCWD anomaly: light gray – 68% range around the median (dot-dash vertical line); intermediate gray – 95% range; dark gray – anomalies exceeding the 95% range.




Figure 9. Average flammability as functions of degradation and climate variability. (a) Scatter plot shows the average flammable area (1980–2016) for each simulated patch across all regions, as a function of canopy height. Density cloud (background color) was produced through a bi-dimensional kernel density estimator; points are the averages used to generate each density cloud. Color ramps (logarithmic) range from 0.1 – 100% of the maximum computed scale. (b) Flammable area at region PRG, as a function of degradation history and drought length (number of consecutive months with water deficit in excess of 20 mm). Points correspond to the median value of 12-month running averages, aggregated into quantiles along the drought length. Bands around the points correspond to the 95% range within each drought length bin. Top panels are the absolute value for intact forests, and bottom panels are the absolute difference between degraded and intact forests. Background shades denote drought-length classes used in the text: seasonal (light gray, less than 12 months); severe (intermediate gray, 12–36 months); extreme (dark gray; more than 36 months). Flammability response to degradation and drought duration for other regions are shown in Figure S15.

dation history, as exemplified by region PRG (Figure 9b). While the time series of flammable area were synchronized across degradation types, ED-2.2 predictions of flammable area were generally higher for burned forests than intact or lightly logged forests (Figures 9b and S15). The one exception was the driest region (TAN), where forests that burned multiple times experienced lower flammability than intact forests (Figure S15d); at TAN, even intact forests were relatively short (Figure 9a), which caused ED-2.2 to predict limited access to deeper soils and increased desiccation.

4 Discussion

4.1 Initialization of forest structure from remote sensing

Our method to derive the vertical structure of the canopy from high-resolution airborne lidar successfully characterized the diversity of forest structures of the Amazon, captured differences in forest structure variability along a precipitation gradient, and described the within-region variability in forest structure caused by forest degradation (Figures 3 and S2-S3). Previous studies have used forest structure derived from remote-sensing data to initialize vegetation demography models in tropical forests (e.g., Hurtt et al., 2004; Antonarakis et al., 2011; Rödiger et al., 2018). However, these studies often assume a relationship between forest structure and canopy height with stand age. While this assumption has been successfully applied to intact and second-growth tropical forests (Hurtt et al., 2004; Antonarakis et al., 2011), the association between forest structure and succession is unlikely to be preserved in degraded forests. For example, understory fires proportionally kill more smaller trees than large trees (Uhl & Kauffman, 1990; Brando et al., 2012; Silva et al., 2018), and selective logging creates complex mosaics of forest structure, with substantial losses of large trees from harvesting, and extensive damage to smaller trees in skid trails (Feldpausch et al., 2005). In contrast, our approach accounts for the entire vertical profile at local (50-m) scale, similarly to Antonarakis et al. (2014), which does not require any assumption on the successional stage of the forest. Importantly, our approach requires only the vertical distribution of returns, and could be adapted to large-footprint, airborne or spaceborne lidar data, including the NASA’s Global Ecosystem Dynamics Investigation (GEDI, Hancock et al., 2019).

We demonstrated that the initialization from airborne lidar profiles captures most of the variability across and within regions, yet it has important assumptions and limitations. First, our approach relies on allometric equations to determine both the diameter at breast height (DBH), and the individual leaf area (L_i , Text S4.3), with the implicit assumption, that the contribution of branches, twigs, and stems to the lidar return signal is negligible. In reality, allometric equations have either large uncertainties (DBH) or limited number of samples (Figure S16). Previous studies using destructive sampling and terrestrial laser scanning suggest that wood area index may constitute 7–15% of the plant area index (Olivas et al., 2013; Schneider et al., 2019). The use of allometric equations that account for regional variation (e.g., Feldpausch et al., 2011, 2012), and the ex-

pansion of open-source databases, such as the Biomass And Allometry Database (BAAD, Falster et al., 2015) used in our study, could further improve the characterization of the vertical structure. In addition, the increased availability of terrestrial laser scanning (TLS) and high-resolution, low-altitude unmanned aerial vehicle lidar could substantially increase the data availability and thus improve the overall quality of allometric equations and constrain the relative contribution of woody tissues to the total plant area (Calders et al., 2015; Stovall et al., 2018; Schneider et al., 2019). Alternatively, techniques that extract individual tree crowns from lidar point clouds readily provide highly accurate local stem density and local size-frequency distributions (e.g., tree height or crown size; Ferraz et al., 2016, 2020). These distributions can be used to attribute DBH to individuals and generate initial conditions akin to forest inventory to the ED-2.2 model, and data-model fusion techniques that leverage the growing availability of data could reduce uncertainties on many model parameters, including allometry (F. J. Fischer et al., 2019). Finally, ED-2.2 overestimated the seasonality of gross primary productivity and evapotranspiration at the driest region (TAN) (Figures S4 and S6). This result suggests that simulated rooting depth for TAN was underestimated in the model. Rooting profiles in tropical forests remain largely uncertain: some site studies have sought to relate individual tree size with rooting depth using isotopic measurements (e.g., Stahl et al., 2013; Brum et al., 2019), whereas regional studies that provide spatial distribution of rooting depth still show important discrepancies in the tropics (e.g., Yang et al., 2016; Fan et al., 2017). Constraining the below-ground allocation of tropical ecosystems should be a priority in future studies.

In our study we inferred the functional diversity from forest structure obtained from existing forest inventory plots. The functional group attribution captured the general characteristics of functional composition along degradation gradients (Figure S1), including the more frequent occurrence of early-successional individuals in degraded forests, consistent with field-based studies (Both et al., 2019); nonetheless, uncertainties in functional attribution from field measurements are high. The increased availability of coordinated airborne laser scanning (ALS) and airborne imaging spectroscopy (AIS) data in mid-latitudes has lead to opportunities to link structural variability with functional diversity (e.g., Antonarakis et al., 2014; Schneider et al., 2017), and previous studies have successfully integrated ALS and AIS data to attribute functional groups in the ED-2 model (e.g., Antonarakis et al., 2014; Bogan et al., 2019). Overlapping ALS and AIS data over

tropical forests are becoming increasingly common (Asner et al., 2014; de Almeida et al., 2019; Laybros et al., 2019) and could provide new opportunities to reduce uncertainties in functional attribution in future studies. Likewise, ongoing and upcoming spaceborne missions at the International Space Station such as GEDI (Hancock et al., 2019), and the Hyperspectral Imaging Suite (HISUI, Matsunaga et al., 2017) will allow for large-scale characterization of structure and function of ecosystems at global scale (Stavros et al., 2017; Schimel et al., 2019).

4.2 Degradation impacts on ecosystem functioning

In addition to carbon losses and structural changes, degradation has substantial impacts on energy and water cycles in Amazonian forests, especially in severely degraded forests with marked dry season. According to the ED-2.2 simulations, ground temperature of logged forests ranged from nearly-identical to intact forests (low-impact logging or old logging disturbances) to 0.7°C warmer (recently logged forests), whereas severely burned forests experienced daytime near-surface temperatures increases of as much as 4°C (Figure S10), and differences between the lowest and highest biomass patches exceeded 9°C (Figure 6). Observed differences in understory temperatures show large variability, but they generally agree with the ED-2.2 results. For example, results of temperature differences between logged and intact areas in the wet forests of Sabah, Malaysia, ranged from negligible to 1.2°C for average maximum temperature (Senior et al., 2018; Jucker et al., 2018). The predicted warmer daytime understory temperatures at recurrently burned forests also yielded drier near-surface conditions: daytime ground vapor pressure deficit was on average 15–25 hPa greater than in intact forests (equivalent to 5–15% reduction in relative humidity), which is within the range observed after the most damaging experimental fire at TAN in 2007 (Brando et al., 2014), and similar to differences in understory relative humidity reported in the dry season between open-canopy seasonally flooded forests and closed-canopy upland forests in the Central Amazon (de Resende et al., 2014). Because temperatures are higher in degraded forests, the simulated changes in energy and water cycle caused by degradation also point to a reduction of entropy production in degraded forests, which is consistent with the results across pastures and intact forests across the Amazon (Holdaway et al., 2010).

ED-2.2 showed various degrees of agreement with the few existing observational studies comparing changes in evapotranspiration due to degradation. Evapotranspira-

tion response to reduced-impact logging was minor (-1.9% reduction relative to intact in BTE), consistent with eddy covariance tower estimates in a logging experiment in the same region (-3.7% reduction after accounting for site differences and interannual variability, S. D. Miller et al., 2011). The model results for the experimental fire at TAN, however, suggested similar wet-season ET between burned and intact forests ($\Delta ET = ET_{\text{Brn}} - ET_{\text{Int}} = 0.002 \text{ mm day}^{-1}$), with stronger depletion of ET in burned forests during the dry season ($\Delta ET = -0.31 \text{ mm day}^{-1}$) (Figures 5 and S6). In contrast, Brando, Silvério, et al. (2019) found higher ET in burned forests over a period of 4 years, albeit ΔET also showed significant interannual variability. A few other studies suggest that the significant decline in dry-season ET in burned forests may be expected in some areas: for example, Hirano et al. (2015) found that evapotranspiration of drained and burned peatlands with second-growth vegetation in Central Kalimantan (Indonesia) was 0.43 mm day^{-1} lower than drained forests; Quesada et al. (2004) inferred ET changes from soil water budget in savannas and found significant reductions following fires in a savanna site in Central Brazil. The advent of high-resolution remote sensing products that quantify energy, water, and carbon fluxes, such as the ECOSystem Spaceborne Thermal Radiometer Experiment on Space Station (ECOSTRESS) and the Orbiting Carbon Observatory 3 (OCO-3), will provide new opportunities to quantify the role of tropical forest degradation on ecosystem functioning at regional scale (Schimel et al., 2019), as well as to provide new benchmark data for ecosystem models.

Our model results indicate that severe degradation substantially alters the magnitude and seasonality of energy, water, and carbon fluxes (Figures 5-7 and S10-S12). In our study, we disabled the vegetation dynamics in ED-2.2 to ensure that predicted differences in ecosystem functioning could be unequivocally attributed to structural diversity, but the differences in ecosystem functioning between degraded and intact forests may diminish over time as the forest recovers from previous disturbance. This pathway is consistent with the relatively small differences in ET and surface temperature (Figures 5-6) observed at logged forests at GYF (25 years since last disturbance) and burned forests at BTE (15 years since last disturbance). However, the recovery trajectory is one out of multiple possible pathways: degraded forests may be more prone to subsequent disturbances (Silvério et al., 2019; Hérault & Piponiot, 2018); the recovery dynamics can be long or not attainable if multiple stable states exist or if succession is arrested (Mesquita et al., 2015; Ghazoul & Chazdon, 2017), potentially prolonging the impacts of forest degra-

dation on energy and water cycles; and feedbacks on precipitation caused by degradation could affect the spatial distribution of rainfall similarly to the effect observed with deforestation (Spracklen et al., 2018), although to our knowledge this impact has not yet been quantified for degraded forests.

In this study, we focused on the effects of forest structure on ecosystem function, and thus we used idealized, homogeneous soil with intermediate hydraulic characteristics in all simulations. In reality, soils across the Amazon are highly heterogeneous and directly affect forest structure across the biome (Quesada et al., 2012). Likewise, soil depth and texture and variability in local topography also modulate the effects of tropical forest degradation on microclimate (Jucker et al., 2018). A previous study using ED-2.2 found that evapotranspiration in Central Amazonia could decrease by 12–16% under scenarios of recurrent yearlong droughts (40% reduction in rainfall), but the severity of the decrease varied by 7% under the same climate scenarios but different soil hydraulic properties (Longo et al., 2018). These results suggest that degraded forests in clay-rich, compact soils and deeper water table could amplify reductions in evapotranspiration and gross primary productivity during the dry season, while degradation effects on energy, water, and carbon cycle would likely be dampened in regions where the water table is near the surface for most of the year, or soils with higher water storage capacity.

4.3 Interactions between forest degradation and climate variability

The predicted reductions in evapotranspiration (ET) in the most degraded areas during the dry season suggest that land-use change impacts on the water cycle may be more widespread and pervasive than indicated by earlier studies. Previous model-based studies showed that biome-wide deforestation could cause ET to decrease by 25–40% relative to intact forests in the Amazon during the dry season (e.g., von Randow et al., 2004; Zemp et al., 2017). These reductions are comparable to the ET reductions predicted by ED-2.2 at the most degraded forests (21–32%, Figure 5). Because tropical forest degradation affects an area comparable to deforestation in the Amazon (Tyukavina et al., 2017), it may further reduce the strength of the Amazon water vapor source to the atmosphere. In our study, we focused on understanding how climate and structure variability impacts the water and energy fluxes, but degradation-driven changes in these fluxes are likely to feed back into the atmosphere. For example, changes in evapotranspiration and sensible heat flux associated with deforestation are known to either redistribute or reduce to-

tal rainfall in tropical forests (Spracklen et al., 2018, and references therein), and a substantial fraction of South American precipitation water comes from evapotranspiration from Amazonian forests (van der Ent et al., 2010). Recent estimates of ET for the Amazon Basin from the Gravity Recovery and Climate Experiment (GRACE) suggest that the basin-wide ET (including intact forests) has decreased by 1.7% between 2002 and 2015 (Swann & Koven, 2017). In addition, several studies suggest that the dry season in the Amazon is becoming longer (Fu et al., 2013; Sena et al., 2018), and land use change is one of the main drivers of the drying trend (Barkhordarian et al., 2018). The role of forest degradation on ongoing and future changes in climate across the Amazon remains uncertain and deserves further investigation, potentially with coupled biosphere-atmosphere models that represent heterogeneity in forest structure and functioning (Swann et al., 2015; Knox et al., 2015; Wu et al., 2017). Likewise, we could not account for cascading effects of climate on the energy, water, and carbon cycle in this study because we disabled dynamic vegetation. However, severe droughts are known to increase mortality rates and canopy turnover in tropical forests (Phillips et al., 2010; Feldpausch et al., 2016; Leitold et al., 2018); such disturbances may increase gap fraction and thus reduce gross primary productivity and evapotranspiration in the years immediately following the drought. Future studies that include dynamic vegetation can provide further insights on the resilience and resistance of degraded and intact forests to climate extreme.

Our results show that structural changes resulting from forest degradation make the forest surface drier and warmer (Figures 5-8 and S10). Drier and warmer conditions near the surface increase flammability (Brando, Paolucci, et al., 2019, and references therein), and it has been long suggested that forest degradation and canopy opening make forests more likely to burn (e.g., Uhl & Buschbacher, 1985; Cochrane et al., 1999; Ray et al., 2005; A. A. C. Alencar et al., 2015). The ED-2.2 simulations indeed predicted higher flammability in degraded (more open-canopy) forests on any given year (Figures 9 and S15). However, our results also suggest that climate strongly drives the variability of flammable area across most of our study regions (Figures 9b and S15), which is consistent with the significant increases in forest fires in the Amazon during extreme drought years (Morton et al., 2013; Aragão et al., 2018). Moreover, our results indicate that differences in flammable area between intact and degraded forests are reduced or even reversed during extreme droughts, which indicates that under extreme conditions, the level of degradation is less critical to create flammable conditions. This effect was predicted for most years at TAN,

which typically experiences severe and longer dry seasons compared to the other study regions (Figure S15).

Previous studies suggest that parts of the Eastern Amazon could become drier by the end of the century and experience more extreme events, including droughts (IPCC, 2014; Duffy et al., 2015), and thus potentially more susceptible to future fires (De Faria et al., 2017; Brando et al., 2020). However, how tropical forest flammability will respond in the long-term to ongoing changes in climate and land use is still uncertain, and recent studies have shown that either climate (Le Page et al., 2017) or land use (Fonseca et al., 2019) could be dominant on predicted shifts in fire regime. Importantly, while our analysis focused on flammability, and ED-2.2 fire model captures the general patterns of fire disturbance across the Amazon (Longo, Knox, Levine, et al., 2019), it does not represent many mechanisms and processes that are critical to describe fire dynamics in tropical forests, such as anthropogenic ignitions, diurnal cycle of fire intensity, and fire termination, therefore we could not quantify the effects of fire on further forest degradation. The use of process-based fire disturbance models within the ED-2.2 (e.g., Thonicke et al., 2010; Le Page et al., 2015) framework could contribute to further improve our understanding of interactions between forest degradation, climate, and flammability across the Amazon.

5 Conclusion

Our study showed that tropical forest degradation can markedly modify the ecosystem functioning in the Amazon, with substantial reductions in evapotranspiration (ET) and gross primary productivity (GPP), and increase in surface temperature (Figures 5–8). Within the regions included in our study, the effects of degradation on energy, water, and carbon cycles were the strongest in the Eastern and Southern Amazon, where the dry season is more pronounced. Notably, in areas where severe forest degradation resulted in substantial changes in forest structure, reductions in dry-season evapotranspiration are similar to those found in deforested areas (Figure 5; von Randow et al., 2004). The area of the Amazon forest impacted by degradation is comparable to the deforested area (Asner et al., 2005; Morton et al., 2013; Souza Jr. et al., 2013; Tyukavina et al., 2017), and thus degradation-driven changes in water, energy, and carbon cycles are potentially important. However, the extent to which degradation affects the biophysical and biogeochemical cycles at regional scale ultimately depends on (1) annual degradation rates;

(2) recovery time of degraded forests; and (3) the likelihood that degraded forests are cleared. For example, (Brando, Silv rio, et al., 2019) found that ET in burned forests was indistinguishable from intact forests 7 years after the last fire. While their result suggests fast recovery of degraded forests, the impacts of degradation on ET can still be regionally relevant if degradation rates are sufficiently high to maintain low average age since last disturbance in degraded forests. Moreover, we found that the impacts of tropical forest degradation on energy, water, and carbon cycles and on flammability are more pronounced during typical years than during extreme droughts (when all forests become flammable), which highlights the complex interactions between climate and forest structure. To understand and reduce uncertainties of climate-structure interactions, it would be valuable to leverage the recent advances in remote sensing of forest structure, including the recently launched GEDI mission (Hancock et al., 2019), and terrestrial biosphere models that can represent complex and heterogeneous ecosystems (Fisher et al., 2018). Our study, while focusing on airborne lidar data, has demonstrated the opportunities to integrate remote sensing and terrestrial biosphere models even in regions with complex forest structure such as degraded forests.

Acknowledgments

Airborne lidar and forest inventory data were obtained from Sustainable Landscapes Brazil (2019), dos-Santos et al. (2019) (Brazil) and Paracou Portal (2016) (French Guiana). MERRA-2 reanalyses are available from GMAO (2015a, 2015b, 2015c, 2015d) and MSWEP-2.2 data were downloaded from <http://www.gloh2o.org>. The ED-2.2 model used in this study is available at Longo et al. (2019) and the scripts and ED-2.2 output are permanently stored at Longo et al. (2020). Trait data are available at the TRY initiative on plant traits (<http://www.try-db.org>), request 2751; at Gu et al. (2016); or as supporting information from the cited references (I. J. Wright et al., 2004; Santiago & Wright, 2007; Bahar et al., 2017).

The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D004), and supported by NASA Earth Sciences grant (NNH16ZDA001N-IDS, grant 16-IDS16-0049). Data recorded at the Guyaflux tower were obtained thanks to the support of two *Investissement d’avenir* grants from the Agence Nationale de la Recherche (CEBA, ref ANR-10-LABX-25-01; ARBRE, ref. ANR-11-LABX-0002-01). Data

in Brazil were acquired by the Sustainable Landscapes Brazil project supported by the Brazilian Agricultural Research Corporation (EMBRAPA), the U.S. Forest Service, and USAID, and the U.S. Department of State. The study has been supported by the TRY initiative on plant traits, which is hosted, developed and maintained by J. Kattge and G. Bönisch (Max Planck Institute for Biogeochemistry, Jena, Germany). TRY is currently supported by DIVERSITAS/Future Earth and the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig.

We thank Xiangtao Xu for sharing the trait plasticity algorithm and discussions on model results; Divino Silvério for processing and sharing the data from Tanguro; Hylke Beck for sharing the MSWEP-2.2 data; and Marcos Scaranello, Fabian Schneider, Alexandra Konings, and A. Anthony Bloom for discussions on the lidar initialization algorithm and interpretation of model results. The model simulations were carried out at the Odyssey cluster, supported by the FAS Division of Science, Research Computing Group at Harvard University; and at the Brazilian National Laboratory for Scientific Computing (LNCC). M.L. was supported by the São Paulo State Research Foundation (FAPESP, 2015/07227-6) and by the NASA Postdoctoral Program, administered by Universities Space Research Association under contract with NASA. M.K. was supported in part by the Next Generation Ecosystem Experiments-Tropics, funded by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research.

References

- Achard, F., Beuchle, R., Mayaux, P., Stibig, H.-J., Bodart, C., Brink, A., ... Simonetti, D. (2014, Aug). Determination of tropical deforestation rates and related carbon losses from 1990 to 2010. *Glob. Change Biol.*, 20(8), 2540–2554. doi: 10.1111/gcb.12605
- Alamgir, M., Campbell, M. J., Turton, S. M., Pert, P. L., Edwards, W., & Laurance, W. F. (2016, Jul). Degraded tropical rain forests possess valuable carbon storage opportunities in a complex, forested landscape. *Sci. Rep.*, 6(30012). doi: 10.1038/srep30012
- Alencar, A., Nepstad, D., & Vera Diaz, M. C. (2006, Feb). Forest understory fire in the Brazilian Amazon in ENSO and non-ENSO years: Area burned and committed carbon emissions. *Earth Interact.*, 10(6), 1–17. doi: 10.1175/EI150.1
- Alencar, A. A. C., Brando, P. M., Asner, G. P., & Putz, F. E. (2015, Sep). Land-

- scape fragmentation, severe drought and the new Amazon forest fire regime. *Ecol. Appl.*, *25*(6), 1493–1505. doi: 10.1890/14-1528.1
- Antonarakis, A. S., Munger, J. W., & Moorcroft, P. R. (2014, Jul). Imaging spectroscopy- and lidar-derived estimates of canopy composition and structure to improve predictions of forest carbon fluxes and ecosystem dynamics. *Geophys. Res. Lett.*, *41*(7), 2535–2542. doi: 10.1002/2013GL058373
- Antonarakis, A. S., Saatchi, S. S., Chazdon, R. L., & Moorcroft, P. R. (2011, Jun). Using lidar and radar measurements to constrain predictions of forest ecosystem structure and function. *Ecol. Appl.*, *21*(4), 1120–1137. doi: 10.1890/10-0274.1
- Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., ... Saatchi, S. (2018, Feb). 21st century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nature Comm.*, *9*(1), 536. doi: 10.1038/s41467-017-02771-y
- Aragão, L. E. O. C., Poulter, B., Barlow, J. B., Anderson, L. O., Malhi, Y., Saatchi, S., ... Gloor, E. (2014, Nov). Environmental change and the carbon balance of Amazonian forests. *Biol. Rev.*, *89*(4), 913–931. doi: 10.1111/brv.12088
- Asner, G. P., Broadbent, E. N., Oliveira, P. J. C., Keller, M., Knapp, D. E., & Silva, J. N. M. (2006, Aug). Condition and fate of logged forests in the Brazilian Amazon. *Proc. Natl. Acad. Sci. U. S. A.*, *103*(34), 12947–12950. doi: 10.1073/pnas.0604093103
- Asner, G. P., Knapp, D. E., Broadbent, E. N., Oliveira, P. J. C., Keller, M., & Silva, J. N. (2005, Oct). Selective logging in the Brazilian Amazon. *Science*, *310*(5747), 480–482. doi: 10.1126/science.1118051
- Asner, G. P., Martin, R. E., Tupayachi, R., Anderson, C. B., Sinca, F., Carranza-Jiménez, L., & Martinez, P. (2014, Apr). Amazonian functional diversity from forest canopy chemical assembly. *Proc. Natl. Acad. Sci. U. S. A.*, *111*(15), 5604–5609. doi: 10.1073/pnas.1401181111
- Asner, G. P., Powell, G. V. N., Mascaro, J., Knapp, D. E., Clark, J. K., Jacobson, J., ... Hughes, R. F. (2010, Sep). High-resolution forest carbon stocks and emissions in the Amazon. *Proc. Natl. Acad. Sci. U. S. A.*, *107*(38), 16738–16742. doi: 10.1073/pnas.1004875107
- Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., & Houghton,

- 865 R. A. (2017, Oct). Tropical forests are a net carbon source based on above-
 866 ground measurements of gain and loss. *Science*, *358*(6360), 230–234. doi:
 867 10.1126/science.aam5962
- 868 Bahar, N. H. A., Ishida, F. Y., Weerasinghe, L. K., Guerrieri, R., O’Sullivan, O. S.,
 869 Bloomfield, K. J., ... Atkin, O. K. (2017, May). Leaf-level photosynthetic ca-
 870 pacity in lowland Amazonian and high-elevation Andean tropical moist forests
 871 of Peru. *New Phytol.*, *214*(3), 1002–1018. doi: 10.1111/nph.14079
- 872 Balch, J. K., Nepstad, D. C., Brando, P. M., Curran, L. M., Portela, O., de Car-
 873 valho, O., & Lefebvre, P. (2008, Oct). Negative fire feedback in a transitional
 874 forest of southeastern Amazonia. *Glob. Change Biol.*, *14*(10), 2276–2287. doi:
 875 10.1111/j.1365-2486.2008.01655.x
- 876 Baraloto, C., Paine, C. E. T., Poorter, L., Beauchene, J., Bonal, D., Domen-
 877 ach, A.-M., ... Chave, J. (2010, Nov). Decoupled leaf and stem eco-
 878 nomics in rain forest trees. *Ecol. Lett.*, *13*(11), 1338–1347. doi: 10.1111/
 879 j.1461-0248.2010.01517.x
- 880 Barkhordarian, A., von Storch, H., Behrangi, A., Loikith, P. C., Mechoso, C. R.,
 881 & Detzer, J. (2018, Jun). Simultaneous regional detection of land-
 882 use changes and elevated GHG levels: the case of spring precipitation in
 883 tropical South America. *Geophys. Res. Lett.*, *45*(12), 6262–6271. doi:
 884 10.1029/2018GL078041
- 885 Barlow, J., Lagan, B. O., & Peres, C. A. (2003, 5). Morphological correlates of
 886 fire-induced tree mortality in a central Amazonian forest. *J. Trop. Ecol.*, *19*(3),
 887 291–299. doi: 10.1017/S0266467403003328
- 888 Barlow, J., Lennox, G. D., Ferreira, J., Berenguer, E., Lees, A. C., Mac Nally, R.,
 889 ... Gardner, T. A. (2016, Jul). Anthropogenic disturbance in tropical forests
 890 can double biodiversity loss from deforestation. *Nature*, *535*, 144–147. doi:
 891 10.1038/nature18326
- 892 Beck, H. E., Wood, E. F., Pan, M., Fisher, C. K., Miralles, D. G., van Dijk, A. I.,
 893 ... Adler, R. F. (2019, Mar). MSWEP V2 global 3-hourly 0.1° precipitation:
 894 methodology and quantitative assessment. *Bull. Am. Meteorol. Soc.*, *100*(3),
 895 473–500. doi: 10.1175/BAMS-D-17-0138.1
- 896 Berenguer, E., Ferreira, J., Gardner, T. A., Aragão, L. E. O. C., de Camargo, P. B.,
 897 Cerri, C. E., ... Barlow, J. (2014, Dec). A large-scale field assessment of

- 898 carbon stocks in human-modified tropical forests. *Glob. Change Biol.*, 20(12),
899 3713–3726. doi: 10.1111/gcb.12627
- 900 Blair, J. B., & Hofton, M. A. (1999, Aug). Modeling laser altimeter return wave-
901 forms over complex vegetation using high-resolution elevation data. *Geophys.*
902 *Res. Lett.*, 26(16), 2509–2512. doi: 10.1029/1999GL010484
- 903 Blanc, L., Echard, M., Herault, B., Bonal, D., Marcon, E., Chave, J., & Baraloto, C.
904 (2009, Sep). Dynamics of aboveground carbon stocks in a selectively logged
905 tropical forest. *Ecol. Appl.*, 19(6), 1397–1404. doi: 10.1890/08-1572.1
- 906 Bogan, S. A., Antonarakis, A. S., & Moorcroft, P. R. (2019, Jul). Imag-
907 ing spectrometry-derived estimates of regional ecosystem composition for
908 the Sierra Nevada, California. *Remote Sens. Environ.*, 228, 14–30. doi:
909 10.1016/j.rse.2019.03.031
- 910 Bonal, D., Bosc, A., Ponton, S., Goret, J.-Y., Burban, B., Gross, P., ... Granier,
911 A. (2008, Aug). Impact of severe dry season on net ecosystem exchange
912 in the Neotropical rainforest of French Guiana. *Glob. Change Biol.*, 14(8),
913 1917–1933. doi: 10.1111/j.1365-2486.2008.01610.x
- 914 Both, S., Riutta, T., Paine, C. E. T., Elias, D. M. O., Cruz, R. S., Jain, A., ...
915 Burslem, D. F. R. P. (2019, Feb). Logging and soil nutrients independently ex-
916 plain plant trait expression in tropical forests. *New Phytol.*, 221(4), 1853–1865.
917 doi: 10.1111/nph.15444
- 918 Braghiere, R. K., Quaife, T., Black, E., He, L., & Chen, J. M. (2019, Nov). Underes-
919 timation of global photosynthesis in earth system models due to representation
920 of vegetation structure. *Global Biogeochem. Cycles*, 33(11), 1358–1369. doi:
921 10.1029/2018GB006135
- 922 Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe,
923 M. T., ... Soares-Filho, B. S. (2014, Apr). Abrupt increases in Amazonian
924 tree mortality due to drought–fire interactions. *Proc. Natl. Acad. Sci. U. S. A.*,
925 111(17), 6347–6352. doi: 10.1073/pnas.1305499111
- 926 Brando, P. M., Nepstad, D. C., Balch, J. K., Bolker, B., Christman, M. C., Coe, M.,
927 & Putz, F. E. (2012, Feb). Fire-induced tree mortality in a neotropical forest:
928 the roles of bark traits, tree size, wood density and fire behavior. *Glob. Change*
929 *Biol.*, 18(2), 630–641. doi: 10.1111/j.1365-2486.2011.02533.x
- 930 Brando, P. M., Paolucci, L., Ummenhofer, C. C., Ordway, E. M., Hartmann, H.,

- 931 Cattau, M. E., ... Balch, J. (2019, May). Droughts, wildfires, and forest
 932 carbon cycling: A pantropical synthesis. *Annu. Rev. Earth Pl. Sc.*, 47(1),
 933 555–581. doi: 10.1146/annurev-earth-082517-010235
- 934 Brando, P. M., Silvério, D., Maracahipes-Santos, L., Oliveira-Santos, C., Levick,
 935 S. R., Coe, M. T., ... Trumbore, S. E. (2019, Sep). Prolonged tropical forest
 936 degradation due to compounding disturbances: implications for CO₂ and H₂O
 937 fluxes of an experimental forest. *Glob. Change Biol.*, 25(9), 2855–2868. doi:
 938 10.1111/gcb.14659
- 939 Brando, P. M., Soares Filho, B., Rodrigues, L., Assunção, A., Morton, D., Tuch-
 940 schneider, D., ... Coe, M. T. (2020, Jan). The gathering firestorm in southern
 941 Amazonia. *Sci. Adv.*, 6(2), eaay1632. doi: 10.1126/sciadv.aay1632
- 942 Brum, M., Vadeboncoeur, M. A., Ivanov, V., Asbjornsen, H., Saleska, S., Alves,
 943 L. F., ... Oliveira, R. S. (2019, Jan). Hydrological niche segregation defines
 944 forest structure and drought tolerance strategies in a seasonal Amazon forest.
 945 *J. Ecol.*, 107(1), 318–333. doi: 10.1111/1365-2745.13022
- 946 Bustamante, M. M. C., Roitman, I., Aide, T. M., Alencar, A., Anderson, L., Aragão,
 947 L., ... Vieira, I. C. (2016, Jan). Towards an integrated monitoring frame-
 948 work to assess the effects of tropical forest degradation and recovery on
 949 carbon stocks and biodiversity. *Glob. Change Biol.*, 22(1), 92–109. doi:
 950 10.1111/gcb.13087
- 951 Calders, K., Newnham, G., Burt, A., Murphy, S., Raunonen, P., Herold, M., ...
 952 Kaasalainen, M. (2015, Feb). Nondestructive estimates of above-ground
 953 biomass using terrestrial laser scanning. *Methods Ecol. Evol.*, 6(2), 198–208.
 954 doi: 10.1111/2041-210X.12301
- 955 Chave, J., Condit, R., Aguilar, S., Hernandez, A., Lao, S., & Perez, R. (2004, Mar).
 956 Error propagation and scaling for tropical forest biomass estimates. *Philos.*
 957 *Trans. R. Soc. B-Biol. Sci.*, 359(1443), 409–420. doi: 10.1098/rstb.2003.1425
- 958 Chave, J., Coomes, D., Jansen, S., Lewis, S. L., Swenson, N. G., & Zanne, A. E.
 959 (2009, Apr). Towards a worldwide wood economics spectrum. *Ecol. Lett.*,
 960 12(4), 351–366. doi: 10.1111/j.1461-0248.2009.01285.x
- 961 Chave, J., Réjou-Méchain, M., Búrquez, A., Chidumayo, E., Colgan, M. S., Delitti,
 962 W. B., ... Vieilledent, G. (2014, Oct). Improved allometric models to esti-
 963 mate the aboveground biomass of tropical trees. *Glob. Change Biol.*, 20(10),

- 3177–3190. doi: 10.1111/gcb.12629
- Cochrane, M. A., Alencar, A., Schulze, M. D., Souza, C. M., Nepstad, D. C., Lefebvre, P., & Davidson, E. A. (1999, Jun). Positive feedbacks in the fire dynamic of closed canopy tropical forests. *Science*, *284*(5421), 1832–1835. doi: 10.1126/science.284.5421.1832
- Davidson, E. A., de Araújo, A. C., Artaxo, P., Balch, J. K., Brown, I. F., Bus-tamante, M. M. C., ... Wofsy, S. C. (2012, Jan). The Amazon basin in transition. *Nature*, *481*(7381), 321–328. doi: 10.1038/nature10717
- de Almeida, C. T., Galvão, L. S., Aragão, L. E. d. O. C. e., Ometto, J. P. H. B., Jacon, A. D., Pereira, F. R. d. S., ... Longo, M. (2019, Oct). Combining LiDAR and hyperspectral data for aboveground biomass modeling in the Brazilian Amazon using different regression algorithms. *Remote Sens. Environ.*, *232*, 111323. doi: 10.1016/j.rse.2019.111323
- De Faria, B. L., Brando, P. M., Macedo, M. N., Panday, P. K., Soares Filho, B. S., & Coe, M. T. (2017, Sep). Current and future patterns of fire-induced forest degradation in Amazonia. *Environ. Res. Lett.*, *12*(9), 095005. doi: 10.1088/1748-9326/aa69ce
- de Resende, A. F., Nelson, B. W., Flores, B. M., & de Almeida, D. R. (2014, Nov). Fire damage in seasonally flooded and upland forests of the Central Amazon. *Biotropica*, *46*(6), 643–646. doi: 10.1111/btp.12153
- d'Oliveira, M. V. N., Reutebuch, S. E., McGaughey, R. J., & Andersen, H.-E. (2012, Sep). Estimating forest biomass and identifying low-intensity logging areas using airborne scanning lidar in Antimary State Forest, Acre state, western Brazilian Amazon. *Remote Sens. Environ.*, *124*, 479–491. doi: 10.1016/j.rse.2012.05.014
- dos-Santos, M., Keller, M., & Morton, D. (2019, Dec). *LiDAR surveys over selected forest research sites, Brazilian Amazon, 2008–2018*. Retrieved 31 Jan 2020, from https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1644 doi: 10.3334/ORNLDAAAC/1644
- Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015, Oct). Projections of future meteorological drought and wet periods in the Amazon. *Proc. Natl. Acad. Sci. U. S. A.*, *112*(43), 13172–13177. doi: 10.1073/pnas.1421010112
- Erb, K.-H., Fetzl, T., Plutzer, C., Kastner, T., Lauk, C., Mayer, A., ... Haberl, H.

- (2016, Sep). Biomass turnover time in terrestrial ecosystems halved by land use. *Nature Geosci.*, *9*(9), 674–678. doi: 10.1038/ngeo2782
- Erb, K.-H., Kastner, T., Plutzer, C., Bais, A. L. S., Carvalhais, N., Fetzel, T., ... Luysaert, S. (2018, Jan). Unexpectedly large impact of forest management and grazing on global vegetation biomass. *Nature*, *553*(7686), 73–76. doi: 10.1038/nature25138
- Falster, D. S., Duursma, R. A., Ishihara, M. I., Barneche, D. R., FitzJohn, R. G., Vårhammar, A., ... York, R. A. (2015, May). BAAD: a biomass and allometry database for woody plants. *Ecology*, *96*(5), 1445–1445. doi: 10.1890/14-1889.1
- Falster, D. S., FitzJohn, R. G., Brännström, Å., Dieckmann, U., & Westoby, M. (2016, Feb). **plant**: A package for modelling forest trait ecology and evolution. *Methods Ecol. Evol.*, *7*(2), 136–146. doi: 10.1111/2041-210X.12525
- Fan, Y., Miguez-Macho, G., Jobbágy, E. G., Jackson, R. B., & Otero-Casal, C. (2017, Oct). Hydrologic regulation of plant rooting depth. *Proc. Natl. Acad. Sci. U. S. A.*, *114*(40), 10572–10577. doi: 10.1073/pnas.1712381114
- Feldpausch, T. R., Banin, L., Phillips, O. L., Baker, T. R., Lewis, S. L., Quesada, C. A., ... Lloyd, J. (2011, May). Height-diameter allometry of tropical forest trees. *Biogeosciences*, *8*(5), 1081–1106. doi: 10.5194/bg-8-1081-2011
- Feldpausch, T. R., Jirka, S., Passos, C. A. M., Jasper, F., & Riha, S. J. (2005, 11). When big trees fall: Damage and carbon export by reduced impact logging in southern Amazonia. *Forest Ecol. Manag.*, *219*(2–3), 199–215. doi: 10.1016/j.foreco.2005.09.003
- Feldpausch, T. R., Lloyd, J., Lewis, S. L., Brien, R. J. W., Gloor, M., Monteagudo Mendoza, A., ... Phillips, O. L. (2012, Aug). Tree height integrated into pantropical forest biomass estimates. *Biogeosciences*, *9*(8), 3381–3403. doi: 10.5194/bg-9-3381-2012
- Feldpausch, T. R., Phillips, O. L., Brien, R. J. W., Gloor, E., Lloyd, J., Lopez-Gonzalez, G., ... Vos, V. A. (2016, Jul). Amazon forest response to repeated droughts. *Global Biogeochem. Cycles*, *30*(7), 964–982. doi: 10.1002/2015GB005133
- Ferraz, A., Saatchi, S., Longo, M., & Clark, D. B. (2020). Tropical tree size demography from airborne lidar. *Ecol. Appl.* (advance online publication) doi: 10

- 1030 .1002/eap.2154
- 1031 Ferraz, A., Saatchi, S., Mallet, C., & Meyer, V. (2016, Sep). Lidar detection of indi-
 1032 vidual tree size in tropical forests. *Remote Sens. Environ.*, *183*, 318–333. doi:
 1033 10.1016/j.rse.2016.05.028
- 1034 Ferraz, A., Saatchi, S., Xu, L., Hagen, S., Chave, J., Yu, Y., ... Ganguly, S. (2018,
 1035 Sep). Carbon storage potential in degraded forests of Kalimantan, Indonesia.
 1036 *Environ. Res. Lett.*, *13*(9), 095001. doi: 10.1088/1748-9326/aad782
- 1037 Fischer, F. J., Maréchaux, I., & Chave, J. (2019). Improving plant allometry by fus-
 1038 ing forest models and remote sensing. *New Phytol.* (advance online publica-
 1039 tion) doi: 10.1111/nph.15810
- 1040 Fischer, R., Bohn, F., de Paula, M. D., Dislich, C., Groeneveld, J., Gutiérrez, A. G.,
 1041 ... Huth, A. (2016, Apr). Lessons learned from applying a forest gap model to
 1042 understand ecosystem and carbon dynamics of complex tropical forests. *Ecol.*
 1043 *Model.*, *326*, 124–133. doi: 10.1016/j.ecolmodel.2015.11.018
- 1044 Fisher, R. A., Koven, C. D., Anderegg, W. R. L., Christoffersen, B. O., Dietze,
 1045 M. C., Farrior, C., ... Moorcroft, P. (2018, Jan). Vegetation demographics in
 1046 Earth system models: a review of progress and priorities. *Glob. Change Biol.*,
 1047 *24*(1), 35–54. doi: 10.1111/gcb.13910
- 1048 Fonseca, M. G., Alves, L. M., Aguiar, A. P. D., Arai, E., Anderson, L. O., Rosan,
 1049 T. M., ... de Aragão, L. E. O. e. C. (2019). Effects of climate and land-use
 1050 change scenarios on fire probability during the 21st century in the Brazilian
 1051 Amazon. *Glob. Change Biol.* doi: 10.1111/gcb.14709
- 1052 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Hauck, J., Peters,
 1053 G. P., ... Zaehle, S. (2019, Dec). Global carbon budget 2019. *Earth Syst. Sci.*
 1054 *Data*, *11*(4), 1783–1838. doi: 10.5194/essd-11-1783-2019
- 1055 Fu, R., Yin, L., Li, W., Arias, P. A., Dickinson, R. E., Huang, L., ... Myneni, R. B.
 1056 (2013, Nov). Increased dry-season length over southern Amazonia in recent
 1057 decades and its implication for future climate projection. *Proc. Natl. Acad.*
 1058 *Sci. U. S. A.*, *110*(45), 18110–18115. doi: 10.1073/pnas.1302584110
- 1059 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., ...
 1060 Zhao, B. (2017, Jul). The Modern-Era Retrospective analysis for Research
 1061 and Applications, version 2 (MERRA-2). *J. Climate*, *30*(14), 5419–5454. doi:
 1062 10.1175/JCLI-D-16-0758.1

- 1063 Ghazoul, J., & Chazdon, R. (2017, Oct). Degradation and recovery in changing for-
1064 est landscapes: A multiscale conceptual framework. *Ann. Rev. Environ. Res.*,
1065 42(1), 161–188. doi: 10.1146/annurev-environ-102016-060736
- 1066 GMAO. (2015a). *MERRA-2 tavg1_2d_flux_Nx: 2d, 1-hourly, time-averaged,*
1067 *single-level, assimilation, surface flux diagnostics V5.12.4.* Retrieved 06
1068 Sep 2018, from <https://doi.org/10.5067/7MCPBJ41Y0K6> doi: 10.5067/
1069 7MCPBJ41Y0K6
- 1070 GMAO. (2015b). *MERRA-2 tavg1_2d_lnd_Nx: 2d, 1-hourly, time-averaged,*
1071 *single-level, assimilation, land surface diagnostics V5.12.4.* Retrieved
1072 06 Sep 2018, from <https://doi.org/10.5067/RKPHT8KC1Y1T> doi:
1073 10.5067/RKPHT8KC1Y1T
- 1074 GMAO. (2015c). *MERRA-2 tavg1_2d_rad_Nx: 2d, 1-hourly, time-averaged, single-*
1075 *level, assimilation, radiation diagnostics V5.12.4.* Retrieved 06 Sep 2018, from
1076 <https://doi.org/10.5067/Q9QMY5PBNV1T> doi: 10.5067/Q9QMY5PBNV1T
- 1077 GMAO. (2015d). *MERRA-2 tavg1_2d_slv_Nx: 2d, 1-hourly, time-averaged,*
1078 *single-level, assimilation, single-level diagnostics V5.12.4.* Retrieved 06
1079 Sep 2018, from <https://doi.org/10.5067/VJAFPLI1CSIV> doi: 10.5067/
1080 VJAFPLI1CSIV
- 1081 Gourlet-Fleury, S., Ferry, B., Molino, J.-F., Petronelli, P., & Schmitt, L. (2004).
1082 Experimental plots: Key features. In S. Gourlet-Fleury, J.-M. Guehl, &
1083 O. Laroussinie (Eds.), *Ecology and management of a Neotropical rainforest:*
1084 *Lessons drawn from Paracou, a long-term experimental research site in French*
1085 *Guiana* (pp. 3–60). Paris: Elsevier.
- 1086 Gourlet-Fleury, S., Guehl, J.-M., & Laroussinie, O. (2004). *Ecology and manage-*
1087 *ment of a Neotropical rainforest: Lessons drawn from Paracou, a long-term*
1088 *experimental research site in French Guiana*. Paris: Elsevier.
- 1089 Grace, J., Mitchard, E., & Gloor, E. (2014, Oct). Perturbations in the carbon bud-
1090 get of the tropics. *Glob. Change Biol.*, 20(10), 3238–3255. doi: 10.1111/gcb.
1091 .12600
- 1092 Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf,
1093 R. T. W., ... McDowell, N. G. (2020, Jun). Plant responses to rising vapor
1094 pressure deficit. *New Phytol.*, 226(6), 1550–1566. doi: 10.1111/nph.16485
- 1095 Gu, L., Norby, R., Haworth, I., Jensen, A., Turner, B., Walker, A., ... Winter, K.

- (2016). *Photosynthetic parameters and nutrient content of trees at the Panama crane sites. 1.0. NGEF Tropics data collection.* Retrieved 12 Sep 2019, from <https://ngt-data.lbl.gov> doi: 10.15486/NGT/1255260
- Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., ... Townshend, J. R. (2015, Mar). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, 1(2), e1500052. doi: 10.1126/sciadv.1500052
- Hancock, S., Armston, J., Hofton, M., Sun, X., Tang, H., Duncanson, L. I., ... Dubayah, R. (2019, Feb). The GEDI simulator: A large-footprint waveform lidar simulator for calibration and validation of spaceborne missions. *Earth Space Sci.*, 6(2), 290–310. doi: 10.1029/2018EA000506
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., ... Townshend, J. R. G. (2013, Nov). High-resolution global maps of 21st-century forest cover change. *Science*, 342(6160), 850–853. doi: 10.1126/science.1244693
- Harris, N. L., Brown, S., Hagen, S. C., Saatchi, S. S., Petrova, S., Salas, W., ... Lotsch, A. (2012, Jun). Baseline map of carbon emissions from deforestation in tropical regions. *Science*, 336(6088), 1573–1576. doi: 10.1126/science.1217962
- Hayek, M. N., Longo, M., Wu, J., Smith, M. N., Restrepo-Coupe, N., Tapajós, R., ... Wofsy, S. C. (2018, Aug). Carbon exchange in an Amazon forest: from hours to years. *Biogeosciences*, 15(15), 4833–4848. doi: 10.5194/bg-15-4833-2018
- Héroult, B., & Piponiot, C. (2018, Feb). Key drivers of ecosystem recovery after disturbance in a neotropical forest. *Forest Ecosyst.*, 5(1), 2. doi: 10.1186/s40663-017-0126-7
- Hess, L. L., Melack, J. M., Affonso, A. G., Barbosa, C., Gastil-Buhl, M., & Novo, E. M. L. M. (2015, Aug). Wetlands of the lowland Amazon Basin: Extent, vegetative cover, and dual-season inundated area as mapped with JERS-1 synthetic aperture radar. *Wetlands*, 35(4), 745–756. doi: 10.1007/s13157-015-0666-y
- Hirano, T., Kusin, K., Limin, S., & Osaki, M. (2015, May). Evapotranspiration of tropical peat swamp forests. *Glob. Change Biol.*, 21(5), 1914–1927. doi: 10.1111/gcb.12653

- 1129 Holdaway, R. J., Sparrow, A. D., & Coomes, D. A. (2010, May). Trends in entropy
1130 production during ecosystem development in the Amazon Basin. *Philos. Trans.
1131 R. Soc. B-Biol. Sci.*, 365(1545), 1437–1447. doi: 10.1098/rstb.2009.0298
- 1132 Hosonuma, N., Herold, M., De Sy, V., De Fries, R. S., Brockhaus, M., Verchot, L.,
1133 ... Romijn, E. (2012, Apr). An assessment of deforestation and forest degrada-
1134 tion drivers in developing countries. *Environ. Res. Lett.*, 7(4), 044009. doi:
1135 10.1088/1748-9326/7/4/044009
- 1136 Huang, M., & Asner, G. P. (2010, Sep). Long-term carbon loss and recovery fol-
1137 lowing selective logging in Amazon forests. *Global Biogeochem. Cycles*, 24(3),
1138 GB3028. doi: 10.1029/2009GB003727
- 1139 Huang, M., Asner, G. P., Keller, M., & Berry, J. A. (2008, Mar). An ecosystem
1140 model for tropical forest disturbance and selective logging. *J. Geophys. Res.-
1141 Biogeosci.*, 113(G1), G01002. doi: 10.1029/2007JG000438
- 1142 Hunter, M. O., Keller, M., Vitoria, D., & Morton, D. C. (2013, Dec). Tree height
1143 and tropical forest biomass estimation. *Biogeosciences*, 10(6), 10491–10529.
1144 doi: 10.5194/bg-10-8385-2013
- 1145 Hurtt, G. C., Dubayah, R., Drake, J., Moorcroft, P. R., Pacala, S. W., Blair, J. B.,
1146 & Fearon, M. G. (2004, Jun). Beyond potential vegetation: combining lidar
1147 data and a height-structured model for carbon studies. *Ecol. Appl.*, 14(3),
1148 873–883. doi: 10.1890/02-5317
- 1149 INMET. (2019, May). *Normais climatológicas do Brasil: 1981-2010*. Retrieved 15
1150 May 2019, from [http://www.inmet.gov.br/portal/index.php?r=clima/
1151 normaisClimatologicas](http://www.inmet.gov.br/portal/index.php?r=clima/normaisClimatologicas) (In Portuguese)
- 1152 IPCC. (2014). *Climate change 2014: impacts, adaptation, and vulnerability. part b:
1153 regional aspects. contribution of working group ii to the fifth assessment report
1154 of the Intergovernmental Panel on Climate Change* (V. R. Barros et al., Eds.).
1155 Cambridge, UK and New York, NY, USA: Cambridge Univ. Press.
- 1156 Jucker, T., Caspersen, J., Chave, J., Antin, C., Barbier, N., Bongers, F., ...
1157 Coomes, D. A. (2017, Jan). Allometric equations for integrating remote
1158 sensing imagery into forest monitoring programmes. *Glob. Change Biol.*, 23(1),
1159 177–190. doi: 10.1111/gcb.13388
- 1160 Jucker, T., Hardwick, S. R., Both, S., Elias, D. M., Ewers, R. M., Milodowski, D. T.,
1161 ... Coomes, D. A. (2018, Nov). Canopy structure and topography jointly con-

- strain the microclimate of human-modified tropical landscapes. *Glob. Change Biol.*, *24*(11), 5243–5258. doi: 10.1111/gcb.14415
- Kapos, V. (1989, May). Effects of isolation on the water status of forest patches in the brazilian amazon. *J. Trop. Ecol.*, *5*(2), 173–185. doi: 10.1017/S0266467400003448
- Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., ... Wirth, C. (2020, Jan). TRY plant trait database — enhanced coverage and open access. *Glob. Change Biol.*, *26*(1), 119–188. doi: 10.1111/gcb.14904
- Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönisch, G., ... Wirth, C. (2011, Sep). TRY – a global database of plant traits. *Glob. Change Biol.*, *17*(9), 2905–2935. doi: 10.1111/j.1365-2486.2011.02451.x
- Kattge, J., Knorr, W., Raddatz, T., & Wirth, C. (2009, Apr). Quantifying photosynthetic capacity and its relationship to leaf nitrogen content for global-scale terrestrial biosphere models. *Glob. Change Biol.*, *15*(4), 976–991. doi: 10.1111/j.1365-2486.2008.01744.x
- Keenan, T. F., & Niinemets, Ü. (2016, Dec). Global leaf trait estimates biased due to plasticity in the shade. *Nat. Plants*, *3*, 16201. doi: 10.1038/nplants.2016.201
- Knox, R. G., Longo, M., Swann, A. L. S., Zhang, K., Levine, N. M., Moorcroft, P. R., & Bras, R. L. (2015, Jan). Hydrometeorological effects of historical land-conversion in an ecosystem-atmosphere model of Northern South America. *Hydrol. Earth Syst. Sci.*, *19*(1), 241–273. doi: 10.5194/hess-19-241-2015
- Koven, C. D., Knox, R. G., Fisher, R. A., Chambers, J., Christoffersen, B. O., Davies, S. J., ... Xu, C. (2019). Benchmarking and parameter sensitivity of physiological and vegetation dynamics using the functionally assembled terrestrial ecosystem simulator (FATES) at Barro Colorado Island, Panama. *Biogeosciences Discuss.*, *2019*, 1–46. (in press) doi: 10.5194/bg-2019-409
- Laybros, A., Schläpfer, D., Féret, J.-B., Descroix, L., Bedeau, C., Lefevre, M.-J., & Vincent, G. (2019, Apr). Across date species detection using airborne imaging spectroscopy. *Remote Sens.*, *11*(7), 789. doi: 10.3390/rs11070789
- Lees, J. M. (2017). RSEIS: Seismic time series analysis tools [Computer software manual]. Retrieved from <https://CRAN.R-project.org/package=RSEIS> (R package version 3.7-4)

- 1195 Lei, Y., Treuhaft, R., Keller, M., dos-Santos, M., Gonçalves, F., & Neumann,
1196 M. (2018, Jun). Quantification of selective logging in tropical forest with
1197 spaceborne SAR interferometry. *Remote Sens. Environ.*, *211*, 167–183. doi:
1198 10.1016/j.rse.2018.04.009
- 1199 Leitold, V., Keller, M., Morton, D., Cook, B., & Shimabukuro, Y. (2015, Feb).
1200 Airborne lidar-based estimates of tropical forest structure in complex terrain:
1201 opportunities and trade-offs for REDD+. *Carbon Balance Manage.*, *10*(1), 3.
1202 doi: 10.1186/s13021-015-0013-x
- 1203 Leitold, V., Morton, D. C., Longo, M., dos-Santos, M. N., Keller, M., & Scaranello,
1204 M. (2018, Aug). El Niño drought increased canopy turnover in Amazon forests.
1205 *New Phytol.*, *219*(3), 959–971. doi: 10.1111/nph.15110
- 1206 Le Page, Y., Morton, D., Bond-Lamberty, B., Pereira, J. M. C., & Hurtt, G. (2015,
1207 Feb). HESFIRE: a global fire model to explore the role of anthropogenic and
1208 weather drivers. *Biogeosciences*, *12*(3), 887–903. doi: 10.5194/bg-12-887-2015
- 1209 Le Page, Y., Morton, D., Hartin, C., Bond-Lamberty, B., Pereira, J. M. C., Hurtt,
1210 G., & Asrar, G. (2017, Dec). Synergy between land use and climate change
1211 increases future fire risk in Amazon forests. *Earth Syst. Dynam.*, *8*(4), 1237–
1212 1246. doi: 10.5194/esd-8-1237-2017
- 1213 Levine, N. M., Zhang, K., Longo, M., Baccini, A., Phillips, O. L., Lewis, S. L., ...
1214 Moorcroft, P. R. (2016, Jan). Ecosystem heterogeneity determines the re-
1215 siliance of the Amazon to climate change. *Proc. Natl. Acad. Sci. U. S. A.*,
1216 *113*(3), 793–797. doi: 10.1073/pnas.1511344112
- 1217 Lewis, S. L., Edwards, D. P., & Galbraith, D. (2015, Aug). Increasing human dom-
1218 inance of tropical forests. *Science*, *349*(6250), 827–832. doi: 10.1126/science
1219 .aaa9932
- 1220 Lloyd, J., Patiño, S., Paiva, R. Q., Nardoto, G. B., Quesada, C. A., Santos, A. J. B.,
1221 ... Mercado, L. M. (2010, Jun). Optimisation of photosynthetic carbon gain
1222 and within-canopy gradients of associated foliar traits for Amazon forest trees.
1223 *Biogeosciences*, *7*(6), 1833–1859. doi: 10.5194/bg-7-1833-2010
- 1224 Lombardozzi, D. L., Smith, N. G., Cheng, S. J., Dukes, J. S., Sharkey, T. D.,
1225 Rogers, A., ... Bonan, G. B. (2018, Jul). Triose phosphate limitation in
1226 photosynthesis models reduces leaf photosynthesis and global terrestrial carbon
1227 storage. *Environ. Res. Lett.*, *13*(7), 074025. doi: 10.1088/1748-9326/aac6f8

- 1228 Longo, M., Keller, M., dos-Santos, M. N., Morton, D., Moorcroft, P. R., Vincent, G.,
1229 ... Saatchi, S. (2020, Feb). Supporting dataset for “Impacts of degradation
1230 on water, energy, and carbon cycling of the Amazon tropical forests”. *Zenodo*.
1231 doi: 10.5281/zenodo.3634131
- 1232 Longo, M., Keller, M., dos Santos, M. N., Leitold, V., Pinagé, E. R., Baccini, A., ...
1233 Morton, D. C. (2016, Nov). Aboveground biomass variability across intact and
1234 degraded forests in the Brazilian Amazon. *Global Biogeochem. Cycles*, *30*(11),
1235 1639–1660. doi: 10.1002/2016GB005465
- 1236 Longo, M., Knox, R., Medvigy, D. M., Levine, N. M., Dietze, M., Swann, A. L. S.,
1237 ... Moorcroft, P. (2019, Aug). Ecosystem Demography model, version 2.2
1238 (ed-2.2). *Zenodo*. doi: 10.5281/zenodo.3365659
- 1239 Longo, M., Knox, R. G., Levine, N. M., Alves, L. F., Bonal, D., Camargo, P. B., ...
1240 Moorcroft, P. R. (2018, Aug). Ecosystem heterogeneity and diversity mitigate
1241 Amazon forest resilience to frequent extreme droughts. *New Phytol.*, *219*(3),
1242 914–931. doi: 10.1111/nph.15185
- 1243 Longo, M., Knox, R. G., Levine, N. M., Swann, A. L. S., Medvigy, D. M., Dietze,
1244 M. C., ... Moorcroft, P. R. (2019, Oct). The biophysics, ecology, and bio-
1245 geochemistry of functionally diverse, vertically and horizontally heterogeneous
1246 ecosystems: the Ecosystem Demography model, version 2.2 – part 2: Model
1247 evaluation for tropical South America. *Geosci. Model Dev.*, *12*(10), 4347–4374.
1248 doi: 10.5194/gmd-12-4347-2019
- 1249 Longo, M., Knox, R. G., Medvigy, D. M., Levine, N. M., Dietze, M. C., Kim, Y., ...
1250 Moorcroft, P. R. (2019, Oct). The biophysics, ecology, and biogeochemistry
1251 of functionally diverse, vertically and horizontally heterogeneous ecosystems:
1252 the Ecosystem Demography model, version 2.2 – part 1: Model description.
1253 *Geosci. Model Dev.*, *12*(10), 4309–4346. doi: 10.5194/gmd-12-4309-2019
- 1254 MacArthur, R. H., & Horn, H. S. (1969, Sep). Foliage profile by vertical measure-
1255 ments. *Ecology*, *50*(5), 802–804. doi: 10.2307/1933693
- 1256 Maréchaux, I., & Chave, J. (2017, Nov). An individual-based forest model to jointly
1257 simulate carbon and tree diversity in Amazonia: description and applications.
1258 *Ecol. Monogr.*, *87*(4), 632–664. doi: 10.1002/ecm.1271
- 1259 Martins, F. S. R. V., Xaud, H. A. M., dos Santos, J. R., & Galvão, L. S. (2012,
1260 Nov). Effects of fire on above-ground forest biomass in the northern Brazilian

- Amazon. *J. Trop. Ecol.*, 28(6), 591–601. doi: 10.1017/S0266467412000636
- Marvin, D. C., Asner, G. P., Knapp, D. E., Anderson, C. B., Martin, R. E., Sinca, F., & Tupayachi, R. (2014, Dec). Amazonian landscapes and the bias in field studies of forest structure and biomass. *Proc. Natl. Acad. Sci. U. S. A.*, 111(48), E5224–E5232. doi: 10.1073/pnas.1412999111
- Mascaro, J., Litton, C. M., Hughes, R. F., Uowolo, A., & Schnitzer, S. A. (2011, Nov). Minimizing bias in biomass allometry: Model selection and log-transformation of data. *Biotropica*, 43(6), 649–653. doi: 10.1111/j.1744-7429.2011.00798.x
- Matsunaga, T., Iwasaki, A., Tsuchida, S., Iwao, K., Tanii, J., Kashimura, O., . . . Tachikawa, T. (2017, Jul). Current status of hyperspectral imager suite (HISUI) onboard international space station (ISS). In *2017 IEEE international geoscience and remote sensing symposium (IGARSS)* (pp. 443–446). doi: 10.1109/IGARSS.2017.8126989
- Mauya, E., Hansen, E., Gobakken, T., Bollandas, O., Malimbwi, R., & Næsset, E. (2015, May). Effects of field plot size on prediction accuracy of aboveground biomass in airborne laser scanning-assisted inventories in tropical rain forests of Tanzania. *Carbon Balance Manage.*, 10(1), 10. doi: 10.1186/s13021-015-0021-x
- Medvigy, D. M., Wofsy, S. C., Munger, J. W., Hollinger, D. Y., & Moorcroft, P. R. (2009, Jan). Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem demography model version 2. *J. Geophys. Res.-Biogeosci.*, 114(G1), G01002. doi: 10.1029/2008JG000812
- Meister, K., Ashton, M. S., Craven, D., & Griscom, H. (2012). Carbon dynamics of tropical forests. In M. S. Ashton, M. L. Tyrrell, D. Spalding, & B. Gentry (Eds.), (pp. 51–75). Dordrecht, Netherlands: Springer Netherlands. doi: 10.1007/978-94-007-2232-3_4
- Mesquita, R. d. C. G., Massoca, P. E. d. S., Jakovac, C. C., Bentos, T. V., & Williamson, G. B. (2015, Aug). Amazon rain forest succession: Stochasticity or land-use legacy? *BioScience*, 65(9), 849–861. doi: 10.1093/biosci/biv108
- Meyer, V., Saatchi, S., Ferraz, A., Xu, L., Duque, A., García, M., & Chave, J. (2019, Mar). Forest degradation and biomass loss along the Chocó region of Colombia. *Carbon Balance Manage.*, 14(1), 2. doi: 10.1186/s13021-019-0117-9

- 1294 Meyer, V., Saatchi, S. S., Chave, J., Dalling, J. W., Bohlman, S., Fricker, G. A., ...
 1295 Hubbell, S. (2013, Aug). Detecting tropical forest biomass dynamics from
 1296 repeated airborne lidar measurements. *Biogeosciences*, 10(8), 5421–5438. doi:
 1297 10.5194/bg-10-5421-2013
- 1298 Miller, A. J. (1984, Nov). Selection of subsets of regression variables. *J. R. Stat. Soc.*
 1299 *A-Gen.*, 147(3), 389–425. doi: 10.2307/2981576
- 1300 Miller, S. D., Goulden, M. L., Hutyrá, L. R., Keller, M., Saleska, S. R., Wofsy, S. C.,
 1301 ... de Camargo, P. B. (2011, Nov). Reduced impact logging minimally alters
 1302 tropical rainforest carbon and energy exchange. *Proc. Natl. Acad. Sci. U. S.*
 1303 *A.*, 108(48), 19431–19435. doi: 10.1073/pnas.1105068108
- 1304 Moorcroft, P. R., Hurtt, G. C., & Pacala, S. W. (2001, Nov). A method for scaling
 1305 vegetation dynamics: The Ecosystem Demography model (ED). *Ecol. Monogr.*,
 1306 71(4), 557–586. doi: 10.1890/0012-9615(2001)071[0557:AMFSVD]2.0
 1307 .CO;2
- 1308 Morton, D. C. (2016, Apr). Forest carbon fluxes: A satellite perspective. *Nature*
 1309 *Clim. Change*, 6(4), 346–348. doi: 10.1038/nclimate2978
- 1310 Morton, D. C., Le Page, Y., DeFries, R. S., Collatz, G. J., & Hurtt, G. C. (2013,
 1311 Jun). Understorey fire frequency and the fate of burned forests in southern
 1312 Amazonia. *Philos. Trans. R. Soc. B-Biol. Sci.*, 368(1619), 20120163. doi:
 1313 10.1098/rstb.2012.0163
- 1314 Ni-Meister, W., Jupp, D., & Dubayah, R. (2001, Sep). Modeling lidar waveforms
 1315 in heterogeneous and discrete canopies. *IEEE T. Geosci. Remote Sens.*, 39(9),
 1316 1943–1958. doi: 10.1109/36.951085
- 1317 Norby, R. J., Gu, L., Haworth, I. C., Jensen, A. M., Turner, B. L., Walker, A. P.,
 1318 ... Winter, K. (2017, Sep). Informing models through empirical relationships
 1319 between foliar phosphorus, nitrogen and photosynthesis across diverse woody
 1320 species in tropical forests of Panama. *New Phytol.*, 215(4), 1425–1437. doi:
 1321 10.1111/nph.14319
- 1322 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven,
 1323 C. D., ... Yang, Z.-L. (2013). *Technical description of version 4.5 of the*
 1324 *community land model (CLM)* (Technical Report Nos. NCAR/TN-503+STR).
 1325 Boulder, CO: NCAR. (420pp.) doi: 10.5065/D6RR1W7M
- 1326 Olivas, P. C., Oberbauer, S. F., Clark, D. B., Clark, D. A., Ryan, M. G., O'Brien,

- 1327 J. J., & Ordoñez, H. (2013, Aug). Comparison of direct and indirect methods
1328 for assessing leaf area index across a tropical rain forest landscape. *Agric. For.*
1329 *Meteorol.*, 177, 110–116. doi: 10.1016/j.agrformet.2013.04.010
- 1330 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell,
1331 G. V. N., Underwood, E. C., ... Kassem, K. R. (2001, Nov). Terrestrial
1332 ecoregions of the world: A new map of life on Earth. *BioScience*, 51(11),
1333 933–938. doi: 10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- 1334 Paracou Portal. (2016, Dec). *Paracou research station, a large-scale forest distur-*
1335 *bance experiment in Amazonia*. Retrieved 31 Jan 2020, from [https://paracou](https://paracou.cirad.fr)
1336 [.cirad.fr](https://paracou.cirad.fr)
- 1337 Pearson, T. R. H., Brown, S., Murray, L., & Sidman, G. (2017, Feb). Greenhouse
1338 gas emissions from tropical forest degradation: an underestimated source. *Car-*
1339 *bon Balance Manage.*, 12(1), 3. doi: 10.1186/s13021-017-0072-2
- 1340 Pellegrini, A. F. A., Franco, A. C., & Hoffmann, W. A. (2016, Mar). Shifts in
1341 functional traits elevate risk of fire-driven tree dieback in tropical savanna and
1342 forest biomes. *Glob. Change Biol.*, 22(3), 1235–1243. doi: 10.1111/gcb.13110
- 1343 Phillips, O. L., van der Heijden, G., Lewis, S. L., López-González, G., Aragão,
1344 L. E. O. C., Lloyd, J., ... Vilanova, E. (2010, Aug). Drought-mortality
1345 relationships for tropical forests. *New Phytol.*, 187(3), 631–646. doi:
1346 10.1111/j.1469-8137.2010.03359.x
- 1347 Pinagé, E. R., Keller, M., Duffy, P., Longo, M., dos Santos, M. N., & Morton, D. C.
1348 (2019, Mar). Long-term impacts of selective logging on Amazon forest dy-
1349 namics from multi-temporal airborne LiDAR. *Remote Sens.*, 11(6), 709. doi:
1350 10.3390/rs11060709
- 1351 Pinto, A., Amaral, P., Souza Jr., C. M., Veríssimo, A., Salomão, R., Gomes,
1352 G., & Balieiro, C. (2009). *Diagnóstico socioeconômico e flore-*
1353 *stal do município de Paragominas* (Technical Report). Belém,
1354 PA, Brazil: Instituto do Homem e Meio Ambiente da Amazônia
1355 (Imazon). (Available at [http://imazon.org.br/publicacoes/](http://imazon.org.br/publicacoes/diagnostico-socioeconomico-e-florestal-do-municipio-de-paragominas/)
1356 [diagnostico-socioeconomico-e-florestal-do-municipio-de-paragominas/](http://imazon.org.br/publicacoes/diagnostico-socioeconomico-e-florestal-do-municipio-de-paragominas/)
1357)
- 1358 Popescu, S. C., Zhao, K., Neuenschwander, A., & Lin, C. (2011, Nov). Satellite lidar
1359 vs. small footprint airborne lidar: Comparing the accuracy of aboveground

- biomass estimates and forest structure metrics at footprint level. *Remote Sens. Environ.*, *115*(11), 2786–2797. doi: 10.1016/j.rse.2011.01.026
- Potapov, P., Hansen, M. C., Laestadius, L., Turubanova, S., Yaroshenko, A., Thies, C., ... Esipova, E. (2017, Jan). The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Sci. Adv.*, *3*(1), e1600821. doi: 10.1126/sciadv.1600821
- Powell, T. L., Galbraith, D. R., Christoffersen, B. O., Harper, A., Imbuzeiro, H. M. A., Rowland, L., ... Moorcroft, P. R. (2013, Oct). Confronting model predictions of carbon fluxes with measurements of Amazon forests subjected to experimental drought. *New Phytol.*, *200*(2), 350–365. doi: 10.1111/nph.12390
- Powers, J. S., & Tiffin, P. (2010, Aug). Plant functional type classifications in tropical dry forests in Costa Rica: leaf habit versus taxonomic approaches. *Funct. Ecol.*, *24*(4), 927–936. doi: 10.1111/j.1365-2435.2010.01701.x
- Priestley, C. H. B., & Taylor, R. J. (1972, Feb). On the assessment of surface heat flux and evaporation using large-scale parameters. *Mon. Wea. Rev.*, *100*(2), 81–92. doi: 10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2
- PRODES-INPE. (2018, Nov). *Taxas anuais de desmatamento*. Retrieved 9 Jan 2019, from <http://www.obt.inpe.br/prodes/index.php> (In Portuguese)
- Purves, D., & Pacala, S. (2008, Jun). Predictive models of forest dynamics. *Science*, *320*(5882), 1452–1453. doi: 10.1126/science.1155359
- Purves, D. W., Lichstein, J. W., Strigul, N., & Pacala, S. W. (2008, Nov). Predicting and understanding forest dynamics using a simple tractable model. *Proc. Natl. Acad. Sci. U. S. A.*, *105*(44), 17018–17022. doi: 10.1073/pnas.0807754105
- Pütz, S., Groeneveld, J., Henle, K., Knogge, C., Martensen, A. C., Metz, M., ... Huth, A. (2014, Oct). Long-term carbon loss in fragmented Neotropical forests. *Nat. Commun.*, *5*. doi: 10.1038/ncomms6037
- Pyle, E. H., Santoni, G. W., Nascimento, H. E. M., Huttyra, L. R., Vieira, S., Curran, D. J., ... Wofsy, S. C. (2008, Mar). Dynamics of carbon, biomass, and structure in two Amazonian forests. *J. Geophys. Res.-Biogeosci.*, *113*(G1), G00B08. doi: 10.1029/2007JG000592
- Quesada, C. A., Miranda, A. C., Hodnett, M. G., Santos, A. J. B., Miranda, H. S., & Breyer, L. M. (2004, Aug). Seasonal and depth variation of soil moisture in

- a burned open savanna (campo sujo) in central Brazil. *Ecol. Appl.*, 14(sp4), 33–41. doi: 10.1890/01-6017
- Quesada, C. A., Phillips, O. L., Schwarz, M., Czimczik, C. I., Baker, T. R., Patiño, S., ... Lloyd, J. (2012, Sep). Basin-wide variations in Amazon forest structure and function are mediated by both soils and climate. *Biogeosciences*, 9(6), 2203–2246. doi: 10.5194/bg-9-2203-2012
- R Core Team. (2019). R: A language and environment for statistical computing [Computer software manual]. Vienna, Austria. Retrieved from <http://www.R-project.org/>
- Rappaport, D., Morton, D., Longo, M., Keller, M., Dubayah, R., & dos-Santos, M. N. (2018, Jun). Quantifying long-term changes in carbon stocks and forest structure from Amazon forest degradation. *Environ. Res. Lett.*, 13(6), 065013. doi: 10.1088/1748-9326/aac331
- Ray, D., Nepstad, D., & Moutinho, P. (2005, Oct). Micrometeorological and canopy controls of fire susceptibility in a forested Amazon landscape. *Ecol. Appl.*, 15(5), 1664–1678. doi: 10.1890/05-0404
- Rödig, E., Cuntz, M., Rammig, A., Fischer, R., Taubert, F., & Huth, A. (2018, Apr). The importance of forest structure for carbon fluxes of the Amazon rain-forest. *Environ. Res. Lett.*, 13(5), 054013. doi: 10.1088/1748-9326/aabc61
- Russo, S., & Kitajima, K. (2016, Mar). The ecophysiology of leaf lifespan in tropical forests: Adaptive and plastic responses to environmental heterogeneity. In G. Goldstein & L. S. Santiago (Eds.), *Tropical tree physiology: Adaptations and responses in a changing environment* (Vol. 6, pp. 357–383). Cham, Switzerland: Springer International Publishing. doi: 10.1007/978-3-319-27422-5_17
- Sabine, C. L., Heimann, M., Artaxo, P., Bakker, D. C. E., Chen, C.-T. A., Field, C. B., ... Valentini, R. (2004). Current status and past trends of the global carbon cycle. In C. B. Field & M. R. Raupach (Eds.), *The global carbon cycle: integrating humans, climate, and the natural world* (Vol. 62, pp. 17–44). Washington, DC, USA: Island Press.
- Santiago, L. S., & Wright, S. J. (2007, Feb). Leaf functional traits of tropical forest plants in relation to growth form. *Funct. Ecol.*, 21(1), 19–27. doi: 10.1111/j.1365-2435.2006.01218.x

- Schimel, D., Schneider, F., & JPL Carbon and Ecosystem Participants. (2019). Flux towers in the sky: global ecology from space. *New Phytol.* (advance online publication) doi: 10.1111/nph.15934
- Schneider, F. D., Kükenbrink, D., Schaepman, M. E., Schimel, D. S., & Morsdorf, F. (2019, Apr). Quantifying 3D structure and occlusion in dense tropical and temperate forests using close-range LiDAR. *Agric. For. Meteorol.*, 268, 249–257. doi: 10.1016/j.agrformet.2019.01.033
- Schneider, F. D., Morsdorf, F., Schmid, B., Petchey, O. L., Hueni, A., Schimel, D. S., & Schaepman, M. E. (2017, Nov). Mapping functional diversity from remotely sensed morphological and physiological forest traits. *Nat. Commun.*, 8(1), 1441. doi: 10.1038/s41467-017-01530-3
- Schwarz, G. (1978, Mar). Estimating the dimension of a model. *Ann. Stat.*, 6(2), 461–464. doi: 10.1214/aos/1176344136
- Sena, E. T., Silva Dias, M. A. F., Carvalho, L. M. V., & Silva Dias, P. L. (2018, Dec). Reduced wet-season length detected by satellite retrievals of cloudiness over Brazilian Amazonia: A new methodology. *J. Climate*, 31(24), 9941–9964. doi: 10.1175/JCLI-D-17-0702.1
- Senior, R. A., Hill, J. K., Benedick, S., & Edwards, D. P. (2018, Mar). Tropical forests are thermally buffered despite intensive selective logging. *Glob. Change Biol.*, 24(3), 1267–1278. doi: 10.1111/gcb.13914
- Shao, G., Stark, S. C., de Almeida, D. R., & Smith, M. N. (2019, Feb). Towards high throughput assessment of canopy dynamics: The estimation of leaf area structure in Amazonian forests with multitemporal multi-sensor airborne lidar. *Remote Sens. Environ.*, 221(221), 1–13. doi: 10.1016/j.rse.2018.10.035
- Silva, C. V. J., Aragão, L. E. O. C., Barlow, J., Espírito-Santo, F., Young, P. J., Anderson, L. O., ... Xaud, H. A. M. (2018, Nov). Drought-induced Amazonian wildfires instigate a decadal-scale disruption of forest carbon dynamics. *Philos. Trans. R. Soc. B-Biol. Sci.*, 373(1760), 20180043. doi: 10.1098/rstb.2018.0043
- Silvério, D. V., Brando, P. M., Bustamante, M. M., Putz, F. E., Marra, D. M., Levick, S. R., & Trumbore, S. E. (2019, Mar). Fire, fragmentation, and windstorms: a recipe for tropical forest degradation. *J. Ecol.*, 107(2), 656–667. doi: 10.1111/1365-2745.13076

- 1459 Souza Jr., C. M., Siqueira, J., Sales, M. H., Fonseca, A., Ribeiro, J., Numat, ...
 1460 Barlow, J. B. (2013, Oct). Ten-year Landsat classification of deforestation and
 1461 forest degradation in the Brazilian Amazon. *Remote Sens.*, 5(11), 5493–5513.
 1462 doi: 10.3390/rs5115493
- 1463 Spracklen, D., Baker, J., Garcia-Carreras, L., & Marsham, J. (2018, Oct). The ef-
 1464 fects of tropical vegetation on rainfall. *Ann. Rev. Environ. Res.*, 43(1), 193–
 1465 218. doi: 10.1146/annurev-environ-102017-030136
- 1466 Stahl, C., Hérault, B., Rossi, V., Burban, B., Bréchet, C., & Bonal, D. (2013,
 1467 Dec). Depth of soil water uptake by tropical rainforest trees during dry
 1468 periods: does tree dimension matter? *Oecologia*, 173(4), 1191–1201. doi:
 1469 10.1007/s00442-013-2724-6
- 1470 Stark, S. C., Leitold, V., Wu, J. L., Hunter, M. O., de Castilho, C. V., Costa,
 1471 F. R. C., ... Saleska, S. R. (2012, Dec). Amazon forest carbon dynamics
 1472 predicted by profiles of canopy leaf area and light environment. *Ecol. Lett.*,
 1473 15(12), 1406–1414. doi: 10.1111/j.1461-0248.2012.01864.x
- 1474 Stavros, E. N., Schimel, D., Pavlick, R., Serbin, S., Swann, A., Duncanson, L., ...
 1475 Wennberg, P. (2017, Jun). ISS observations offer insights into plant function.
 1476 *Nature Ecol. Evol.*, 1, 0194. doi: 10.1038/s41559-017-0194
- 1477 Stovall, A. E., Anderson Teixeira, K. J., & Shugart, H. H. (2018, Nov). Assessing
 1478 terrestrial laser scanning for developing non-destructive biomass allometry.
 1479 *Forest Ecol. Manag.*, 427, 217–229. doi: 10.1016/j.foreco.2018.06.004
- 1480 Strigul, N., Pristinski, D., Purves, D., Dushoff, J., & Pacala, S. (2008, Nov). Scal-
 1481 ing from trees to forests: tractable macroscopic equations for forest dynamics.
 1482 *Ecol. Monogr.*, 78(4), 523–545. doi: 10.1890/08-0082.1
- 1483 Sullivan, M. J. P., Lewis, S. L., Affum-Baffoe, K., Castilho, C., Costa, F., Sanchez,
 1484 A. C., ... Phillips, O. L. (2020, May). Long-term thermal sensitivity of earth’s
 1485 tropical forests. *Science*, 368(6493), 869–874. doi: 10.1126/science.aaw7578
- 1486 Sustainable Landscapes Brazil. (2019, Nov). Retrieved 9 Jan 2019, from [https://](https://www.paisagenslidar.cnptia.embrapa.br/webgis/)
 1487 www.paisagenslidar.cnptia.embrapa.br/webgis/
- 1488 Swann, A. L. S., & Koven, C. D. (2017, Aug). A direct estimate of the seasonal
 1489 cycle of evapotranspiration over the Amazon Basin. *J. Hydrometeor.*, 18(8),
 1490 2173–2185. doi: 10.1175/JHM-D-17-0004.1
- 1491 Swann, A. L. S., Longo, M., Knox, R. G., Lee, E., & Moorcroft, P. R. (2015,

- Dec). Future deforestation in the Amazon and consequences for South American climate. *Agric. For. Meteorol.*, 214–215, 12–24. doi: 10.1016/j.agrformet.2015.07.006
- Tang, H., & Dubayah, R. (2017, Mar). Light-driven growth in Amazon evergreen forests explained by seasonal variations of vertical canopy structure. *Proc. Natl. Acad. Sci. U. S. A.*, 114(10), 2640–2644. doi: 10.1073/pnas.1616943114
- Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L., & Carmona-Moreno, C. (2010, Jun). The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model. *Biogeosciences*, 7(6), 1991–2011. doi: 10.5194/bg-7-1991-2010
- Tyukavina, A., Hansen, M. C., Potapov, P. V., Krylov, A. M., & Goetz, S. J. (2016, Feb). Pan-tropical hinterland forests: mapping minimally disturbed forests. *Global Ecol. Biogeogr.*, 25(2), 151–163. doi: 10.1111/geb.12394
- Tyukavina, A., Hansen, M. C., Potapov, P. V., Stehman, S. V., Smith-Rodriguez, K., Okpa, C., & Aguilar, R. (2017, Apr). Types and rates of forest disturbance in Brazilian Legal Amazon, 2000–2013. *Sci. Adv.*, 3(4), e1601047. doi: 10.1126/sciadv.1601047
- Uhl, C., & Buschbacher, R. (1985, Dec). A disturbing synergism between cattle ranch burning practices and selective tree harvesting in the eastern Amazon. *Biotropica*, 17(4), 265–268. doi: 10.2307/2388588
- Uhl, C., & Kauffman, J. B. (1990, Apr). Deforestation, fire susceptibility, and potential tree responses to fire in the Eastern Amazon. *Ecology*, 71(2), 437–449. doi: 10.2307/1940299
- van der Ent, R. J., Savenije, H. H. G., Schaeffli, B., & Steele-Dunne, S. C. (2010, Sep). Origin and fate of atmospheric moisture over continents. *Water Resour. Res.*, 46(9), W09525. doi: 10.1029/2010WR009127
- VanWey, L. K., D’Antona, A. O., & Brondízio, E. S. (2007, Jan). Household demographic change and land use/land cover change in the Brazilian Amazon. *Popul. Environ.*, 28(3), 163–185. doi: 10.1007/s11111-007-0040-y
- Veríssimo, A., Barreto, P., Mattos, M., Tarifa, R., & Uhl, C. (1992, Dec). Logging impacts and prospects for sustainable forest management in an old Amazonian frontier: The case of Paragominas. *Forest Ecol. Manag.*, 55(1–4), 169–199. doi: 10.1016/0378-1127(92)90099-U

- 1525 Vincent, G., Antin, C., Laurans, M., Heurtebize, J., Durrieu, S., Lavalley, C., &
 1526 Dauzat, J. (2017, Sep). Mapping plant area index of tropical evergreen for-
 1527 est by airborne laser scanning. a cross-validation study using LAI2200 optical
 1528 sensor. *Remote Sens. Environ.*, 198, 254–266. doi: 10.1016/j.rse.2017.05.034
- 1529 von Caemmerer, S. (2000). *Biochemical models of leaf photosynthesis* (No. 2).
 1530 Collingwood, VIC, Australia: CSIRO Publishing. doi: 10.1006/anbo.2000
 1531 .1296
- 1532 von Randow, C., Manzi, A. O., Kruijt, B., de Oliveira, P. J., Zanchi, F. B., Silva,
 1533 R. L., ... Kabat, P. (2004, Jun). Comparative measurements and sea-
 1534 sonal variations in energy and carbon exchange over forest and pasture
 1535 in south west amazonia. *Theor. Appl. Climatol.*, 78(1–3), 5–26. doi:
 1536 10.1007/s00704-004-0041-z
- 1537 Wagner, F., Rossi, V., Stahl, C., Bonal, D., & Hérault, B. (2013, Nov). Asyn-
 1538 chronism in leaf and wood production in tropical forests: a study combining
 1539 satellite and ground-based measurements. *Biogeosciences*, 10(11), 7307–7321.
 1540 doi: 10.5194/bgd-10-8247-2013
- 1541 Walker, W. S., Gorelik, S. R., Baccini, A., Aragon-Osejo, J. L., Josse, C., Meyer,
 1542 C., ... Schwartzman, S. (2020). The role of forest conversion, degradation,
 1543 and disturbance in the carbon dynamics of Amazon indigenous territories and
 1544 protected areas. *Proc. Natl. Acad. Sci. U. S. A.* (advance online publication)
 1545 doi: 10.1073/pnas.1913321117
- 1546 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F.,
 1547 ... Villar, R. (2004, Apr). The worldwide leaf economics spectrum. *Nature*,
 1548 428(6985), 821–827. doi: 10.1038/nature02403
- 1549 Wright, J. S., Fu, R., Worden, J. R., Chakraborty, S., Clinton, N. E., Risi, C., ...
 1550 Yin, L. (2017, Aug). Rainforest-initiated wet season onset over the south-
 1551 ern Amazon. *Proc. Natl. Acad. Sci. U. S. A.*, 114(32), 8481–8486. doi:
 1552 10.1073/pnas.1621516114
- 1553 Wu, M., Schurgers, G., Ahlström, A., Rummukainen, M., Miller, P. A., Smith, B.,
 1554 & May, W. (2017, May). Impacts of land use on climate and ecosystem pro-
 1555 ductivity over the Amazon and the South American continent. *Environ. Res.*
 1556 *Lett.*, 12(5), 054016. doi: 10.1088/1748-9326/aa6fd6
- 1557 Yang, Y., Donohue, R. J., & McVicar, T. R. (2016, Oct). Global estimation of ef-

- 1558 fective plant rooting depth: Implications for hydrological modeling. *Water Re-*
1559 *sour. Res.*, *52*(10), 8260–8276. doi: 10.1002/2016WR019392
- 1560 Zemp, D. C., Schleussner, C.-F., Barbosa, H. M. J., Hirota, M., Montade, V., Sam-
1561 paio, G., ... Rammig, A. (2017, Mar). Self-amplified Amazon forest loss
1562 due to vegetation-atmosphere feedbacks. *Nat. Commun.*, *8*, 14681. doi:
1563 10.1038/ncomms14681
- 1564 Zhang, K., Castanho, A. D. d. A., Galbraith, D. R., Moghim, S., Levine, N., Bras,
1565 R. L., ... Moorcroft, P. R. (2015, Jul). The fate of Amazonian ecosystems
1566 over the coming century arising from changes in climate, atmospheric CO₂ and
1567 land-use. *Glob. Change Biol.*, *21*(7), 2569–2587. doi: 10.1111/gcb.12903

Figure 1.







Figure 2.



3

	Unscaled Property	w_y	e_y
Biomass	19.48	0.28	1.28
Basal Area	33.79	0.25	1.10
Demographic Density	3.287×10^{-2}	0.18	1.51
Leaf area index	4.37	0.29	1.33
	e		1.29





Figure 3.








Figure 4.



- ◇ Gross primary productivity
- ◆ Net ecosystem productivity
- ▽ Evapotranspiration
- ▼ Sensible heat flux
- Outgoing shortwave radiation
- Outgoing longwave radiation
- Paracou (GYF) – Intact
- Belterra (BTE) – Intact
- Tanguro (TAN) – Control
- Tanguro (TAN) – Burned

Figure 5.

Evapotranspiration [mm day⁻¹]









Figure 6.




Figure 7.




Figure 8.







Figure 9.

