

SOLAR ENERGETIC PARTICLE RADIATION DOSAGE NEAR A SIMPLE LUNAR CRATER.

P. H. Phipps^{1,2,3}, T. J. Stubbs², M. D. Looper⁴, H. E. Spence⁵, and L. W. Townsend⁶ ¹Center for Space Sciences and Technology, University of Maryland, Baltimore County, Baltimore, Maryland, USA (email: phhipps@umbc.edu), ²Planetary Magnetospheres Laboratory, NASA/GSFC, Greenbelt, Maryland, USA, ³Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD, USA, ⁴The Aerospace Corporation, El Segundo, CA, USA, ⁵Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, New Hampshire, USA, ⁶Department of Nuclear Engineering, University of Tennessee, Knoxville, TN, USA.

Introduction: The Moon has a harsh radiation environment that poses significant challenges to future science and exploration activities. Exposure hazards from space radiation are primarily due to galactic cosmic rays (GCRs) and solar energetic particles (SEPs) that are incident at the lunar surface from all directions. The Lunar Reconnaissance Orbiter's (LRO) Cosmic Ray Telescope for Effects of Radiation (CRaTER) instrument has been observing space radiation around the Moon since 2009 [1]. The CRaTER observations show a steady GCR flux with intermittent SEP events that have much higher fluxes. During solar minimum GCRs have a higher flux, while SEP events are less common. On the other hand, during solar maximum the SEP events have a higher rate, but the GCR flux is lower. This is due to variations in solar activity. GCRs have characteristic energies spanning from 1 MeV to 10s of GeV [2]. SEPs, however, have much lower energy ranges of 50 keV to 100s of MeV.




Figure 1: Illustration of how natural shielding from surrounding terrain effects radiation exposure at the lunar surface in and around a crater during a solar energetic particle (SEP) event. Inside the crater, the high elevation of the crater walls blocks SEPs incident at shallow angles.

The level of exposure at a given location on the Moon is dependent on the amount of space radiation incident from above the local horizon (Figure 1). This means that radiation dosage depends on the surrounding terrain for any location on the surface, so it can vary substantially from point to point. Here we consider the radiation exposure around simple lunar craters that are representative of the types of landforms that will be encountered by future landed missions (e.g., the

Artemis program) [3]. Of particular concern will be radiation exposure to biological targets, such as astronauts, and to critical electronic systems.

Methods: We use Geant4 Monte Carlo simulations [4] to compute the dose response for spherical targets composed of water (H₂O) and silicon (Si), as proxies for biological and electronic systems, respectively. These targets are surrounded by shells of aluminum of varying thickness to approximate the influence of localized shielding from space suits, rovers, and habitats. To determine the dose rates from SEP proton radiation (e.g., in rads), we convolve the Geant4-computed dose responses with representative SEP spectra for the sum of the October 1989 SEP event [5]. This spectrum has been labeled as best event to use for worst case dose.

To determine the topographical effects, we created a 20 km diameter simple crater, similar to Shackleton Crater at the lunar South Pole. This provides a good proxy for the location of Artemis Base Camp 004, which is planned to be near the rim of Shackleton Crater. Measuring the local horizon for each location on a grid we calculate the solid angle of the visible sky. This fraction can be multiplied by the dose rates determined from the Geant4 convolution with SEP spectrum. This gives the radiation dose received from SEP protons at each surface point.

Discussion and Conclusions: In comparison with dose received on a planar surface, the dose can be reduced by at least 40 percent for regions inside craters, while for areas outside the crater rim it can be reduced by at least 10 percent. Radiation from SEP events are much larger than from the steady flux of GCRs, and occur over much shorter time scales. Thus, acute radiation exposure can be expected to occur during SEP events. Therefore, for protection from SEP events, the shielding effects of surrounding terrain is an important consideration when selecting sites for permanent habitats, as well as for choosing routes and contingency planning during surface operations.

References: [1] Schwadron, N. A., et al. (2018) *Space Weather*, 16, 289–303. [2] Case, A. W., et al. (2013), *Space Weather*, 11, 361–368. [3] NASA's Lunar Exploration Program Overview (Sept 2020) NP-2020-05-2853-HQ. [4] Allison, J., et al. (2006) *IEEE Trans. Nucl. Sci.*, 53 (1), 270–278. [5] Townsend, L.W. et al (2018) *Life Sciences in Space Research*, 17, 32-29