

# Demonstration of Chromospheric Magnetic Mapping with CLASP2.1

David E. McKenzie<sup>1</sup>, Ryohko Ishikawa<sup>2</sup>, Javier Trujillo Bueno<sup>3</sup>, Frederic Auchere<sup>4</sup>, Ken Kobayashi<sup>5</sup>, Amy Winebarger<sup>6</sup>, Ryouhei Kano<sup>7</sup>, Donguk Song<sup>2,5</sup>, Takenori Joten Okamoto<sup>2</sup>, Laurel Rachmeler<sup>8</sup>, Bart De Pontieu<sup>7</sup>, Genevieve Vigil<sup>1</sup>, Luca Belluzzi<sup>9</sup>, Ernest Alsina Ballester<sup>3</sup>, Tanausú del Pino Aleman<sup>3</sup>, Christian Bethge<sup>9</sup>, Taro Sakao<sup>10</sup>, Jiri Stepan<sup>11</sup>

<sup>1</sup>NASA MSFC; <sup>2</sup>NAOJ; <sup>3</sup>IAC; <sup>4</sup>IAS; <sup>5</sup>KASI; <sup>6</sup>NOAA; <sup>7</sup>LMSAL; <sup>8</sup>IRISOL; <sup>9</sup>CU-Boulder; <sup>10</sup>ISAS/JAXA; <sup>11</sup>Academy of Sciences of the Czech Republic



**Introduction:** Probing the magnetic nature of the Sun's atmosphere requires measurement of the Stokes I, Q, U and V profiles of relevant spectral lines (of which Q, U and V encode the magnetic field information). Many of the magnetically sensitive lines formed in the chromosphere and transition region are in the ultraviolet spectrum, necessitating observations above the absorbing terrestrial atmosphere. The Chromospheric Layer Spectro-Polarimeter ("CLASP2") sounding rocket was flown successfully in April 2019, as a follow-on to the successful flight in September 2015 of the Chromospheric Lyman-Alpha Spectro-Polarimeter ("CLASP1"). In October of 2021, we re-flew the CLASP2 experiment with a modified observing program to further demonstrate the maturity of the UV spectropolarimetry techniques, and readiness for development into a satellite observatory. During the reflight, called "CLASP2.1", the spectrograph slit was scanned across an active region plage to acquire a two-dimensional map of Stokes V/I, to demonstrate the ability of UV spectropolarimetry to yield chromospheric magnetic fields over a large area.

**What are the CLASP missions about?** The CLASP1 and CLASP2 missions demonstrated the necessary technology to spectroscopically resolve the linear and circular polarization in UV lines formed in the chromosphere (Figures 1-3). CLASP1 and CLASP2 showed, through the use of the Hanle and Zeeman effects, that magnetic measurements could be achieved in the chromosphere, and that those magnetic fields have spatial variations that relate to – but are not identical to – the variations in the photosphere. *Routine satellite mapping of the spatially varying chromospheric magnetic field will therefore dramatically improve our understanding of the chromosphere and its connection to the rest of the solar atmosphere.*

Although a suborbital space experiment can only provide a few minutes of observing time, the CLASP1 and CLASP2 observations demonstrate the enormous diagnostic potential of spectropolarimetry in the UV spectral region. As clearly shown in Fig. 3, the CLASP2 observations have allowed us to directly determine the longitudinal component of the magnetic field from the low to the very top layers in the chromospheres of active region plages and enhanced network features, at each position along the spatial direction of the spectrograph slit. Figure 3 is essentially a 1-dimensional, multi-height magnetogram of the plage target on 11 April 2019.

To further demonstrate the power of UV spectropolarimetry, and as a test of the maturity and readiness for development into a satellite observatory, we flew CLASP2.1 with the objective to produce a two-dimensional multi-height magnetogram, similar to those commonly displayed for the solar photosphere.

**The CLASP2.1 mission:** With no changes to the instrument since the 2019 flight, CLASP2.1 was launched 08 October 2021 to observe AR 12882. Maintaining the instrument configuration ensured the same wavelength range, spectral resolution, and spatial resolution as the previous investigation of the Mg II lines. Using the SPARCS attitude control system of NASA's Sounding Rocket Program, the spectrograph slit was placed initially at the southern edge of AR 12882, and then subsequently re-targeted the slit 15 times, with separations of slightly less than 2". The result is a sampling of AR 12882 over an area of approximately 200" x 30", significantly expanding the magnetic measurements made by CLASP2. As with the previous flights, co-observations were made in the same target with *Hinode* and *IRIS* satellites. These co-observations are extremely valuable for the further interpretation of the CLASP1/2.1 data. Although the analysis of the CLASP2.1 spectra and Stokes profiles is not yet complete, preliminary examination indicates a fully successful mission. Further reports will be coming soon!

## CLASP2.1




Figure 4: Area of AR 12882 scanned by CLASP2.1. Left-hand panel: slitjaw image with actual locations of the CLASP2.1 slit. A total of 16 slit pointings were achieved. Right-hand panel: The white box shows the planned location of the CLASP2.1 scan, and the black box shows the planned location of IRIS scans, overlaid on an HMI magnetogram of AR 12882.

## CLASP1




Figure 1: A: Lyman- $\alpha$  image taken by the CLASP1 slitjaw camera. B: The Stokes I, Q/I, and U/I profiles of Lyman- $\alpha$  observed by CLASP1 along the slit direction: (a) Stokes I in logarithmic scale; (b) & (c) Stokes-Q/I in two different scales +/- 0.6% & +/- 0.6%, and (d) & (e) the same for Stokes-U/I. The Q/I wings show a clear center-to-limb variation, from -5% close to the limb to -1% near the disk center. The positive reference direction for Stokes Q is parallel to the limb.

## CLASP2




Figure 2: A: CLASP2 slitjaw image of plage on 11 April 2019. B: Orientation of the CLASP2 measurements with respect to *Hinode*/SOT photospheric magnetogram. The spectrograph slit is the faint dark line in the center. C: Mg II k intensity measured by CLASP2. D: Stokes V/I derived from CLASP2 observations of chromospheric plage; key spectral lines are labeled.




Figure 3: A: The longitudinal component of the magnetic field ( $B_{los}$ ) inferred from the Stokes V/I signals shown in Fig. 2. The black and red points show  $B_{los}$  in the middle and upper chromosphere, respectively; they have been determined from the external lobes of Mg II h & k (black points) and from the inner lobes of Mg II h & k (red points). The blue points give  $B_{los}$  in the lower chromosphere, inferred from the Mn I lines. CLASP2 thus measured  $B_{los}$  at three heights in the plage atmosphere. Error bars are  $\pm 1\sigma$ . The green line is the photospheric  $B_{los}$  measured by *Hinode*. This thin black curves show the Mg II k line intensity along the spectrograph slit, at the  $k_3$  line center (solid) and at the  $k_{av}$  emission peak (dash-dotted).

### Important references:

- Belluzzi, L. & Trujillo Bueno, J., 2012 "The Polarization of the Solar Mg II h and k Lines", *ApJ*, 750:L11
- Ishikawa et al, 2021 "Mapping Solar Magnetic Fields from the Photosphere to the Base of the Corona", *Science Advances*, vol.7, issue 8
- Kano, R., Trujillo Bueno, J., Winebarger, A., Auchere, F., et al., 2017 "Discovery of Scattering Polarization in the Hydrogen Ly-alpha Line of the Solar Disk Radiation", *ApJ*, 839(1):L10
- Trujillo Bueno, J., Stepan, J., & Casini, R., 2011 "The Hanle Effect of the Hydrogen Ly-alpha Line for Probing the Magnetism of the Solar Transition Region", *ApJ*, 738:L11

