


Oxidative stress and neuroconsequences of the spaceflight environment – A 5-ion GCRsim dose-response study

A. Ronca, A. Paul, S. Mhatre, J. Iyer, S. Puukila, M. Lowe, S. Tabares-Ruiz, J. Hayes, O. Sui, I. Korosteskiij, M. Semel, M. Dubose, M. Burke, K. Wong, Y. Talyansky, A. Ratri, L.K. Christenson, L. Rubinstein, Y. Shirazi, J. Alwood & C. Tahimic




2022 Human Research Program Investigator's Workshop Enabling the Future: The Pathway to the Moon and Mars (virtual)
Discipline Session 3: Multistressor Hazard Implementation - GCRsim, Hind Limb Unloading, and Isolation, Abstract No. 1105-002458



Artemis and Lunar Gateway



"the first woman and the first person of color" on the Moon



J Women's Health, 2014

Effects of Sex and Gender on Adaptation to Space: Neurosensory Systems

Millard F. Reschke, PhD¹ Helen S. Cohen, EdD² Jody M. Cerisano, BS³ Janine A. Clayton, PhD⁴
Ronita Cromwell, PhD⁵ Richard W. Danielson, PhD² Emma Y. Hwang, PhD³
Candace Tingen, PhD⁶ John R. Allen, PhD⁷ and David L. Tomko, PhD⁸

JOURNAL OF WOMEN'S HEALTH
Volume 23, Number 11, 2014
© Mary Ann Liebert, Inc.
DOI: 10.1089/jwh.2014.4911


Effects of Sex and Gender on Adaptation to Space: Behavioral Health

Namni Goel, PhD¹ Tracy L. Bale, PhD^{1,2} C. Neill Epperson, MD^{1,3} Susan G. Kornstein, MD⁴
Gloria R. Leon, PhD⁵ Lawrence A. Palinkas, PhD⁶ Jack W. Stuster, PhD⁷ and David F. Dinges, PhD¹

Figure Credit: Mark et al., JWH 2014


Central Hypothesis

Spaceflight hazards combine synergistically to trigger oxidative stress that alters immune homeostasis, brain structure and function, and neurobehavioral and cognitive performance.



Mhatre, Iyer et al., Neuroscience and Biobehavioral Reviews Special Issue on Fifty Years of Space Neuroscience as a Prelude to Exploration and Colonization, 2021


HU X Low Dose Radiation



Spaceflight hazards and ground-analog models


Brain and Behavior

- Oxidative damage, increased apoptosis, remodeling of brain microvasculature, increased expression of Nox2 (NADPH oxidase), BBB dysfunction, and increased risk-taking behavior (Bellone et al., 2016; Mao et al., 2016, 2017; Overbey et al., 2019a)
- Increased AQP4 (green) in astrocytes (red) 9mo after return to normal ambulation (Bellone, et al, 2016)
- Space flown mice show similar outcomes in AQP4 and oxidative stress (Mao, et al, 2020)



- Altered pathways in neurogenesis and neuroplasticity, regulation of neuropeptides and cellular signaling revealed by a NASA Transcriptomic analysis (Overbey et al., 2019b)

HU Alone



Spaceflight hazards and ground-analog models

Sensorimotor

- Acute gait impairment (Canu, et al, 1998)
- Inhibition of Righting Reflex (Fox, et al, 1998; Kawano, et al, 2004)
- Somatosensory Cortex adaptation (Dupont, et al, 2010)


Cognitive

- Learning and memory impairments with MWM & Shuttlebox after 14d HU; Cholinergic and antioxidant dysfunction/oxidative damage (Zhang, et al, 2018)
- Learning and memory impairments with NOR (Zhai, et al, 2020; Xu, et al, 2021); Reduced hippocampal theta oscillation; Glutamatergic / GABAergic imbalance (Zhai, et al, 2020; Xu, et al, 2021)

Anxiety & Risk Taking

- Increased anxiety (open field) and risk-taking behavior (Zero Maze); Oxidative damage 9-mo post return to normal ambulation (Bellone, et al, 2016)

Social Isolation Alone



Spaceflight hazards and ground-analog models

- ROS production and oxidative stress, altered brain morphology (hippocampal neuron replacement; remodeling of dendrites and synapses; prefrontal cortical changes; behavioral alterations consistent with the pathogenesis of neurologic and psychiatric diseases (Sorce and Krause, 2009).
- Decreased NK cell activity, reduced CD4⁺ T cells (Hawley and Cacioppo, 2010).
- Weakened BBB → peripheral cytokines cross into CNS (Sylvia and Demas, 2018; Takahashi et al., 2018).
- Chronically elevated sympathetic tone and HPA activation; impaired inflammatory control; immunity; poor sleep quality, and the expression of glucocorticoid-regulating genes → Increased morbidity and mortality rates in older adults (Cacioppo et al., 2011; Holt-Lunstad et. 2015)

Project Aims

Determine ***sex-specific effects*** of single vs. combined:

- Ionizing Radiation (**IR, 5-ion GCRsim**)
- Simulated microgravity (head-down tilt via hindlimb unloading, **(HU)**)
- Social isolation (**SI**)

Crew age-matched male and female mice

→ Assess oxidative stress, immune system, central nervous system (CNS) changes, and behavioral health/cognitive performance

Specific Aim 1 (SA1)

Determine IR dose dependence

Specific Aim 2 (SA2)

Determine combined effects


- IR+HU+SI vs single factor

Specific Aim 3 (SA3)


Test dietary antioxidant countermeasure, Nicotinamide Mononucleotide (NMN)





Overall Study Design




SA1: 5-Ion GCRsim Dose-Response Study (0, 5, 15, or 50cGy)






Mouse Body and Organ Weights





No 5-ion GCRsim Dose Effects
Predictable Sex Differences




Immune: Acute (IR+3) and Intermediate (IR+14)



Cytokine Expression: Intermediate Plasma (IR+14)



Neuroscore Test: Sensorimotor Function (IR+3)



Subtest	Score			
	0	1	2	3
Spontaneous activity (3 min)	No movement	Minimal movement	Touches 1–2 walls	Touches 3–4 walls while standing on hind limbs
Side stroking	No response	Unilateral response	Bilateral weak response or strong ipsilateral response	Strong bilateral response
Vibrissae touch (stroke whiskers)	No response	Unilateral response	Weak bilateral response	Strong bilateral response
Limb extension	Both contralateral; limb completely flexed	One contralateral; limb extended and other flexed	Mid flexion of either contralateral limb	Contralateral forelimbs and hind limbs completely extended
Forepaw outstretching	No movement	Moves in circles	Moves to one side	Straight/curved path
Climbing (1 min)	Weak grip and fall down	Climbs but does not reach top and weak grip	Climbs to top and weak grip or does not climb to top and good grip	Climbs to top and strong grip

- Neurological deficits
- Sensorimotor deficits
- Neuromotor skills
- Reflex test
- Limb use

No 5-ion GCRsim Dose or Sex Differences

Neuroscore Spontaneous Activity (IR+3 & IR+91)




Digging and Rearing - 3min samples



Significant duration differences:

- Female sham vs. 15 cGy digging
- Female sham vs. 5 cGy rearing



Nest-Building: Sensorimotor Function (IR+3)

Deacon Score

- █ 3 Least developed nestlet build
- █ 4
- █ 5 Most developed nestlet build





Deacon Score 3



Deacon Score 4



Deacon Score 5



- females > males
- 50cGy females - no '3's
- 15cGy males < sham


Credit: Siu, Rubinstein, Puukila ¹⁵

Catwalk Gait Analysis: Sensorimotor Function (IR+37)

DELAYED NEUROBIOLOGICAL AND NEUROBEHAVIORAL EFFECTS OF FIVE-ION GCRSIM EXPOSURE IN MALE AND FEMALE MICE

S. Puukila^{1,2}, S. Mhatre^{2,3}, M. Lowe^{2,4}, S. Tabares Ruiz^{2,4}, L. Rubinstein¹, J. Alwood², A. Ronca²
 1Universities Space Research Association, Columbia, MD, USA, 21046, 2Space Biosciences Division, NASA Ames Research Center, Moffett Field, CA, USA, 94035, 3KBR, Houston, TX, USA, 77002, 4Blue Marble Space Institute of Science, Seattle, WA, USA, 98154



2022 NASA Human Research Program Investigators' Workshop



	Sham	5 cGy	15 cGy	50 cGy	
Males	Speed (cm/s)	18.6 ± 2.9	15.4 ± 3.7	17.1 ± 3.8	18.2 ± 2.5
	Regularity Index	61.7 ± 23.1	59.0 ± 21.6	69.5 ± 19.1	68.8 ± 20.9
	Stance Ratio	1.2 ± 0.2	1.1 ± 0.2	1.2 ± 0.3	1.3 ± 0.3
	Base of Support	Front Hind	1.5 ± 0.2 2.8 ± 0.2	1.5 ± 0.3 2.8 ± 0.3	1.4 ± 0.2 2.8 ± 0.2
Females	Speed (cm/s)	31.3 ± 5.2 [#]	29.4 ± 4.6 [#]	29.3 ± 3.1 [#]	28.8 ± 5.6 [#]
	Regularity Index	94.7 ± 7.1 [#]	97.0 ± 3.4 [#]	93.6 ± 6.9 [#]	91.4 ± 12.8 [#]
	Stance Ratio	1.1 ± 0.1	1.0 ± 0.1	1.1 ± 0.1	1.1 ± 0.1 [#]
	Base of Support	Front Hind	1.6 ± 0.2 3.9 ± 0.4 [#]	1.7 ± 0.2 [#] 4.0 ± 0.3 [#]	1.7 ± 0.3 [#] 3.8 ± 0.4 [#]

- Sex differences at all doses
- Females > Males:
Speed, Regularity Index & Base of Support
- No 5-ion GCRsim effects at any dose

Adhesive Removal Test: Sensorimotor Function (IR+63) VNSCOR





5-ion GCRsim Dose x Sex Interactions


Zero Maze: Anxiety and Risk Taking (IR+42)




No 5-ion GCRsim Dose or Sex Differences



Open Field + Novel Object: Memory (IR+84) VNSCOR



No 5-ion GCRsim Dose or Sex Differences



Memory: Barnes Maze (IR+115)



No GCRsim Dose but Sex Differences

- Females faster than males
- Females had fewer errors
- No 5-ion GCRsim dose differed from sham

		Sham	5cGy	15cGy	50cGy
Males	Total Latency (s)				
	Day 1	157.8 ± 40.2	161.5 ± 36.5	157.0 ± 39.8	143.2 ± 46.1
	Day 2	147.1 ± 48.6	143.5 ± 47.0	151.3 ± 40.3	143.9 ± 48.8
	Day 3	153.0 ± 52.1	168.0 ± 28.8	150.8 ± 44.2	164.1 ± 36.1
	Day 4	122.3 ± 63.2*	140.9 ± 53.6	150.1 ± 51.7	152.3 ± 43.1
	Probe	131.0 ± 49.9	118.2 ± 60.1*	122.6 ± 62.5*	142.7 ± 49.1
	Total Errors				
	Day 1	24.5 ± 14.8	26.4 ± 14.9	24.1 ± 11.3	23.3 ± 11.4
	Day 2	25.9 ± 17.1	22.3 ± 13.3	21.5 ± 9.6	23.2 ± 12.5
	Day 3	36.9 ± 20.5	38.7 ± 22.2*	32.8 ± 18.9	32.5 ± 12.8
Females	Total Latency (s)				
	Day 1	161.8 ± 37.4	164.5 ± 37.7	173.5 ± 30.3	157.5 ± 47.3
	Day 2	112.0 ± 58.5	111.0 ± 62.8*	111.9 ± 60.3**	134.7 ± 58.1
	Day 3	102.7 ± 61.1**	95.1 ± 62.1**	98.4 ± 57.4**	112.1 ± 65.3*
	Day 4	77.5 ± 64.0**	86.7 ± 56.7**	116.6 ± 60.4**	112.1 ± 63.0**
	Probe	80.0 ± 46.9**	65.2 ± 42.6**	82.7 ± 56.8**	112.1 ± 63.0**
	Total Errors				
	Day 1	30.7 ± 14.8	28.7 ± 14.6	26.6 ± 14.2	27.1 ± 14.6
	Day 2	20.4 ± 13.3	15.5 ± 13.2	14.6 ± 9.2*	20.0 ± 12.7
	Day 3	21.9 ± 14.9*	20.3 ± 17.1*	22.9 ± 15.3	22.3 ± 17.1
	Day 4	10.8 ± 10.5**	14.0 ± 12.7**	19.5 ± 16.4	18.1 ± 13.9
	Probe	15.7 ± 13.2**	13.2 ± 9.6**	17.6 ± 17.1	16.5 ± 14.5*

= p<.05 vs. males

* = p<.05 vs. Day 1



Summary

- ❑ Early findings from our 5-Ion GCRsim Dose-Response study revealed consistent sex differences in most immune and behavioral analyses, highlighting the importance of male-to-female comparisons.
- ❑ 5-ion GCRsim + isolation produced relatively few effects on immune and behavioral endpoints, although some IR dose effects were observed.
- ❑ Our next steps include:
 - Full analysis of our data, including correlational comparisons, to facilitate selection of the optimal 5-ion GCRsim dose for our upcoming studies.
 - Collaboration with the VNSCOR Team to apply common predictive modeling approaches.