

# CORRELATING THE SEASONAL BEHAVIOR OF POLAR WARMING AND O<sub>2</sub> IR NIGHTGLOW

**A. S. Brecht** ([amanda.s.brecht@nasa.gov](mailto:amanda.s.brecht@nasa.gov)), *NASA Ames Research Center, Moffett Field, CA, USA*, **L. Gkouvelis**, *NASA Postdoctoral Program, NASA Ames Research Center, Moffett Field, CA, USA*, **R. J. Wilson**, **C. E. Harman**, **M. A. Kahre**, *NASA Ames Research Center, Moffett Field, CA, USA*, **A. Kling**, *Bay Area Environmental Research Institute, Moffett Field, CA, USA*.

## Introduction:

Mars' meridional circulation impacts the climate through the transport of heat, water, dust, and trace gases. The meridional circulation in the middle atmosphere is instrumental in the exchange of water between hemispheres and the expansion phase of large regional- and global-scale dust storms. The overall nature and structure of the mean meridional circulation (i.e., the Hadley cell) in the Martian atmosphere has been widely studied but the detailed behavior is still being investigated. To provide a more detailed examination of the meridional circulation, two features driven by the mean circulation will be analyzed as proxies. The first feature is polar warming (PW), which is dynamically induced due to the compressional heating of air in the descending branch of the Hadley cell. The second feature is O<sub>2</sub> IR nightglow emission in the 1.27 micron band, which is the result of a three-body recombination ( $O+O+CO_2 \rightarrow O_2^*+CO_2$ ) and is largely dependent on the transport of dayside produced chemistry. By characterizing their seasonal sensitivities due to known lower atmosphere circulation drivers and correlating them, we will advance our knowledge of the meridional circulation.

## Motivation:

Polar warming is characterized by a reversed (poleward) meridional temperature gradient in the middle-to-high latitudes all year, except local summer. It usually occurs between ~50 km and 80 km and ranges in magnitude from several to tens of Kelvin. It has been observed and discussed since the early 1970's (e.g., Conrath et al., 1973; Conrath et al., 1981; Martin and Kieffer, 1979; Jakosky and Martin, 1987; Deming et al., 1986; Smith et al., 2001; Keating et al., 2003; Tolson et al., 2007; Bougher et al., 2006; McCleese et al., 2007, 2008, 2010; Kleinbohl et al., 2009; McDunn et al., 2013; Forget et al., 2009). McDunn et al. (2013) provides the most quantitative set of parameters of observed PW based upon MRO/MCS-derived atmospheric temperatures. The study identifies PW trends that suggest that the structure of the mean meridional circulation is complex and not fully understood. Such trends are: (a) the southern hemisphere annual maximum PW occurs during late local winter of Martian year (MY) 29 and MY 30, (b) southern winter and northern winter solstices have similar PW magnitudes, and (c) the equinoctial seasons demonstrated hemispheric asymmetry in PW magnitude (McDunn et al., 2013).

The O<sub>2</sub> IR nightglow emission is a product of 3-body reaction that occurs in areas of converging circulation. The nightglow becomes a valid tracer of circulation due to the chemical sources (atomic oxygen) largely produced on the dayside through photolysis (CO<sub>2</sub> and O<sub>3</sub>). Chemical species (CO<sub>2</sub>, O<sub>3</sub>, O, etc.) are then transported by the circulation (day to night; equator to pole) to a region of favorable condition for recombination and subsequent de-excitation (i.e., emission). The O<sub>2</sub> IR nightglow emission at Mars produces a lower intensity compared to Venus and Earth, however it has been observed by orbital missions; Mex/OMEGA (Bertaux et al., 2012), Mex/SPICAM (Fedorova et al., 2012), and MRO/CRISM (Clancy et al., 2012, 2013). As with the PW, the nightglow is observed over high northern and southern latitudes between 40 – 60 km in altitude. The observational trends of the vertically integrated emission are: (1) the northern winter solstice has more intense emission than the southern winter solstice, and (2) the lowest peak altitude occurs during northern winter solstice. These trends seem to support the general concept of the role of the mean meridional circulation, unlike some of the PW trends (Clancy et al., 2012).

Past numerical modeling has been focused on either PW or the O<sub>2</sub> IR nightglow, but not both. In general, numerical models seem to under-predict the PW and over-predict the O<sub>2</sub> IR nightglow emission in comparison to observations (e.g., Haberle et al., 1982; Barnes and Haberle, 1996; Wilson, 1997; Forget et al., 1999; Clancy et al. 2012, 2013).

The observations are suggesting these two features, which are driven by the same circulation and are co-located, respond to the mean meridional circulation differently with respect to seasons. We will investigate the seasonal behavior of these features with respect to different atmospheric conditions and evaluate the trends.

## Methods:

The current NASA Ames Mars Global Climate Model (MGCM), as described in Kahre et al., (this meeting), is utilized to characterize the seasonal response of PW and the O<sub>2</sub> IR nightglow emission to different dynamical drivers and their correlations simultaneously. Examining the drivers of PW and O<sub>2</sub> IR nightglow emission will provide a deeper understanding of the meridional circulation and how it impacts transport of other elements (e.g., water, dust). The work shown here uses a single MGCM set up. We




Figure 1: Polar warming for cloud case. Night time (22-2 LT), zonal average, binned  $10^\circ L_s$ ,  $5^\circ \text{Lat}$ .

use a background dust scenario, which is constructed from the Montabone scenarios (MY29-34) by identifying the spatially and seasonally varying minimum value of the multi-year record to filter out the influence of episodic regional dust storms. For this specific application, the dust scenario is zonally averaged. The vertical dust distribution is governed by a latitudinally and seasonally varying prescription similar to Forget et al. (1999). The water ice clouds are radiatively active and solar flux is set for solar moderate conditions. Orographic and non-orographic gravity waves are not enabled for this simulation.

The simulated fields are binned in  $L_s$  ( $10^\circ$ ) and latitude ( $5^\circ$ ), zonally averaged, specifically looking at nightside (local time = 22-02).

The polar warming magnitude is quantified as described in McDunn et al., 2013:

$\Delta_p T(\text{Lat}) = [T_{\text{Lat}} - T_2]_p$ ;  $\text{Lat}_{T2} < \text{Lat} \leq T_1$ , where  $T_1$  is the largest bin-averaged temperatures in each hemisphere,  $T_2$  is the smallest bin-averaged temperature between  $\text{Lat}_{T1}$  and the equator,  $\text{Lat}_{T1}$  is the latitude of  $T_1$  and  $\text{Lat}_{T2}$  is the latitude of  $T_2$ . Following McDunn et al. (2013) temperature enhancement larger than 15 K is used to identify PW.

The  $\text{O}_2$  IR nightglow volume emission rate (VER) is calculated using the following expression:

$E(\text{O}_2^*) = (a \cdot k_1 \cdot [\text{O}] \cdot [\text{O}] \cdot [\text{CO}_2]) / (1 + t \cdot (k_{\text{co}2} \cdot [\text{CO}_2]))$  where  $a$  is the total yield of  $\text{O}_2^*$ ,  $k_1$  is the rate coefficient for the three-body recombination reaction,  $[\text{O}]$

and  $[\text{CO}_2]$  are concentrations for each specie,  $t$  is the lifetime by radiative relaxation of  $\text{O}_2^*$ ,  $k_{\text{co}2}$  is the quenching coefficient for  $\text{CO}_2$ . Current values for this simulation are:  $a = 0.75$ ,  $k_1 = 2.5 \cdot 5.2 \cdot 10^{-35} \cdot \exp(900/T) \text{ cm}^6 \text{ s}^{-1}$ ,  $t = 3800 \text{ s}$ ,  $k_{\text{co}2} = 2.0 \cdot 10^{-20} \text{ cm}^3 \text{ s}^{-1}$ . Some of these values are not well known, so sensitivity tests will need to be conducted.

### Preliminary Results:

**Polar Warming.** Figure 1 demonstrates the nightside polar warming calculated from the MGC. For this simulation,  $L_s = 90^\circ$  (Fig. 1 top) has minimal ( $\sim 15 \text{ K}$  in northern hemisphere) to no PW. The MCS derived PW for  $L_s = 90^\circ$  was less than 15 K during MY 29 in both hemispheres. However, MY 30 had a maximum of  $\sim 30 \text{ K}$  in the southern hemisphere and  $\sim 15-20 \text{ K}$  in the northern hemisphere (McDunn et al. 2013).




Figure 2:  $\text{O}_2$  IR nightglow volume emission rate for cloud case. Night time (22-2 LT), zonal average, binned  $10^\circ L_s$ ,  $5^\circ \text{Lat}$ .

The northern winter solstice ( $L_s = 270^\circ$ ; Fig. 1 bottom) has PW magnitudes as large as 70 K in the northern hemisphere and  $\sim 20 \text{ K}$  in the southern hemisphere. The MCS derived PW for  $L_s = 270^\circ$  was largest overall in MY 28 with the maximum northern hemisphere  $\sim 50 \text{ K}$  and southern hemisphere  $\sim 30 \text{ K}$ . This Mars year also had a solstitial-season global dust storm. Both MY 29 and MY 30 resulted in northern hemisphere PW  $\sim 30 \text{ K}$  and southern hemisphere PW  $\sim 15 \text{ K}$  (McDunn et al., 2013).

This general trend of the simulated larger PW near

northern winter solstice compared to southern winter solstice is consistent with the expected seasonal variation in the strength and extent of the mean meridional




Figure 3: Maximum values of polar warming and log O<sub>2</sub> IR nightglow volume emission rate with respect to solar longitude ( $L_s$ ). Night time (22-2 LT), zonal average, binned  $10^\circ L_s$ ,  $5^\circ$  Lat. [Top] Northern hemisphere and [Bottom] Southern hemisphere.

circulation. However, the simulated behavior of the northern hemisphere PW at  $L_s=90^\circ$  being larger than the southern hemisphere is not expected.

**O<sub>2</sub> IR Nightglow Emission.** Figure 2 demonstrates the simulated nightside O<sub>2</sub> IR nightglow VER. This simulation demonstrates the nightglow to be largest in the winter hemisphere with respect to solar longitude. The maximums are similar in magnitude for the southern winter solstice ( $L_s=90^\circ$ ; Fig. 2, top) and northern winter solstice ( $L_s=270^\circ$ ; Fig. 2, bottom) which would seem counter to what would be expected. The mean meridional circulation is observed to be stronger during  $L_s = 270^\circ$ , as the PW magnitudes in Figure 1 supports. Further examination and results will be compared to Clancy et al. 2012 and 2013, which discusses the MARCI CRISM O<sub>2</sub> IR nightglow emission observations.

**Correlation.** As discussed above, these features are driven by the same circulation and have been observed in similar spatial regions. Figure 3 shows how the simulated maximum polar warming and O<sub>2</sub> IR VER vary seasonally; with the top panel representing the northern hemisphere and the bottom panel representing the southern hemisphere. In the north, the O<sub>2</sub> IR VER has a sinusoidal pattern, with a minimum

near  $L_s \sim 100^\circ$  and a maximum near  $L_s \sim 270^\circ$ . PW has a similar sinusoidal pattern but with an extra sizable decrease  $L_s \sim 205^\circ$ . The southern hemisphere shows the two features trending in opposite directions around  $L_s = 90^\circ$  and then both trending toward a minimum near  $L_s = \sim 290^\circ$ . Clancy et al. (2012, 2013) presented the correlation between CRISM observed O<sub>2</sub> IR nightglow emission and MCS temperatures for  $L_s = 74-137^\circ$ , Lat= $70-90^\circ$  S, and altitude range of 50-60 km. They found the two features to be anticorrelated, as did the GCM results within that study. They claim, based on the southern hemisphere observations, that the correlation with adiabatic heating with O<sub>2</sub> IR nightglow emission is not very strong, but it seems to be largely dependent on the temperature dependent rate constant.

### Future Work:

We will examine the behavior of the PW and O<sub>2</sub> IR nightglow VER and correlate their seasonal behaviors to provide more knowledge on the meridional circulation. We will extend the work of Clancy et al. (2012, 2013) by studying the full Martian year. The thermal and wind field, along with mass stream functions will also be reviewed. Moreover, several sensitivity cases will be conducted to investigate the relative importance of thermal (clouds and dust) and wave-induced forcing on the correlation between PW and O<sub>2</sub> IR nightglow VER, thus the meridional circulation.

### References:

- Barnes and Haberle, *J. Atmos. Sci.*, 53, 1996; Bertaux et al., *J. Geophys. Res.*, 117, 2012; Bougher et al., *Planet. Space Sci.*, 54, 2006; Clancy et al., *J. Geophys. Res.*, 117, 2012; Clancy et al., *J. Geophys. Res.*, 118, 2013; Conrath et al., *J. Geophys. Res.*, 78, 1973; Conrath et al., *Icarus*, 24, 1981; Deming et al., *Icarus*, 66, 1986; Fedorova et al., *Icarus*, 219, 2012; Forget et al., *J. Geophys. Res.*, 104, 1999; Forget et al., *J. Geophys. Res.*, 114, 2009; Haberle et al., *Icarus*, 50, 1982; Jakosky and Martin, *Icarus*, 72, 1987; Keating et al., *Third International Mars Polar Science Conference*, 8033, 2003; Kleinbohl et al., *J. Geophys. Res.*, 114, 2009; Martin and Kieffer, *J. Geophys. Res.*, 84, 1979; McCleese et al., *J. Geophys. Res.*, 112, 2007; McCleese et al., *Nature Geoscience*, 1, 2008; McCleese et al., *J. Geophys. Res.*, 115, 2010; McDunn et al., *J. Geophys. Res. Planets*, 118, 2013; Smith et al., *J. Geophys. Res. Planets*, 106, 2001; Tolson et al., *J. Spacecraft and Rockets*, 44, 2007; Wilson, *Geophys. Res. Lett.*, 24, 1997