

# **Comparative Study on a Variety of High Order Spatial Discretizations with the Entropy Split Methods for MHD**

**H.C. Yee, NASA Ames Research Center**  
*(Joint work: B. Sjogreen)*

**ECCOMAS-2022**  
**June 5-9, 2022, Oslo, Norway**

# Objectives

- Present Our **Recent Development of Entropy Split Methods (ES)** for the **Ideal MHD**
  - > Two ES methods are proposed
- **Comparison** of **8<sup>th</sup>-Order Entropy Split Methods** with **Six** Other **8<sup>th</sup>-Order Methods**
  - > Entropy Conserving Methods
  - > Momentum Conserving Method
  - > Kinetic Energy Preserving Method
  - > Combination of Two of These Properties Methods

# Motivation

- **Improvement in Nonlinear Stability (reduce aliasing errors)**
  - > More efficient than Tadmor-type entropy conserving methods of same order
  - > Methods cater to **long time integration** of compressible turbulence
- **Improvement in Accuracy (for a wide spectrum of flow speeds)**
  - > **Efficient Nonlinear Filter** methods with adaptive local **flow sensors** designed to **minimize** the use of numerical dissipation
- **Mass Conservation Issue for shock-free & flows with shocks**
  - > Entropy split method is entropy conserving but it is not in conservation form
  - > Two proposed methods to overcome the issue

## Comparison Among Seven 8<sup>th</sup>-Order Methods

- **ECLOG**: Tadmor-type entropy conserving method using the Tadmor entropy function  $E_L = -\rho \log(p\rho^{-\gamma})$ .
- **ECLOGKP**: Tadmor-type entropy conserving method using the Tadmor entropy function with Ranocha's kinetic energy preserving modification [32].
- **ECHKP**: Entropy conserving using the Harten class of entropy functions  $E_H = -\frac{\gamma+\alpha}{\gamma-1}\rho(p\rho^{-\gamma})^{\frac{1}{\alpha+\gamma}}$  [50, 23]. In its base form also satisfies Ranocha's kinetic energy preservation condition (KEP); only one variant for this method [43, 32].
- **KGP**: Kennedy-Gruber-Pirozzoli (KGP) skew-symmetric splitting of the inviscid flux derivative that is kinetic energy preserving [26, 31, 13].

## Comparison Among **Seven** 8<sup>th</sup>-Order Methods (*Cont.*)

- **ESSW**: Entropy split with Ducros et al. splitting but switch to regular central near discontinuities [46].
- **DSKP**: Ducros Split with KEP.
- **DS**: Momentum conserving Ducros et al. skew-symmetric split of the inviscid flux derivative [14].

# Treatment of Discontinuities for Entropy Split Schemes

Accurate schemes developed for **short** time integration (or rapidly developing flows) usually **SUFFER** from **nonlinear instability for long time integration**

- Standard shock-capturing methods are too **diffusive** for long time integration
- **Careful** design of appropriate nonlinear numerical dissipations with flow sensors can **improve accuracy**

## Yee et al. Nonlinear Filter Approach:

- **High order entropy split schemes as the base scheme**
- **Local flow sensor to examine the regularity of the computed data after each full time step of the base scheme.**
- **Determine the location and amount of numerical dissipation needed**
- **Nonlinearly filter the computed data at these locations only**

# Preliminary

$$\mathbf{U}_t + \mathbf{F}(\mathbf{U})_x = 0$$

Entropy function:  $E(\mathbf{U})$

Entropy variables:  $\mathbf{W} = \nabla_{\mathbf{U}} E$  (require  $E_{\mathbf{U},\mathbf{U}} > 0$ )

Entropy flux:  $f = f(\mathbf{U})$

Entropy conservation if

$$\mathbf{W}^T \frac{\partial \mathbf{F}}{\partial \mathbf{U}} = \nabla_{\mathbf{U}} f$$

Take conservation law & multiply by  $\mathbf{W}$  to give

$$\mathbf{W}^T \mathbf{U}_t + \mathbf{W}^T \mathbf{F}_x = 0 \quad \rightarrow \quad E_t + \mathbf{W}^T \frac{\partial \mathbf{F}}{\partial \mathbf{U}} \mathbf{U}_x = 0 \quad \rightarrow \quad E_t + f_x = 0$$

Alternatively, if  $f$  is known, denote  $\psi = \mathbf{W}^T \mathbf{F} - f$  and require

$$\psi_x = \mathbf{W}_x^T \mathbf{F}.$$

Entropy conservation shown by

$$\mathbf{W}^T \mathbf{F}_x = (\mathbf{W}^T \mathbf{F})_x - \mathbf{W}_x^T \mathbf{F} = (\mathbf{W}^T \mathbf{F})_x - (\mathbf{W}^T \mathbf{F} - f)_x = f_x$$

# Entropy Conserving Discretization

Numerical flux:  $\mathbf{H}_{j+1/2}$

Entropy numerical flux:  $\mathbf{h}_{j+1/2}$

Semi-discretized

$$\frac{d}{dt}\mathbf{U}_j + \frac{1}{\Delta x}(\mathbf{H}_{j+1/2} - \mathbf{H}_{j-1/2}) = \mathbf{0}$$

If

$$(\mathbf{W}_{j+1} - \mathbf{W}_j)^T \mathbf{H}_{j+1/2} = \psi_{j+1} - \psi_j$$

Then can derive discrete entropy conservation law

$$\frac{d}{dt}E_j + \frac{1}{\Delta x}(h_{j+1/2} - h_{j-1/2}) = 0$$

with

$$h_{j+1/2} = \frac{1}{2}(\mathbf{W}_{j+1} + \mathbf{W}_j)^T \mathbf{H}_{j+1/2} - \frac{1}{2}(\psi_{j+1} + \psi_j)$$

[1] E.Tadmor, Math. Comput., 43(1984) pp. 369–381.

# Local & Global Entropy Stability

Local entropy conservation:

$$\frac{d}{dt}E_j + \frac{1}{\Delta x}(h_{j+1/2} - h_{j-1/2}) = 0$$

Local entropy stability:

$$\frac{d}{dt}E_j + \frac{1}{\Delta x}(h_{j+1/2} - h_{j-1/2}) \leq 0$$

Global entropy conservation:

$$\frac{d}{dt} \sum_j \Delta x E_j = 0$$

Global entropy stability:

$$\frac{d}{dt} \sum_j \Delta x E_j \leq 0$$

**Entropy Splitting of the Euler Flux Derivatives  
Leads to Global Entropy Conservation (Stability) by the Energy Norm**

# Entropy Split Methods (ECHKP)

Entropies considered by Harten (1983)

$$E = -\rho(p\rho^{-\gamma})^{1/(\alpha+\gamma)}$$

Symmetrizable & Homogeneous property of the Euler Flux lead to

$$\mathbf{F}_w \mathbf{W} = \beta \mathbf{F} \quad (1)$$

for  $\beta = \frac{\alpha+\gamma}{1-\gamma}$ . Use the split form with  $\beta \neq -1$

$$\mathbf{F}_x = \frac{\beta}{\beta+1} \mathbf{F}_x + \frac{1}{\beta+1} \mathbf{F}_w \mathbf{W}_x$$

in a semi-discrete approximation:

$$\frac{d}{dt} \mathbf{U}_j + \frac{\beta}{\beta+1} D \mathbf{F}_j + \frac{1}{\beta+1} (\mathbf{F}_w)_j D \mathbf{W}_j = 0$$

where  $D$  is a SBP difference operator.

**ECHKP:** *Entropy Split Methods automatically satisfies the Ranocha's kinetic energy preservation condition*

## Entropy Splitting Leads to Global Entropy Conservation & Stability

Form the SBP scalar product by  $\mathbf{W}_j^T$  to obtain the entropy

$$\frac{d}{dt} \sum_j \omega_j \Delta x E_j + \frac{\beta}{\beta + 1} (\mathbf{W}, D\mathbf{F}) + \frac{1}{\beta + 1} (\mathbf{W}, \mathbf{F}_W D\mathbf{W}) = 0$$

( $\omega_j$  are SBP norm weights) where SBP and property (1) give

$$\begin{aligned} \beta(\mathbf{W}, D\mathbf{F}) &= -\beta(D\mathbf{W}, \mathbf{F}) - \beta\mathbf{W}_1^T \mathbf{F}_1 + \beta\mathbf{W}_N^T \mathbf{F}_N = \\ &\quad - (D\mathbf{W}, \mathbf{F}_W \mathbf{W}) - \beta\mathbf{W}_1^T \mathbf{F}_1 + \beta\mathbf{W}_N^T \mathbf{F}_N \end{aligned}$$

and symmetry of  $\mathbf{F}_W$  gives  $(\mathbf{W}, \mathbf{F}_W D\mathbf{W}) = (\mathbf{F}_W \mathbf{W}, D\mathbf{W})$ , so that

$$\frac{d}{dt} \sum_j \omega_j \Delta x E_j = \frac{\beta}{\beta + 1} (\mathbf{W}_1^T \mathbf{F}_1 - \mathbf{W}_N^T \mathbf{F}_N)$$

i.e., global entropy conservation with boundary contributions.

Note, property (1) does not hold for  $E = -\rho \log(p\rho^{-\gamma})$  commonly used in the literature

# Entropy Splitting of Euler Flux Derivatives

(Oliger & Olsson, Olsson, Gerittsen & Olsson, Yee et al. (1994-1999))

Entropy stable (generalized energy norm) with SBP for Central Schemes

Physical Entropy, Homogeneous of Degree One of Euler Flux Derivative  $F_x$

$$\underline{F}_x = \frac{\beta}{\beta + 1} \underline{F}_x + \frac{1}{\beta + 1} \underline{F}_w W_x, \quad \beta \neq -1$$

$$\begin{aligned} W &= [w_1, w_2, w_3, w_3, w_5]^T \\ &= \frac{p^*}{p} \left[ e + \frac{\alpha - 1}{\gamma - 1} p, -\rho u, -\rho v, \rho w, \rho \right]^T, \end{aligned}$$

where

$$p^* = -(p\rho^{-\gamma})^{\frac{1}{\alpha+\gamma}}$$

and

$$\beta = \frac{\alpha + \gamma}{1 - \gamma}, \quad \alpha > 0 \text{ or } \alpha < -\gamma. \quad \text{Negative Beta} \rightarrow \text{unphysical}$$

# Entropy Conserving Discretization Tadmor-type (ECLOG) with Boundaries

**Entropy Split Schemes:** Achieving global entropy conservation & stability automatically

by standard SBP & energy norm estimate (periodic or non-periodic BC)

**Entropy Conserving Schemes - Tadmor-type:** Achieving global entropy conservation & stability  
require generalized SBP & dissipation

Domain  $j = 1, 2, \dots$ , with boundary at  $j = 1$ . Consider the scheme

$$\frac{d}{dt} \mathbf{U}_j + \frac{1}{\Delta x} (\mathbf{H}_{j+1/2}^{EC} - \mathbf{H}_{j-1/2}^{EC}) = \mathbf{0}$$

with the E.C. 2nd-order numerical flux  $\mathbf{H}_{j+1/2}^{EC} = \mathbf{H}^{EC}(\mathbf{U}_j, \mathbf{U}_{j+1})$ , defined at  $j = 1, 2, \dots$ ,

Define  $\mathbf{H}_{1/2}^{EC} = 2\mathbf{F}_1 - \mathbf{H}_{3/2}^{EC}$ , then the local entropy conservation law

$$\frac{d}{dt} E_j + \frac{1}{\Delta x} (h_{j+1/2} - h_{j-1/2}) = 0$$

holds for  $j = 1, 2, \dots$ , with

$$h_{1/2} = \mathbf{W}_1^T \mathbf{F}_1 - \psi_1$$

and

$$h_{j+1/2} = \frac{1}{2} (\mathbf{W}_{j+1} + \mathbf{W}_j)^T \mathbf{H}_{j+1/2} - \frac{1}{2} (\psi_{j+1} + \psi_j)$$

for  $j \geq 1$  (as previously).

# Entropy Conserving Discretization of Tadmor-type (ECLOG) with Boundaries (Cont.)

The global entropy conservation with boundaries

$$\Delta x \frac{d}{dt} \left( \frac{1}{2} E_1 + \sum_{j=2} E_j \right) + \mathbf{W}_1^T \mathbf{F}_1 - \psi_1 = 0$$

follows, where  $1/2$  is the SBP weight of the boundary entropy, and  $f_1 = \mathbf{W}_1^T \mathbf{F}_1 - \psi_1$  is the boundary entropy flux.

For a construction that generalizes to higher order accuracy, see:

Parsani, Carpenter, Nielsen, J.Comput.Phys., 292(2015),pp. 88–113.

# Advantage of Entropy Split Method over Standard Tadmor-Type Entropy Conserving Methods

**Tadmor-type entropy conserving method** (logarithmic entropy or physical entropy)  
**requires 2 times CPU** than entropy split method

+

**Both Approaches Exhibit Similar Accuracy & Stability**  
(Same order of central discretization)

# Mass Conservation issue by Entropy Split Methods

**Fact:** Entropy splitting of the Euler inviscid flux derivatives is not in conservation form

- > ES method in its original form is entropy conservative but is not a conservative method
- > Mass conservation is not a problem on smooth flow test cases
- > Mass conservation error increases as a function of the duration of time integration for flows with discontinuities

**Two Proposed New Split Methods to Correct Mass Conservation Error:**

*Sjogreen & Yee, “Construction of Conservative Numerical Fluxes for the Entropy Split Method”, HONOM-2019 special CAMC volume, 2022*

**Test Cases on the new approach with improve stability and accuracy:**

- > 2D Vortex Convection
- > 3D Taylor-Green Vortex
- > 3D Isotropic Turbulence with Shocklets
- > 1D Shu Osher Shock-Turbulence
- > 2D Alfven Wave
- > Orzag-Tang MHD Vortex
- > Brio-Wu MHD Shock-Tube

# Kennedy & Gruber Splitting (2008)

Split the derivative of triple products, kinetic energy preservation

$$\frac{\partial}{\partial x}(abc) = \alpha \frac{\partial}{\partial x}(abc) + \beta \left[ a \frac{\partial}{\partial x}(bc) + bc \frac{\partial}{\partial x}(a) \right] + \kappa \left[ b \frac{\partial}{\partial x}(ac) + ac \frac{\partial}{\partial x}(b) \right] + \delta \left[ c \frac{\partial}{\partial x}(ab) + ab \frac{\partial}{\partial x}(c) \right] \\ + \epsilon \left[ bc \frac{\partial}{\partial x}(a) + ac \frac{\partial}{\partial x}(b) + ab \frac{\partial}{\partial x}(c) \right],$$

$$\text{where } \epsilon = 1 - \alpha - \beta - \kappa - \delta$$

**Common parameter:**

$$\alpha = \beta = \kappa = \delta = \frac{1}{4} \text{ and } \epsilon = 0,$$

$$(abc)_x = \frac{1}{4} (abc)_x + \frac{1}{4} (a_x (bc) + b_x (ac) + c_x (ab)) + \frac{1}{4} (a (bc)_x + b (ac)_x + c (ab)_x)$$

# Dispersion Relation-Preserving (DRP) Schemes

**Three DRP schemes in CAA:** **DRP4S7** (Tam & Webb), **DRP4S9**, **ST09** (Bogey & Bailly)

$$Du_j = \frac{1}{h} \sum_{k=1}^q a_k (u_{j+k} - u_{j-k})$$

## DRP4S7

Coefficients of DRP4S7, optimized over  $[0, 1.1]$ .

| $k$ | $a_k$                |
|-----|----------------------|
| 1   | 0.77088238051822552  |
| 2   | -0.16670590441458047 |
| 3   | 0.02084314277031176  |

## DRP4S9

Coefficients of DRP4S9, optimized over  $[\pi/16, \pi/2]$ .

| $k$ | $a_k$              |
|-----|--------------------|
| 1   | 0.846863763009931  |
| 2   | -0.251240526849904 |
| 3   | 0.063181723773749  |
| 4   | -0.008481970157843 |

## ST09

Coefficients of ST09, optimized over  $[\pi/16, \pi/2]$ , from [18].

| $k$ | $a_k$              |
|-----|--------------------|
| 1   | 0.841570216389881  |
| 2   | -0.244678789340406 |
| 3   | 0.059463699920073  |
| 4   | -0.007650934367322 |

# 3-D Compressible MHD (Ideal)

$$\begin{pmatrix} \rho \\ \rho u \\ \rho v \\ \rho w \\ e \\ B_x \\ B_y \\ B_z \end{pmatrix}_t + \operatorname{div} \begin{pmatrix} \rho \mathbf{u} \mathbf{u}^T + (p + \frac{1}{2} B^2) \mathbf{I} - \mathbf{B} \mathbf{B}^T \\ \mathbf{u} (e + p + \frac{1}{2} B^2) - \mathbf{B} (\mathbf{u}^T \mathbf{B}) \\ \mathbf{u} \mathbf{B}^T - \mathbf{B} \mathbf{u}^T \end{pmatrix} = 0$$

**Conservative**

$$\begin{pmatrix} \rho \\ \rho u \\ \rho v \\ \rho w \\ e \\ B_x \\ B_y \\ B_z \end{pmatrix}_t + \operatorname{div} \begin{pmatrix} \rho \mathbf{u} \mathbf{u}^T + (p + \frac{1}{2} B^2) \mathbf{I} - \mathbf{B} \mathbf{B}^T \\ \mathbf{u} (e + p + \frac{1}{2} B^2) - \mathbf{B} (\mathbf{u}^T \mathbf{B}) \\ \mathbf{u} \mathbf{B}^T - \mathbf{B} \mathbf{u}^T \end{pmatrix} = -(\nabla \cdot \mathbf{B}) \begin{pmatrix} 0 \\ B_x \\ B_y \\ B_z \\ \mathbf{u}^T \mathbf{B} \\ u \\ v \\ w \end{pmatrix}$$

**Non-conservative  
(Symmetrizable -  
Godunov, Powell)**

$$\mathbf{u} = (u, v, w)^T$$

$$\mathbf{B} = (B_x, B_y, B_z)^T$$

$$B^2 = B_x^2 + B_y^2 + B_z^2$$

$$p = (\gamma - 1) \left( e - \frac{1}{2} \rho (u^2 + v^2 + w^2) - \frac{1}{2} (B_x^2 + B_y^2 + B_z^2) \right)$$

Conservative form is not symmetrizable

Two Entropy Split forms (ES & ES2) include the Godunov term are considered

# Entropy Split Methods (ES) for MHD

$$\mathbf{q}_t + \mathbf{f}_x^{(x)} + \mathbf{f}_y^{(y)} + \mathbf{f}_z^{(z)} + \mathbf{e} \operatorname{div} \mathbf{B} = 0.$$

The entropy splitting on; e.g., the  $x$ -flux derivative of the form

$$\mathbf{f}_x^{(x)} = \frac{\beta}{\beta + 1} \mathbf{f}_x^{(x)} + \frac{1}{\beta + 1} \mathbf{f}_{\mathbf{v}}^{(x)} \mathbf{v}_x,$$

and similarly for the other directions, to obtain

$$\mathbf{q}_t + \frac{\beta}{\beta + 1} (\mathbf{f}_x^{(x)} + \mathbf{f}_y^{(y)} + \mathbf{f}_z^{(z)} + \mathbf{e} \operatorname{div} \mathbf{B}) + \frac{1}{\beta + 1} (\mathbf{f}_{\mathbf{v}}^{(x)} \mathbf{v}_x + \mathbf{f}_{\mathbf{v}}^{(y)} \mathbf{v}_y + \mathbf{f}_{\mathbf{v}}^{(z)} \mathbf{v}_z + \mathbf{e} \operatorname{div} \mathbf{B}) = 0.$$

The Jacobians with respect to the entropy variables,  $\mathbf{f}_{\mathbf{v}}$ , are not symmetric, but thanks to the presence of the source term

$$\mathbf{f}_{\mathbf{v}}^{(x)} \mathbf{v}_x + \mathbf{e} \mathbf{B}_x = A^{(x)} \mathbf{v}_x,$$

where  $A^{(x)}$  is symmetric. The  $y$ - and  $z$ - directions are similar. Hence, the split form of the equations

$$\textbf{(ES)} \quad \mathbf{q}_t + \frac{\beta}{\beta + 1} (\mathbf{f}_x^{(x)} + \mathbf{f}_y^{(y)} + \mathbf{f}_z^{(z)}) + \frac{1}{\beta + 1} (A^{(x)} \mathbf{v}_x + A^{(y)} \mathbf{v}_y + A^{(z)} \mathbf{v}_z) + \frac{\beta}{\beta + 1} \mathbf{e} \operatorname{div} \mathbf{B} = 0,$$

## Entropy Split Methods (ES2) for MHD

A different form of the equations is obtained by replacing the weight  $\beta/(1 + \beta)$  on the Godunov term in (46) by 1,

$$\mathbf{q}_t + \frac{\beta}{\beta + 1} (\mathbf{f}_x^{(x)} + \mathbf{f}_y^{(y)} + \mathbf{f}_z^{(z)}) + \frac{1}{\beta + 1} (A^{(x)} \mathbf{v}_x + A^{(y)} \mathbf{v}_y + A^{(z)} \mathbf{v}_z) + \mathbf{e} \operatorname{div} \mathbf{B} = 0. \quad (47)$$

We refer to this alternative form as ES2.

It is noted that by looking at the non-split form of the equations it is seen that ES2 is equivalent with

$$\mathbf{q}_t + \mathbf{f}_x^{(x)} + \mathbf{f}_y^{(y)} + \mathbf{f}_z^{(z)} + \frac{2 + \beta}{1 + \beta} \mathbf{e} \operatorname{div} \mathbf{B} = 0.$$

The weight in front of the Godunov term varies between 2 and 1 when  $\beta$  varies between 0 and  $\infty$ .

# Semi Discrete Entropy Split Approximation

$$\frac{d}{dt}\mathbf{q}_j + \frac{\beta}{\beta+1}D\mathbf{f}_j + \frac{1}{\beta+1}A_jD\mathbf{v}_j + \frac{\beta}{\beta+1}\mathbf{e}_jDB_1 = 0, \quad j = 1, \dots, N, \quad (48)$$

where  $D$  is a linear finite difference operator; e.g.,  $\mathbf{f}_x(x_j) \approx Df_j = \frac{1}{\Delta x} \sum_{k=1}^p \alpha_k (\mathbf{f}_{j+k} - \mathbf{f}_{j-k})$  and  $\beta > 0$  is a parameter related to  $\alpha$  by  $\beta = (\alpha + \gamma)/(1 - \gamma)$ .

Although this entropy split method is formally defined in the same way as in the case of the gas dynamics equations, because the MHD flux function is not homogeneous, entropy conservation cannot be proved in the same way

# Non-uniqueness of Ducros et al. Splitting for MHD

*(Minimize the use of numerical dissipation for high order central schemes)*

- MHD inviscid (ideal) flux derivatives consist of **triple** products of conservative variables & their derivatives
- Splitting such triple products is **not unique**  
*(See Sjogreen & Yee, ICOSAHOM-2016 for the chosen forms)*
- Splitting all 8 flux derivatives, partial or just the gas dynamic portion
- Solving the Conservative MHD
- Solving the Godunov/Powell symmetrizable MHD (Non-conservative)

**The above consists of 8 combinations**

## 2D Alfven Wave Test Case

$$\begin{aligned}\rho &= 1 \quad p = 1 \quad \mathbf{v} = A(-\sin \alpha \sin \phi, \cos \alpha \sin \phi, \cos \phi) \\ \mathbf{B} &= (\cos \alpha - A \sin \alpha \sin \phi, \sin \alpha + A \cos \alpha \sin \phi, A \cos \phi) . \\ \phi &= 2\pi(x \cos \alpha + y \sin \alpha + t)\end{aligned}$$

where  $\alpha = 30^\circ$  and  $A = 0.1$  are chosen with periodic boundary conditions

=====

The computational domain is  $1.3546 \times 2.1915$

The flow is solved up to time 300 until all methods eventually diverge


$$\Delta x = \Delta y = 0.022$$

$$\Delta x = \Delta y = 0.011$$

=====




# Alfven Wave Beta Study (L2-Norm Error vs. Time)

(Comparison between *ES* & *ES2* by 8<sup>th</sup>-order Entropy Split Method)




# Alfven Wave (Max-Norm Error vs. Time)

(Beta Parameter Study: 8<sup>th</sup>-order Entropy Split Method)



# Alfven Wave (Max-Norm Error vs. Time)

## (Comparison Among Seven 8<sup>th</sup>-order Methods)



## Brio-Wu MHD Shock Tube Problem


$$(\rho_L, u_L, p_L, B^{(x)}, B^{(y)}, B^{(z)}) = (1, 0, 1, 0.75, 1, 0)$$

$$(\rho_R, u_R, p_R, B^{(x)}, B^{(y)}, B^{(z)}) = (0.125, 0, 0.1, 0.75, -1, 0).$$


# Brio-Wu MHD Shock Tube Problem

*(Comparison Among Seven Methods + 7th-order nonlinear filter)*


Density





Density (Zoom)



Pressure




Pressure (Zoom)





# Brio-Wu MHD Shock Tube Problem

*(Comparison Among Seven Methods + 7th-order nonlinear filter)*

$B_y$  magnetic field



$B_y$  magnetic field (Zoom)



# Summary

## Entropy Split (ES) Schemes & Entropy Conservative (EC) Methods

### GAS dynamics: Classical Central, DRP Central & Pade (Compact) Spatial Schemes

- **ES** can improve nonlinear stability for **smooth flows** in general without added numerical dissipation
- Nonlinear filter version of **ES** schemes can improve stability & accuracy for DNS & LES with shocks
- High order entropy conserving methods (centered or nonlinear filter version) **ES & EC** provide similar stability & accuracy improvement.
- **Tadmor-type EC methods are approximately twice the CPU than ES**

### Plasma:

- **ES** can improve nonlinear stability in general for **smooth flows**  
**but MHD equations dependent**
- Nonlinear filter version of split schemes can improve stability & accuracy for flows with discontinuities **but MHD equations dependent**
- High order entropy conserving methods (centered or nonlinear filter version) can provide **different stability & accuracy improvement**, depending on the **forms of the MHD equations & the choice of entropy fluxes**

---

## References

1. Adam Y.: Highly accurate compact implicit methods and boundary conditions. *J Comput Phys.* **24** 10-22 (1977).
2. Arakawa, A.: Computational design for long-term numerical integration of the equations of fluid motion: two-dimensional incompressible flow, Part I. *J. Comput. Phys.* **1**, 119–143 (1966).
3. Balsara, D., Shu, Chi-Wang: Monotonicity Preserving Weighted Essentially Non-oscillatory Schemes with Increasingly High Order of Accuracy. *J. Comput. Phys.* **160** 405-452 (2000).
4. Blaisdell, G.A., Spyropoulos, E.T., Qin, J.H.: The Effect of the Formulation of Nonlinear Terms on Aliasing Errors in Spectral Methods. *Appl. Num. Math.* **21**, 207-219 (1996).
5. Bogey, C, Bailly, C.: A Family of Low dispersive and Low Dissipative Explicit Schemes for Computing the Aerodynamic Noise. AIAA-paper 2002-2509, 8th ARAA/CEAS Aeroacoustics Conference & Exhibit, June 17-19, 2002, Breckenridge Colorado.
6. Brambley, E.J.: Optimized finite-difference (DRP) schemes perform poorly for decaying or growing oscillations. *J. Comput. Phys.*, **324**, 258-274 (2015).
7. Brio, M. and Wu, C.C.: An Upwind Differencing Scheme for the Equations of Ideal Magneto-hydrodynamics, *Journal of Computational Physics*, **75**, 400-422 (1988).
8. Cano, B., Sanz-Serna, J.M.: Error growth in the numerical integration of periodic orbits, with application to Hamiltonian and reversible systems. *SIAM J. Numer. Anal.* **34** 1391417 (1997).
9. Chang, C.-C., Chen, M.-H., and Teng, C.-T.: High-Order Summation-by-Parts Implicit Difference Operators for Wave Problems: Presentation, url [https://www.math.ncku.edu.tw/~mhchen/HOFD/HOSBP\\_slide.pdf](https://www.math.ncku.edu.tw/~mhchen/HOFD/HOSBP_slide.pdf)
10. Carpenter, M.H., Gottlieb, D., Arbabanel, S., and Don, W.-S.: The Theoretical Accuracy of Runge-Kutta Time Discretizations for the Initial Boundary Value Problem: A Study of the Boundary Error: *SIAM J. Sci. Comput.* **16**, 1241-1252, (1995).
11. Castro, M.J., Gallardo, J. M., and Marquina, A.: Jacobian-Free Incomplete Riemann Solvers, *Proceedings of Theory, Numerics and Application ProblemsI: Aachen Germany*, 292-307 (2016).
12. Colella, P. and Woodward, P.R.: The Piecewise Parabolic Method (PPM) for Gas-Dynamical Simulations. *J. Comput. Phys.*, **54**, 174-201 (1984).
13. Coppola, G., Capuano, F., Pirozzoli, S., de Luca, L.: Numerically Stable Formulations of Convective Terms for Turbulent Compressible Flows. *J. Comput. Phys.*, 2019, DOI:10.1016/j.jcp.2019.01.007.
14. Ducros, F., Laporte, F., Soulères, T., Guinot, V., Moinat, P., Caruelle, B.: High-Order Fluxes for Conservative Skew-Symmetric-Like Schemes in Structured Meshes: Application to Compressible Flows, *J. Comput. Phys.* **161**, 114-139 (2000).
15. Fang, J., Gao, F., Moulinec, C., and Emerson, D.R.: An Improved Parallel Compact Scheme for Domain-Decoupled Simulation of Turbulence. *IJNUMF*, DOI: 10.1002/fld.4731 (2019).
16. Flash Manual - UserManual.wiki, University of Chicago, 2021.
17. Gassner, G., Winters, A.R.: A Novel Robust Strategy for Discontinuous Galerkin Methods in Computational Fluid Mechanics: Why? When? What? Where?, *Frontiers in Physics* **8**:500690, DOI:10.3389/fphy.2020.500690, Jan. 2021.
18. Gerritsen, M, Olsson, P.: Designing an Efficient Solution Strategy for Fluid Flows. I. A Stable High Order Finite Difference Scheme and Sharp Shock Resolution for the Euler Equations. *J. Comput. Phys.* **129**, 245-262 (1996).
19. Godunov, S.K.: Symmetric Form of the Equations of Magnetohydrodynamics. *Numerical Methods for Mechanics of Continuum Medium*, **13**, no. 1, 26-34 (1972).
20. Gottlieb, D., Gustafsson, B., Olsson, P., and Strand B.: On the Superconvergence of Galerkin Methods for Hyperbolic IBVP. *SIAM J. Numer. Anal.* **33**, 1778–1796, (1996).
21. Gurski, K.F.: An HLLC-Type Approximate Riemann Solver for Ideal Magnetohydrodynamics. *SIAM J. Sci. Comput.* **25**, 2165-2187 (2004).
22. Gurski, K.F.: An HLLC-Type Approximate Riemann Solver for Ideal Magnetohydrodynamics. *SIAM J. Sci. Comput.* **25**(6), 2165187 (2004).
23. Harten, A: On the Symmetric Form of Systems for Conservation Laws with Entropy: *J. Comput Phys.* **49**, 151 (1983).
24. Hirsh, R.S.: Higher Order Accurate Difference Solutions of Fluid Mechanics Problems by a Compact Differencing Technique. *J Comput Phys.* **19**, 90-109 (1975).
25. Johansson, S.: High Order Summation by Parts Operator Based on a DRP Scheme Applied to 2D, Technical Report 2004-050, Uppsala University, Sweden.

26. Kennedy, C.A. and Gruber, A.: Reduced Aliasing Formulations of the Convective Terms Within the Navier-Stokes Equations. *J. Comput. Phys.*, **227** 1676-1700 (2008).
27. Li, S.: An HLLC Riemann Solver for Magneto-Hydrodynamics. *J. Comput. Phys.* **203** (1) 344-357 (2005).
28. Olsson, P., Oliger, J.: Energy and maximum norm estimates for nonlinear conservation laws. *RIACS Technical Report 94.01*, (1994).
29. Palha, A., Gerritsma, M.: A Mass Energy, Enstrophy and Vorticity Conserving (MEEVC) Mimetic Spectral Element Discretization for the 2D Incompressible Navier-Stokes Equations. DOI:10.1016/j.jcp.2016.10.009.
30. Parsani, M., Carpenter, M.H., Nielsen, E.J.: Entropy stable wall boundary conditions for the three-dimensional compressible Navier-Stokes equations. *J. Comput. Phys.* **292**, 88-113 (2015).
31. Pirozzoli, S.: Generalized conservative approximations of split convective derivative operators. *J. Comput. Phys.*, **219**, 7180-90 (2010).
32. Ranocha, H.: Entropy Conserving and Kinetic Energy Preserving Numerical Methods for the Euler Equations Using Summation-by-Parts Operators. *Proceedings of the ICOSAHOM-2018*, Imperial College, London, UK, July 9-13, 2018.
33. Sandham, N.D., Li, Q., Yee, H.C.: Entropy Splitting for High-Order Numerical Simulation of Compressible Turbulence. *J. Comput. Phys.* **23**, 307-322 (2002).
34. Sengupta, T.K., Dipankar, A., and Rao, K.: "A New Compact Scheme for Parallel Computing using Domain Decomposition. *J. Comput. Phys.*, **220** 654-677 (2007).
35. Shu, C.-W., Osher, S.J.: Efficient Implementation of Essentially Non-oscillatory Shock Capturing Schemes. *J. Comput. Phys.*, **83**, 32-78 (1989).
36. Sjögreen, B., Yee, H.C.: Multiresolution Wavelet Based Adaptive Numerical Dissipation Control for High Order Methods. *J. Scientific Computing*, **20**, 211-255 (2004).
37. Sjögreen, B., Yee, H.C. and Vinokur, M.: On high order finite-difference metric discretizations satisfying GCL on moving and deforming grids. *J. Comput. Phys.*, **265**, 211-220 (2014)
38. Sjögreen, B., Yee, H.C.: Accuracy consideration by DRP schemes for DNS and LES of compressible flow Computations. *Computers & fluids*, **159**, 123-136 (2017).
39. Sjögreen, B., Yee, H.C.: An entropy stable method revisited: Central differencing via entropy splitting and SBP. *Entropy Splitting for Central Schemes with SBP Energy Estimates Skew-Symmetric Splitting Revist*. *Proceeding of ICOSAHOM-2018*, July 9-13, 2018, Imperial College, London, UK.
40. Sjögreen, B., Yee, H.C.: High order entropy conservative central schemes for wide ranges of compressible gas dynamics and MHD flows. *J. Comput. Phys.*, **364**, 153-185 (2018).
41. Sjögreen, B., Yee, H.C.: Skew-Symmetric Splitting for Multiscale Gas Dynamics and MHD Turbulence Flows. *Extended version of Proceedings of ASTRONUM-2016*, June 6-10, 2016, Monterey, CA, USA, *J. Scientific Computing*, **83** 43 (2020).
42. Sjögreen, B., Yee, H.C.: Accuracy Consideration by DRP Schemes for DNS and LES of Compressible Flow Computations. *Special issue in Computers & Fluids in honor of Prof. Toro's 70th birthday*, **159** 123-136 (2017).
43. Sjögreen, B., Yee, H.C.: . High order entropy conserving central schemes for wide ranges of compressible gas dynamics and MHD flows. *J. Comput. Phys.*, **364**, 153-185 (2018).
44. Sjögreen, B., Yee, H.C.: Entropy Stable Method for the Euler Equations Revisited: Central Differencing via Entropy Splitting and SBP. *Proceeding of ICOSAHOM-2018*, July 9-13, 2018, Imperial College, London, UK; *J. Scientific Computing*, doi.org/10.1007/s10915-019-01013-1, **81** 1359-1385 (2019) .
45. Sjögreen, B., Yee, H.C.: Entropy Stable Method for the Euler Equations Revisited: Central Differencing via Entropy Splitting and SBP. *J. Scientific Computing*, **81**:1359-1385 (2019), <https://doi.org/10.1007/s10915-019-01013-1>.
46. Sjögreen, B., Yee, H.C.: Construction of Conservative Numerical Fluxes for the Entropy Split Method. *Comm. Applied Mathematics and Computation (CAMC)*, 2021.
47. Sjögreen, B., Yee, H.C., Kotov, D.V., and A. Kritsuk, A. G.: Skew-Symmetric Splitting for Multiscale Gas Dynamics and MHD Turbulence Flows. *J. Scientific Computing*, **83**:1-43, (2020)
48. Sjögreen, B., Yee, H.C.: High Order Compact Central Spatial Discretization Under the Framework of Entropy Split Methods. *Proceedings of the ICOSAHOM21*, July 12-16, 2021.
49. Strand, B.: Summation by Parts for Finite Difference Approximations for  $d/dx$ . *J. Comput. Phys.* **110**, 47-97, (1996).
50. Tadmor, E.: Entropy Stability Theory for Difference Approximations of Nonlinear Conservation Laws and Related Time-Dependent Problems. *Acta Numerica* **12**, 451-512 (2003).

51. Taylor, G., Green, A.: Mechanism of the Production of Small Eddies from Large Ones. *Proc. R. Soc. Lond. A* **158**, 499–521 (1937).
52. Tam, C.K.W.: A CAA Primer for Practicing Engineers. AEDC-TR-08-2, April 2008, Arnold Engineering Development Center, Arnold Air Force Base, Tennessee.
53. Tam, C.K.W.: Computational Aeroacoustics: A Wave Number Approach. Cambridge Aerospace series, Vol. 33, Cambridge University Press, New York, 2012.
54. Vinokur, M., Yee, H.C.: Extension of Efficient Low Dissipation High-Order Schemes for 3D Curvilinear Moving Grids. *Front. Comput. Fluid Dyn.* 12964 (2002); Also, Proceedings of the Robert MacCormack 60th Birthday Conference, June 26-28, 2000, Half Moon Bay, CA, NASA/TM-2000-209598.
55. Winters, A.R., Gassner, G.J.: Affordable, entropy conserving and entropy stable flux functions for the ideal MHD equations. *J. Comput. Phys.*, **304** 72-108, (2016).
56. Yee, H. C., Sandham, N.D., Djomehri, M. J.: Low-Dissipative High Order Shock-Capturing Methods Using Characteristic-Based Filters. *J. Comput. Phys.*, **150**, 199-238 (1999).
57. Yee, H. C., Vinokur, M., Djomehri, M. J.: Entropy Splitting and Numerical Dissipation. *J. Comp. Phys.* **162**, 33-81 (2000).
58. Yee, H. C., Sjögren, B.: Efficient Low Dissipative High Order Schemes for Multiscale MHD Flows, II: Minimization of  $\text{div } \mathbf{B}$  Numerical Error. *J. Sci. Comput.*, Vol. 29, No. 1, October 2006 ( 2005), DOI: 10.1007/s10915-005-9004-5
59. Yee, H. C., Sjögren, B.: Development of Low Dissipative High Order Filter Schemes for Multiscale Navier-Stokes and MHD Systems. *J. Comput. Phys.*, **225** 910-934 (2007).
60. Yee, H. C., Sjögren, B.: Simulation of Richtmyer-Meshkov Instability by Sixth-order Filter Methods. *Shock Waves J.* 17, 185-193 (2007), DOI 10.1007/s00193-007-0104-z.
61. Yee, H. C., Sjögren, B.: Adaptive Filtering and Limiting in Compact High Order Methods for Multiscale Gas Dynamics and MHD Systems. *Computers & Fluid*, **37** 593-619 (2008).
62. Yee, H. C., Sjögren, B.: High Order Filter Methods for Wide Range of Compressible Flow Speeds. Proceedings of the ICOSAHOM09, June 22-26, 2009, Trondheim, Norway.
63. Yee, H. C., Sjögren, B.: On Entropy Conservation and Kinetic Energy Preservation Methods. Proceedings of the ICOSAHOM-2019, July 1-5, 2019, Paris, France.
64. Yee, H. C., Sjögren, B.: Comparative Study on a Variety of Structure-Preserving High Order Spatial Discretizations with the Entropy Split Methods for MHD. Proceedings of the ICOSA-HOM21, July 12-16, 2021.

# Skew-Symmetric Splitting the Derivative of Products

*(Minimize the use of numerical dissipation for high order central schemes)*

**Splitting of the derivative of a product in conservative & non-conservative part:**

$$(ab)_x = \frac{1}{2}(ab)_x + \frac{1}{2}ab_x + \frac{1}{2}a_xb \quad \text{Entropy splitting of Euler flux, Yee et al}$$

**Split approximation can be written into conservative form:**

$$\frac{1}{2}D_0(ab)_j + \frac{1}{2}a_jD_0b_j + \frac{1}{2}b_jD_0a_j = \frac{1}{4}D_+(a_j + a_{j-1})(b_j + b_{j-1})$$

$D_0$ : 2<sup>nd</sup>-order central,  $D_+u_j = (u_{j+1} - u_j)/\Delta x$

**Can be generalized to 2p<sup>th</sup> order accurate: Ducros et al. 2000**

$$D_{0p}u_j = \sum_{k=1}^p \alpha_k^{(p)} D_0(k)u_j \quad D_0(k)u_j = (u_{j+k} - u_{j-k})/(2k\Delta x)$$
$$\sum_{k=1}^p \alpha_k^{(p)} = 1 \quad \sum_{k=1}^p \alpha_k^{(p)} k^{2n} = 0, \quad n = 1, \dots, p-1$$