Understanding Atmospheric Absorption Effects on UV Spectra from Sounding Rockets using a Spherical-Shells Model


Summary Abstract

Our team is working on building and calibrating the FURST sounding rocket, with an expected launch in mid-2023. The goal is to image the most complete and highest resolution UV spectra to date. To do this, precise radiometric and wavelength calibration techniques have been developed. We describe below our model of O2 atmospheric absorption and couple that with simulated FURST images. With a high-enough SNR, we can estimate our ability to use absorption peaks for calibration, or for back-calculating atmospheric properties. If data is available, this method could be applied to older sounding rocket data to find hidden science.

Motivation

- **For FURST:**
  - To obtain high-resolution UV spectra [3].
  - To close the large data gap [17, 18].
  - To directly compare with HST & JWST [11].
  - To show the value of FURST as a satellite.
- **For this Research:**
  - To investigate potential SNR issues [23].
  - To aid in spectral calibration [7, 5, 6].
  - To better understand this effect [10, 22, 21].
  - To discover science hidden inside data.

Instrument




Spectral Range: 120-181nm
Resolution: $\mathcal{R}$>2·10^4 (ΔV<15km/s)
Launch: 2023 from WSMR, NM
Altitudes: 110-260km

Methods

- Transmission T is calculated from optical depth τ via Beer-Lambert's Law and the Differential Path Length ΔL:


$$T = e^{-\tau},\ \tau_{i,j} = \sum_{i'=i}^{\infty} \sigma_{i',j}\eta_{i'}\Delta L_{i'}$$



- Simulate results to match [15].
- Geometric and Trigonometric relations used to find ΔL.
- Earth's radius and Snell's Law allow for changing zenith angle [2, 8, 14, 19].
- Density data from NRL-MSISE [16].
- Absorption cross-section data from MPI-Mainz UVVIS [9, 13, 20, 12].
- HRTS data is folded into FURST optics simulate a UV spectra [1].
- Altitude / Exposure time effect SNR.

Results

- The spherical-shells model improves our estimated SNR.
- In the range 120-130.4nm, we show absorption lines for calibration.
  - Using 2 data point pairs (pixel and wavelength values) we calculate the spectral plate scale to within 1%.
- Running a forward model of density, we can to use the ratio between signals for density recalculation.
  - Using 2 images we fit a density model to within 16% RMSE [15].
- Further improvements include:
  - Seeing how varying exposure times effect SNR.
  - Using more exposures/wavelength ranges together.
  - Calculating the O2 resonant absorption contribution [4].
  - Applying these methods to existing actual raw data to verify.



Acknowledgements

The primary author wishes to thank their advisors, mentors, and colleagues for advice and support along the way, all of the staff at UAH for the funding support, and his wife Crystal. *This material is based upon work supported by the NSF EPSCoR RII-Track-1.2a (Non-invasive plasma diagnostics for LTP) Cooperative Agreement OIA-1655280. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.*

References **[1]** P Brekke. In: *The Astrophysical Journal Supplement Series* 87 (1993). **[2]** C. Cornwall, A. Horiuchi, and C. Lehman. 2022. **[3]** W Curdt et al. In: *Astronomy & Astrophysics* 375.2 (2001). **[4]** D. A. Degenstein. The University of Saskatchewan, 2000. **[5]** N. Donders et al. In: *AGU Fall Meeting Abstracts*. Vol. 2020. 2020. **[6]** N. Donders et al. In: *Proceedings of the International Astronautical Congress*. 2020. **[7]** N. Donders et al. In: *AGU Fall Meeting Abstracts*. Vol. 2019. 2019. **[8]** N. ESRL. 2005. **[9]** I. E. Gordon et al. In: *Journal of Quantitative Spectroscopy and Radiative Transfer* 203 (2017). **[10]** W. He et al. In: *Remote Sensing* 11.22 (2019). **[11]** P. G. Judge et al. In: *The Astrophysical Journal* 848.1 (2017). **[12]** H. Keller-Rudek et al. 2014. **[13]** H. Keller-Rudek et al. In: *Earth System Science Data* 5.2 (2013). **[14]** Y. Liu and P. H. Daum. In: *Journal of Aerosol Science* 39.11 (2008). **[15]** R. Meier. In: *Space Science Reviews* 58.1 (1991). **[16]** 2022. **[17]** H Peter. In: *The Astrophysical Journal* 516.1 (1999). **[18]** H. Peter and P. Judge. In: *The Astrophysical Journal* 522.2 (1999). **[19]** P Phillips. In: *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 97.684 (1920). **[20]** R. Sander et al. In: *EGU General Assembly Conference Abstracts*. 2014. **[21]** K. Sun et al. In: *Atmospheric Measurement Techniques Discussions* (2022). **[22]** K. Sun et al. In: *Geophysical Research Letters* 45.11 (2018). **[23]** G. D. Vigil et al. In: *Journal of Astronomical Telescopes, Instruments, and Systems* 7.3 (2021).

Nicolas Donders[1], Amy Winebarger[2], Charles Kankelborg[3], Larry Paxton[4], Genevieve Vigil[2], Gary Zank[1]

[1]Department of Space Science and The Center for Space Plasma and Aeronomic Research, The University of Alabama in Huntsville,
[2]NASA Marshall Space Flight Center, [3]Montana State University, [4]Johns Hopkins Applied Physics Laboratory