



Algorithm Stability and the Long-Term Geospace Data Record from TIMED/SABER



M. G. Mlynczak¹

¹NASA Langley Research Center



EGU Meeting 2024 – Vienna Austria


Acknowledgement: NASA Heliophysics Division TIMED Project

Concept of Geospace Data Record

- A Geospace Data Record (GDR) is analogous to the well-known Climate Data Record (CDR) of tropospheric climate science
- A CDR defined as a “*time series of sufficient length, consistency, and continuity to determine climate variability and change*”
- Geospace* is defined as the region of the atmosphere from the mesosphere to the exobase (60 km to 600 km)
- A Geospace Data Record (GDR) is a time series of sufficient length, consistency, and continuity to determine *geospace variability and change*
- Geospace is changing due to increasing CO₂
- Solar variability complicates interpretation of long-term change
- Knowledge of Geospace change is critical for future space law, space policy, and the space insurance industry (Mlynczak et al., 2001)

The 21-Year Geospace Data Record from SABER on TIMED

Daily Global Average Temperature, Geopotential Height, and Thickness




Temperature, GPH, and Thickness Trends

Change attributed to increasing CO₂ and declining solar activity

Table 1 Observed Changes From 2002 to 2019 in Global Annual Mean Temperature, Geopotential Height (GPH), and Thickness (Relative to 1 hPa) (Obs. Change); the Trend in Each Quantity Due to Increasing CO ₂ (Derived Trend); the Estimated Magnitude of the Change in Each Quantity Due to Increasing CO ₂ (Δ CO ₂); and the Estimated Magnitude of the Change in Each Quantity Due to Declining Solar Activity (Δ Solar)					
Quantity	P (hPa)	Obs. Change	Derived trend	Δ CO ₂	Δ Solar
Temperature	1	-1.77 K	-0.55 ± 0.07 K/dec	-0.99 K	-0.78 K
	10 ⁻¹	-2.05 K	-0.63 ± 0.11 K/dec	-1.13 K	-0.92 K
	10 ⁻²	-3.01 K	-0.53 ± 0.17 K/dec	-0.95 K	-2.06 K
	10 ⁻³	-5.12 K	-0.51 ± 0.17 K/dec	-0.92 K	-4.20 K
	10 ⁻⁴	-19.05 K	-2.60 ± 0.67 K/dec	-4.68 K	-14.37 K
GPH	1	-121 m	-8 ± 12.3 m/dec	-14.4 m	-106.6 m
	10 ⁻¹	-240 m	-47 ± 12.9 m/dec	-84.6 m	-155.4 m
	10 ⁻²	-397 m	-84 ± 14.9 m/dec	-151.2 m	-245.8 m
	10 ⁻³	-761 m	-127 ± 20.0 m/dec	-228.6 m	-532.4 m
	10 ⁻⁴	-1,454 m	-197 ± 33.6 m/dec	-354.6 m	-1,099.4 m
Thickness	1	-118 m	-40 ± 3.7 m/dec	-72 m	-46 m
	10 ⁻¹	-276 m	-77 ± 13.4 m/dec	-138.6 m	-137.4 m
	10 ⁻²	-640 m	-119 ± 18.5 m/dec	-214.2 m	-425.8 m
	10 ⁻³	-1,333 m	-190 ± 31.6 m/dec	-342 m	-991 m
	10 ⁻⁴	-1,333 m	-190 ± 31.6 m/dec	-342 m	-991 m

Note. One-sigma trend uncertainties are also listed in the derived trend column.

Global Annual Means



Geopotential Height

Temperature

Thickness

- Mesosphere and Lower Thermosphere (MLT) cooled substantially from 2002 to 2022
- MLT also contracted by over 1400 meters
- Cause is determined to be increasing CO₂ and declining solar activity over the time period

Challenges to creating GDRs from satellite observations

Generating a consistent dataset with changes that are solely due to the variability and change in the atmosphere

Several factors can contribute to ‘instability’ of a data record that are instrument or algorithm induced:

- Changing satellite orbit
- Changes to algorithm parameters/inputs over time
- Changing calibration (despite on-board verification)

SABER has been shown to have a remarkably stable calibration, with changes in temperature of ~ 0.1 K/decade

Example of Algorithm Instability in SABER

SABER Temp (2020 – 2009)

Large, unanticipated difference above 85 km




SABER Temp retrieval uses CO₂ concentrations from WACCM model

WACCM version changed inadvertently in Dec. 2019

Larger WACCM CO₂ (red curve) led to lower Temps

SABER reverted to original WACCM version


Temps now consistent with CO₂ increase and natural variability



GDRs and The Challenge of Achieving Continuity

- GDRs must be of sufficient length to ‘determine geospace variability and change’
- The fleet of satellites launched since early 2000’s is now aging and many satellites are decommissioned
- Recent loss of ICON and AIM magnify the challenge
- SABER continues to operate nominally
- However, a gap seems likely this decade
- Community advocacy for continuity is critical !!!

Mind The Gap!



Mind The Gap!

SABER Temp. stability determined by comparison with COSMIC GPS Temps

GPS Temps are inherently stable due to their traceability to the SI definition of the second

