

Gas Sources from the Coma and Nucleus of Comet 46P/Wirtanen Observed Using ALMA

M. A. CORDINER,^{1,2} N. X. ROTH,^{1,2} S. N. MILAM,¹ G. L. VILLANUEVA,³ D. BOCKELÉE-MORVAN,⁴ A. J. REMIJAN,⁵
S. B. CHARNLEY,¹ N. BIVER,⁴ D. C. LIS,⁶ C. QI,⁷ B. P. BONEV,⁸ J. CROVISIER,⁴ AND J. BOISSIER⁹

¹*Astrochemistry Laboratory, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771, USA.*

²*Department of Physics, Catholic University of America, Washington, DC 20064, USA.*

³*Planetary Systems Laboratory, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771, USA.*

⁴*LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, 5 place Jules Janssen, F-92195 Meudon, France.*

⁵*National Radio Astronomy Observatory, Charlottesville, VA 22903, USA.*

⁶*Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Drove Drive, Pasadena, CA 91109, USA.*

⁷*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MS 42, Cambridge, MA 02138, USA.*

⁸*Department of Physics, American University, Washington D.C., USA.*

⁹*Institut de Radioastronomie Millimétrique, 300 rue de la Piscine, F-38406, Saint Martin d’Heres, France.*

ABSTRACT

Gas-phase molecules in cometary atmospheres (comae) originate primarily from (1) outgassing by the nucleus, (2) sublimation of icy grains in the near-nucleus coma, and (3) coma (photo-)chemical processes. However, the majority of cometary gases observed at radio wavelengths have yet to be mapped, so their production/release mechanisms remain uncertain. Here we present observations of six molecular species towards comet 46P/Wirtanen, obtained using the Atacama Large Millimeter/submillimeter Array (ALMA) during the comet’s unusually close (~ 0.1 au) approach to Earth in December 2018. Interferometric maps of HCN, CH₃OH, CH₃CN, H₂CO, CS and HNC were obtained at an unprecedented sky-projected spatial resolution of up to 25 km, enabling the nucleus and coma sources of these molecules to be accurately quantified. The HCN, CH₃OH and CH₃CN spatial distributions are consistent with production by direct outgassing from (or very near to) the nucleus, with a significant proportion of the observed CH₃OH originating from sublimation of icy grains in the near-nucleus coma (at a scale-length $L_p = 36 \pm 7$ km). On the other hand, H₂CO, CS and HNC originate primarily from distributed coma sources (with L_p values in the range 550–16,000 km), the identities of which remain to be established. The HCN, CH₃OH and HNC abundances in 46P are consistent with the average values previously observed in comets, whereas the H₂CO, CH₃CN and CS abundances are relatively low.

Keywords: Comets, individual: 46P/Wirtanen — Radio interferometry — Molecular lines — Astrochemistry

1. INTRODUCTION

Measurements of cometary compositions provide a unique tool for investigating ice chemistry in the protosolar disk midplane during the epoch of planet formation, and can therefore provide insight into the reagents available for prebiotic chemistry in the early Solar System (Mumma & Charnley 2011). A wealth of molecular species were recently detected in comet 67P by the Rosetta spacecraft (Altwegg et al. 2019), yet remote observations of coma gases remain the most common method for determining cometary compositions (Cochran et al. 2015).

The coma is typically understood in terms of a quasi-spherical expanding outflow of ‘parent’ species, sublimating directly from the nucleus, with ‘daughter’ species originating from photolysis of the parents in the coma (Haser

1957). Several coma molecules, on the other hand, exhibit ‘distributed’ (spatially extended) sources, some of which are believed to arise from the breakdown of macromolecular or dust-grain precursors (Meier et al. 1993; Cottin & Fray 2008; Cordiner et al. 2014, 2017a), the precise identity of which remains unknown, but could be related to the organic-rich, refractory material identified in comet 67P by the Rosetta mission (Capaccioni et al. 2015; Fray et al. 2016; Bardyn et al. 2017). An analysis of a sample of 26 comets led Mumma et al. (2017) to propose that thermal dissociation of ammonium salts (NH_4^+X^- , where X^- is a deprotonated acid) could be another source of gas-phase coma molecules, and this was found to be a plausible explanation for the abundances of several organics observed in the coma of 67P (Altwegg et al. 2020; Poch et al. 2020).

To-date, interferometric observations using the Atacama Large Millimeter/submillimeter Array (ALMA) have confirmed the presence of daughter (or distributed) sources of H_2CO in three comets, while the HNC and CS molecules have been found to exhibit distributed sources in two comets (Cordiner et al. 2014; Roth et al. 2021a; Biver et al. 2022). Earlier single-dish mapping work (Biver et al. 1999) and interferometric observations (Milam et al. 2006) identified extended H_2CO spatial distributions in comets C/1996 B2 (Hyakutake) and C/1995 O1 (Hale-Bopp), respectively. Infrared spectroscopic studies have also indicated the presence of both nucleus and coma sources for H_2CO and CO (Disanti et al. 1999; DiSanti et al. 2006), although no compelling evidence for a distributed CO source was found by Bockelée-Morvan et al. (2010). Despite decades of investigations into the chemical compositions of cometary comae and nuclei, it is surprising that the chemical origins of these commonly-detected (and relatively simple) coma gases still remain to be conclusively determined.

A common way to parameterize molecular production in comets as a function of distance from the nucleus is using the Haser daughter formula (Haser 1957). Through radiative transfer modeling of coma mapping observations, it is then possible to derive the characteristic distance scale at which a given species is produced (commonly expressed as a production scale length, or parent scale length, L_p ; Biver et al. 1999; Cordiner et al. 2014; Roth et al. 2021b). Due to the paucity of detailed studies to-date, and relatively large uncertainties on the derived L_p values for HNC, H_2CO and CS, combined with strong variability in the H_2CO parent scale lengths derived for different comets (even after correcting for heliocentric distance), further studies are needed to help improve our understanding of the distributions of these molecules in cometary comae. By comparison with laboratory measurements and detailed numerical models (e.g. Meier et al. 1993; Fray et al. 2006; Cordiner & Charnley 2021), observational characterization of molecular production as a function of distance from the nucleus allows proposed identities of the parent species to be tested and validated. Measurements of the parent scale lengths of coma daughter species are therefore important in our quest to better understand the native chemical constituents of comets, from which we gain new insights into the chemical processes that occurred during the earliest history of the Solar System.

Previous interferometric studies of molecular production scales in comets have been restricted to bright, long-period comets from the Oort cloud. But these objects represent only part of the wider population of small icy bodies available for study in our Solar System today. Gravitational scattering by the giant planets is believed to have redistributed comets **within** the inner Solar System and into their various present-day reservoirs: in particular, the Oort cloud and the Kuiper Belt or scattered disk. Most comets discovered each year come from the Oort cloud, while the scattered Kuiper disk is considered to be the source of Jupiter-family comets (JFCs). However, dynamical models have various predictions regarding the formative regions of comets, spanning a diverse range of heliocentric distances (e.g. Gomes et al. 2005; Tsiganis et al. 2005). It is therefore vitally important to study both Oort cloud comets (OCCs) and JFCs because it is unclear whether comets from each reservoir were formed in entirely overlapping regions in the protosolar disk. Comparative studies of JFCs and OCCs can also provide insights into the impact of Solar irradiation and thermal processing on the nucleus composition, since JFCs are typically subject to repeated cycles of irradiation, heating and cooling during their relatively frequent, periodic perihelion passages (Meech & Svoren 2004; Gkotsinas et al. 2022). Due to its extremely close approach to Earth ($\Delta = 0.077$ au on UT 2018-12-16), the 2018 apparition of comet 46P/Wirtanen provided a unique opportunity to observe the molecular coma of a Jupiter-family comet, at unprecedented spatial resolution from the ground.

In this study, we present ALMA observations of molecular emission from comet 46P/Wirtanen, conducted in early December 2018 just before its closest approach to Earth. The resulting spectral images include molecules previously identified as daughter/distributed species in Oort cloud comets (H_2CO , HNC and CS), as well as suspected parent species (HCN and CH_3OH). We also present the first map of CH_3CN , which is yet to be imaged in any comet, so its association with sublimating ices in the nucleus, or chemistry in the coma, remains unexplored. Through application of our recently-developed, 3D radiative transfer and excitation model (SUBLIME; Cordiner et al. 2022), the coma

Table 1. Observational parameters

Set.	UT Date	UT Time	Int. ^a (min)	r_H^b (au)	Δ^c (au)	$\dot{\Delta}^d$ (km s ⁻¹)	ν_R^e (GHz)	Band ^f	Baselines ^g (m)	θ_{min}^h ($''$)	θ_{PB}^i ($''$)	PWV ^j (mm)	ϕ_{STO}^k ($^\circ$)	PS Ang. ^l ($^\circ$)
1	2018-12-02	04:16–05:12	43	1.07	0.115	-7.2	354.5	7	15.1–952	0.45×0.31	16.4	0.63	44.1	36.3
2	2018-12-02	02:30–03:52	63	1.07	0.115	-7.4	351.8	7	15.1–952	0.41×0.31	16.5	0.90	44.2	36.2
3	2018-12-07	02:34–04:06	63	1.06	0.096	-5.9	241.8	6	15.1–784	0.62×0.55	24.0	0.78	42.2	38.5

^a On-source observing time.

^b Heliocentric distance of the comet.

^c Geocentric distance of the comet.

^d Comet’s mean topocentric radial velocity.

^e Representative frequency.

^f **ALMA receiver band 6 range is 211–275 GHz; band 7 covers 275–370 GHz.**

^g Range of antenna baseline lengths.

^h Angular resolution (dimensions of Gaussian fit to PSF) at ν_R .

ⁱ Primary beam FWHM at ν_R .

^j Median precipitable water vapor column length at zenith.

^k Sun-Target-Observer (phase) angle.

^l Position angle (in the plane of the sky) of the extended Sun-target vector, counter-clockwise from north.

93 temperature distribution is derived, parent scale-lengths and abundances are calculated, leading to new insights into
94 the nucleus *vs.* coma contributions of the observed molecules.

95 2. OBSERVATIONS

96 ALMA observations of 46P/Wirtanen were carried out on UT 2018-12-02 and 2018-12-07, when the comet was
97 ≈ 0.1 au from Earth (1.1 au from the Sun), using forty-three 12 m antennas in an intermediate array configuration
98 (with baselines in the range 15-952 m; see Table 1). The comet was tracked, and the position of the array phase center
99 on the sky was updated in real-time using JPL Horizons orbital solution #K181/6. Weather conditions were very good
100 throughout, with a vertical precipitable water vapor column (PWV) of less than 1 mm. This resulted in good phase
101 stability, which was checked and corrected for with regular visits (every 6–7 minutes) to the nearby phase-calibration
102 quasar J0241-0815.

103 The ALMA correlator was configured to observe three spectral setups in receiver bands 6 and 7, covering lines from
104 HCN (setting 1), HNC and H₂CO (setting 2), and CH₃OH, CH₃CN and CS (setting 3), with spectral resolutions in
105 the range 122–977 kHz (0.1–1.1 km s⁻¹). Integration times on source and other observational parameters are given in
106 Table 1, while the detected spectral line frequencies of interest to the present study are shown in Table 2.

107 **Noisy outlier data points were identified and flagged (removed) through inspection by Joint ALMA**
108 **Observatory (JAO) staff.** The raw data (visibilities) were further flagged and calibrated using the CASA software
109 (version 5.4; [CASA Team et al. 2022](#)), using standard scripts supplied by the JAO. Prior to imaging, the visibilities
110 were continuum-subtracted using a 2nd-order polynomial fit to the line-free channels in each spectral window. The
111 time-resolved interferometric data series was Doppler-corrected to the rest frame of the comet using the CASA `cvel`
112 task, with cubic spline interpolation between the frequency channels.

113 Imaging was performed using the CASA `tclean` (Clark) algorithm with natural weighting. A pixel size of 0.1 $''$
114 was used for the band 6 data and 0.05 $''$ for band 7. Deconvolution of the spatial point spread function (PSF) was
115 performed within an 8 $''$ -diameter circular mask centered on the comet, and stopping at a flux threshold of twice the
116 RMS noise level (σ). The resulting image cubes were corrected for the response of the ALMA primary beam and then
117 transformed from celestial coordinates to sky-projected distances with respect to the center of the comet, which was
118 determined from the peak of the HCN emission (in settings 1 and 2) and the peak CH₃OH emission (in setting 3).
119 The coma gas and continuum (dust plus nucleus) emission peaks were both found to be consistent (within 0.2 $''$) with
120 the JPL Horizons ephemeris position.

121 Autocorrelation (total power) spectra were also extracted from the ALMA data following the method of [Cordiner](#)
122 [et al. \(2019\)](#) and [Cordiner et al. \(2020\)](#). The complete set of autocorrelation scans for all antennas was averaged
123 together to form a single total-power spectrum for each molecule, which was then corrected for atmospheric opacity
124 at the mean elevation angle of the observations, and converted to a flux scale (in Janskys) using the beam size and
125 aperture efficiencies from the ALMA Technical Handbook ([Cortes et al. 2022](#)).

Table 2. Observed spectral line parameters

Species	Transition	Freq. (MHz)	E_u (K)	Res. (kHz)
CH ₃ OH	5 ₀ -4 ₀ E	241700.1590(0040)	48	244
CH ₃ OH	5 ₋₁ -4 ₋₁ E	241767.2340(0040)	40	244
CH ₃ OH	5 ₀ -4 ₀ A^+	241791.3520(0040)	35	244
CH ₃ OH	5 ₄ -4 ₄ A^\pm	241806.5240(0040)	115	244
CH ₃ OH	5 ₋₄ -4 ₋₄ E	241813.2550(0040)	123	244
CH ₃ OH	5 ₃ -4 ₃ A^\pm	241833.1060(0040)	85	244
CH ₃ OH	5 ₂ -4 ₂ A^-	241842.2840(0040)	73	244
CH ₃ OH	5 ₃ -4 ₃ E	241843.6040(0040)	83	244
CH ₃ OH	5 ₋₃ -4 ₋₃ E	241852.2990(0040)	98	244
CH ₃ OH	5 ₁ -4 ₁ E	241879.0250(0040)	56	244
CH ₃ OH	5 ₂ -4 ₂ A^+	241887.6740(0040)	73	244
CH ₃ OH	5 ₋₂ -4 ₋₂ E	241904.1470(0040)	61	244
CH ₃ OH	5 ₂ -4 ₂ E	241904.6430(0040)	57	244
CS	5-4	244935.5565(0028)	35	488
CH ₃ CN	14 ₄ -13 ₄	257448.1282(0002)	207	977
CH ₃ CN	14 ₃ -13 ₋₃	257482.7919(0002)	157	977
CH ₃ CN	14 ₋₃ -13 ₃	257482.7919(0002)	157	977
CH ₃ CN	14 ₂ -13 ₂	257507.5619(0002)	121	977
CH ₃ CN	14 ₁ -13 ₁	257522.4279(0002)	100	977
CH ₃ CN	14 ₀ -13 ₀	257527.3839(0002)	93	977
H ₂ CO	5 _{1,5} -4 _{1,4}	351768.6450(0300)	62	244
HCN	4-3	354505.4773(0005)	43	122
HNC	4-3	362630.3030(0090)	44	244

Note — All spectral line data were obtained from the Cologne Database for Molecular Spectroscopy (CDMS) (Endres et al. 2016). Transition quantum numbers are $J'_{K'}-J''_{K''}$ for CH₃OH and CH₃CN, $J'_{K'_a, K'_c}-J''_{K''_a, K''_c}$ for H₂CO and $J'-J''$ for CS, HCN and HNC. Uncertainties on the trailing digits of the spectral line frequencies are given in parentheses.

3. RESULTS

Spectral line emission was identified with at least 3σ confidence for all the transitions listed in Table 2. For these lines, the spectral channels with fluxes above 2σ were integrated to produce the flux maps shown in Figures 1 and 2. A centrally-peaked morphology is clearly evident for HCN, CH₃OH and CH₃CN (Figure 1), whereas the H₂CO and CS fluxes (Figure 2) are more spatially distributed, lacking any prominent, well-defined emission peak. HNC does not show any significant emission in the interferometric data.

The inset panels in the upper-right corners of Figures 1 and 2 show the spectral line fluxes of each species as a function of cometocentric Doppler velocity, integrated within a $5''$ -diameter circle centered on the comet. For CH₃OH, the inset spectrum represents an average of the five strongest, spectroscopically distinct (unblended) lines, whereas for CH₃CN, the four strongest lines were averaged. The autocorrelation spectra for each species are shown in Figure 3. For HNC, emission is clearly detected in the autocorrelation spectrum, but not in the interferometric data.

In contrast to the interferometric data, which have a spatial resolution ~ 0.3 – $0.6''$ (see Table 1), or 26–38 km at the distance of the comet, and are insensitive to any structures on the sky larger than ~ 3 – $6''$ (260–410 km), the autocorrelation spectra contain flux from the entirety of the ALMA primary beam (which has a FWHM = 16–24''). Due to the extended nature of the cometary coma (spanning hundreds of arc-seconds), the autocorrelation spectra contain intrinsically more flux per beam, especially for gases that have increasing abundances as a function of distance from the nucleus. The weaker, more spatially extended appearance of the H₂CO, CS and HNC emission maps, combined with the relative strength of their autocorrelation spectra, is therefore characteristic of a more extended spatial distribution for these species, consistent with their release in the coma as daughter/distributed species. On




Figure 1. Spectrally-integrated ALMA emission maps of HCN, CH₃OH and CH₃CN, centered on their respective emission peaks. Beam size (FWHM of a Gaussian fit to the PSF) is shown lower left. Inset plots show the molecular spectra on a cometocentric velocity scale, integrated within a 5''-diameter aperture centered on the brightness peak. The (sky-projected) direction of the comet's orbital trail and comet-Sun vector are shown for CH₃CN in the lower right. Five times the RMS noise level (5σ) is indicated on the color bar for each species ($\sigma(\text{HCN}) = 5.0 \text{ mJy km s}^{-1}$, $\sigma(\text{CH}_3\text{OH}) = 7.1 \text{ mJy km s}^{-1}$, $\sigma(\text{CH}_3\text{CN}) = 2.1 \text{ mJy km s}^{-1}$).




Figure 2. Same as Figure 1, but for H₂CO, CS and HNC. The H₂CO and HNC maps are centered on the position of the HCN emission peak, relative to the phase center, while the CS map is centered on the CH₃OH peak. In the inset spectrum, the H₂CO data have been binned over 8 spectral channels for display. The RMS noise level (σ) is indicated on the color bar for each species ($\sigma(\text{H}_2\text{CO}) = 3.0 \text{ mJy km s}^{-1}$, $\sigma(\text{CS}) = 0.5 \text{ mJy km s}^{-1}$, $\sigma(\text{HNC}) = 1.6 \text{ mJy km s}^{-1}$).

145 the other hand, the strongly centrally peaked morphologies for HCN, CH₃OH and CH₃CN are indicative of their
 146 production as parent species, directly from the nucleus. However, considering the observed flux distributions result
 147 from a complex interplay between molecular excitation and emission processes in a three-dimensional outflowing coma,
 148 detailed radiative transfer modeling is required to reliably determine the molecular origins and derive production scale
 149 lengths.

150 4. SUBLIME RADIATIVE TRANSFER MODELING

151 4.1. Model physics and geometry

152 Our modeling approach follows a similar method to the previous comet interferometry studies by Boissier et al.
 153 (2007), Cordiner et al. (2014) and Roth et al. (2021a). Here, the interferometric visibilities are modeled using a non-
 154 LTE radiative transfer code called SUBLIME (SUBLimating gases in LIME, where LIME is the LIne Modeling Engine;
 155 Brinch & Hogerheijde 2010), which takes into account the detailed molecular excitation and emission processes that
 156 occur in the presence of varying coma density, temperature and abundance distributions. The model incorporates a
 157 Haser (1957) radial density profile, and treats coma molecules as parent species, photochemical daughter species, or
 158 a mixture of both, expanding outward at a constant velocity. However, our method differs from the 1D modeling
 159 performed by earlier studies as we adopt a two-component description of the outflowing gases in three dimensions,
 160 similar to that used for the analysis of single-dish CO spectral/spatial observations of comet C/2016 R2 (PanSTARRS)
 161 by Cordiner et al. (2022).




Figure 3. ALMA autocorrelation (total power) spectra, in the rest frame of the comet. The spectra have been baseline-subtracted and corrected for the nominal ALMA beam efficiency (see text). CH₃OH is the mean of the five strongest unblended lines.

162 The coma is divided into two solid-angle regions (Ω_1 , Ω_2), each with an independent water production rate (Q_1 ,
 163 Q_2), outflow velocity (v_1 , v_2) and kinetic temperature as a function of radius ($T_1(r)$, $T_2(r)$). As shown in Appendix
 164 A Figure 8, the first solid angle region (Ω_1) is defined by a cone of half-opening angle θ_{jet} , with its apex at the center
 165 of the nucleus, and its axial vector pointing at a phase angle ϕ with respect to the observer, and at a position angle
 166 ψ in the plane of the sky. The second region (Ω_2) represents the remaining (ambient) coma. This geometry, although
 167 still likely to be highly simplified compared to reality, represents an evolution from the two-hemisphere coma model of
 168 Roth et al. (2021b), which was successfully used for the analysis of asymmetric CH₃OH spectra observed in 46P using
 169 the ALMA 7 m array (ACA). As explained by Cordiner et al. (2022), the SUBLIME model provides a sufficiently
 170 good approximation to the structure of a coma with a single dominant, rapidly expanding (near-) sunward-facing jet,
 171 embedded in a slower-moving, ambient coma. The adopted geometry is found to reproduce very well the spectral
 172 line profiles (including asymmetries) observed for all our detected species, while keeping the number of variable model
 173 parameters at a minimum.

174 The radial distribution of a daughter species is governed by its parent scale length (L_p), which, in a uniformly-
 175 expanding coma of outflow velocity v , is related to the photodissociation rate (Γ) of its parent species by $L_p = v/\Gamma$.
 176 For consistency, we assume that Γ is constant for a given parent species across both coma solid angle regions (Ω_1 ,
 177 Ω_2), so the ratio of production scale-lengths for a daughter species in those two regions is determined by the ratio of
 178 outflow velocities (v_1/v_2). Consequently, rather than reporting parent scale lengths for both regions, we report only the
 179 L_p value corresponding to the jet component (Ω_1). Outflow velocities for daughter species in our model are allowed
 180 to differ from the parents. Daughter production rates in the two coma solid angle regions are also independently
 181 optimized; reported daughter abundances are therefore taken as averages over the entire coma.

182 The excitation calculation in our model is time-dependent and takes into account radiative cooling of rotational
 183 levels, (de-)excitation by collisions with H₂O and electrons, and pumping by the Solar radiation field as the gas moves
 184 outward (for more details see Cordiner et al. 2022). The impact of coma opacity on the molecular excitation is negligible
 185 for the observed species in this comet, so radiation trapping effects are not included. We adopt H₂O–HCN collision
 186 rates from Dubernet & Quintas-Sánchez (2019) (also assumed to apply to HNC), while collision rates of H₂O with
 187 CH₃OH, H₂CO, CH₃CN and CS are assumed to be the same as for H₂, and have been taken from the LAMDA database
 188 (van der Tak et al. 2020). An electron density scaling factor of $x_{ne} = 0.2$ is used, following the recommendations of
 189 Hartogh et al. (2010) and Biver et al. (2019, 2021). Pumping rates for HCN, HNC, H₂CO, and CS were calculated
 190 as described by Cordiner et al. (2019) and Roth et al. (2021b), based on the method of Crovisier & Encrenaz (1983),
 191 with rovibrational transition data from the latest versions of the HITRAN and GEISA catalogues (Gordon et al.

2022; Delahaye et al. 2021). CH₃OH pumping rates were calculated similarly. However, the Einstein A coefficients for CH₃OH in the HITRAN catalogue were found to be incorrect because they were calculated considering corrupted statistical weights and partition functions. On the other hand, HITRAN line intensities at 296 K are accurate, so by converting these to A coefficients, employing accurate statistical weights and partition functions (Villanueva et al. 2012), the required spectroscopic information could be recovered. We further complemented the HITRAN data by adding rovibrational transitions from the strong v_2 , v_3 and v_9 bands from the quantum band models of Villanueva et al. (2012). The data were then homogenized based on the CH₃OH quantum numbers and energies reported for the ground-state Hamiltonian by Mekhtiev et al. (1999).

4.2. Model optimization strategy

Our ALMA observations were obtained over two dates: 2018-12-02 and 2018-12-07 (see Table 1). On 2018-12-02, the coma physical structure (characterized by the ratio of production rates in the sunward jet *vs.* ambient coma, Q_1/Q_2 , and the jet properties, θ_{jet} , ϕ and ψ) was determined using a model fit to the bright HCN $J = 4 - 3$ line observations. On 2018-12-07, multiple lines from the CH₃OH ($J_K = 5_K - 4_K$) band were used instead, since HCN was not observed on this date. Both these species correlate well with H₂O (Dello Russo et al. 2016a; Bockelée-Morvan & Biver 2017), and therefore provide a reasonable proxy for the overall coma outflow velocity and Q_1/Q_2 ratio in the absence of spectrally-resolved H₂O data. The H₂O coma physical structure derived from fitting the HCN data was therefore used in our models for H₂CO and HNC (also obtained on 2018-12-02), whereas the structure derived using CH₃OH was applied to CH₃CN and CS. For CH₃CN, the observed ALMA data were of insufficient spectral resolution to adequately constrain the values of v_1 and v_2 , whereas for H₂CO, HNC and CS, the correlation with H₂O is insufficiently demonstrated, and the data were of insufficient S/N to reliably constrain the coma structure by themselves.

The total H₂O production rate on each date was obtained from a linear fit to the measurements between 2018-12-02 and 2018-12-10 by Combi et al. (2020) using Ly- α observations by the Solar and Heliospheric Observatory (SOHO) satellite, which gave $Q(\text{H}_2\text{O}) = 6.1 \times 10^{27} \text{ s}^{-1}$ on 2018-12-02 and $Q(\text{H}_2\text{O}) = 7.2 \times 10^{27} \text{ s}^{-1}$ on 2018-12-07. These values are consistent with a slow increase in $Q(\text{H}_2\text{O})$ as the comet approached perihelion on 2018-12-12, and are in line with the average values of $8 \times 10^{27} \text{ s}^{-1}$ measured by Lis et al. (2019) using the Stratospheric Observatory for Infrared Astronomy (SOFIA) between 2018-12-14 and 2018-12-20, and $7 \times 10^{27} \text{ s}^{-1}$ measured using IRTF between 2018-12-06 and 2018-12-21 (Khan et al. 2023).

Our models were set up on a 3D Delaunay grid containing 10,000 points distributed pseudo-randomly with a density of points proportional to the logarithm of the radial distance (r) from the center of the nucleus (excluding the nucleus itself). The model domain therefore consisted of a spherical coma region extending from the surface of the nucleus (assumed to be a sphere of radius 500 m), to an outer boundary at $r = 2 \times 10^5 \text{ km}$. The outer boundary was chosen to be large enough so that further increasing it had no significant impact on our model results. Raytracing was performed along the line-of-sight vectors through each grid point, on a frequency grid with a uniform channel spacing of 100 m s^{-1} , which was then interpolated in two (spatial) dimensions onto the image grid, which consisted of 768×768 pixels $0.1''$ in size. The innermost 4×4 pixel region of the image was further super-sampled using a 30×30 point Cartesian grid, to accurately capture the nonlinear behavior of the coma flux on the smallest relevant radial scales. To simulate the response of the ALMA primary beam, each plane of the synthetic image cubes was multiplied by a 2D Gaussian of $\text{FWHM} = 1.13\lambda/D$, where λ is the wavelength and D is the antenna diameter. For each molecule, the frequency axis of the resulting synthetic image cube was convolved to the spectral resolution of the ALMA observations, followed by cubic spline interpolation to the observed frequency grid. This allowed a channel-by-channel comparison of the model with the observations.

Interferometric observations inherently suffer from incomplete spatial sampling, and the resulting, Fourier-transformed images can be subject to artifacts introduced by gridding, interpolation, and numerical deconvolution. To facilitate accurate modeling of the data, and avoid our model fits becoming biased by image artifacts, we chose to perform all model fitting in the Fourier domain. This requires taking the Fourier transform of the simulated coma image cubes, then sampling each spectral channel with the same set of uv points (antenna baseline lengths and orientations) as the observations, which was performed using the `vis_sample` code (Loomis et al. 2018). To make the problem computationally tractable, the observed visibilities were first averaged along the time-axis to produce a single (complex) visibility point per baseline, per channel. The chi-squared statistic

$$\chi^2 = \sum_{i,j} \left(\frac{(\Re(V_{i,j}^{obs}) - \Re(V_{i,j}^{mod}))^2}{\sigma_{\Re}^2} + \frac{(\Im(V_{i,j}^{obs}) - \Im(V_{i,j}^{mod}))^2}{\sigma_{\Im}^2} \right) + \sum_i \frac{((S_i^{obs}) - (S_i^{mod}))^2}{\sigma_S^2} \quad (1)$$

was minimized using `lmfit` (Newville et al. 2016), by application of the Levenberg-Marquardt algorithm. Summation of the interferometric residuals (the difference between V^{obs} for the observations, and V^{mod} for the model) was performed over the set of baselines $j = N(N-1)$, where N is the number of ALMA antennas (43), and i is the number of spectral channels. Due to the presence of coma asymmetries, both the real (\Re) and imaginary (\Im) parts of the complex visibilities were included in the χ^2 calculation. The difference between the model total power spectra (S_i^{mod}) and the observed autocorrelation spectra (S_i^{obs}), provide additional, strong constraints on the production scale length (L_p) of each species due to the larger angular scales probed by these data. During fitting, the residuals were scaled by the respective standard deviations of the real and imaginary visibilities, and of the autocorrelation spectra (σ_{\Re} , σ_{\Im} and σ_S , respectively), which were calculated from line-free data regions adjacent to each spectral line.

To allow for errors in the comet ephemeris coordinates, a positional (RA, dec.) offset of the model origin from the image center was included as a further pair of free parameters in the model fits. Statistical uncertainties on all model parameters were obtained from the diagonal elements of the `lmfit` covariance matrix.

4.3. Coma temperature structure derived from CH_3OH visibilities

Model optimization was performed first for CH_3OH , because the multiple transitions observed for this species span a broad range of upper-state energy levels (see Table 2), which allow its rotational excitation state to be determined, from which the coma kinetic temperature is derived (see Cordiner et al. 2017b or Biver et al. 2021, for example). Our method differs from previous studies, however, due to our model's ability to interpret the coma temperature structure in three dimensions, by fitting the variations in individual CH_3OH spectral line channels (V_i , S_i) as a function of spatial (uv) coordinate.

We began with the simplest assumption of a constant kinetic temperature (T) as a function of radius, then added complexity to the $T(r)$ profile until a good fit to the data was obtained. This strategy keeps the number of free parameters at a minimum (therefore keeping the χ^2 minimization computationally feasible), and ensures that there are enough degrees of freedom in the model to reproduce the data, but not so many that the model becomes ill-constrained. To obtain a good fit to the observations, it was necessary to implement different temperature profiles as a function of radius ($T_1(r)$, $T_2(r)$), within the two different coma solid angle regions. Temperature variability was implemented using a segmented linear function (in $\log(r)$ - T space, within the domain $r = \theta_{min}$ to θ_{PB}), with a variable number of segments (n), of equal length (l_s) in $\log(r)$ space. The segmented function was smoothed (in $\log(r)$ space) by convolving it with a Gaussian of FWHM equal to l_s . The temperature was set constant inside a radius corresponding to half the minor axis of the ALMA beam ($r = \theta_{min}/2$), and outside a radius corresponding to half the primary beam FWHM ($r = \theta_{PB}/2$). A good fit to the entire CH_3OH dataset was obtained using $n = 5$ variable points in the temperature profiles as a function of radius (see Figure 4 and Appendix B, Figure 9).

Due to the high S/N of our CH_3OH observations, the best-fitting coma model is tightly constrained by the data, and has a jet half-opening angle of $\theta_{jet} = 70 \pm 5^\circ$, with a phase angle of $\phi = 38 \pm 1^\circ$ and position angle of $\psi = 33 \pm 1^\circ$. The resulting model jet emanates from close to (within 10° of) the sub-solar point on our (assumed) spherical nucleus, and is therefore consistent with preferential outgassing in the general sunward direction. The best-fitting jet outflow velocity is $v_1 = 0.729 \pm 0.002 \text{ km s}^{-1}$, with the remaining (ambient/nightside) coma expanding at $v_2 = 0.395 \pm 0.003 \text{ km s}^{-1}$. We initially attempted to fit the observed ALMA data assuming CH_3OH was solely a parent species, but a significant (15σ) improvement in the final χ^2 value was obtained using a composite (parent + daughter) model. Independent optimization of the parent and daughter CH_3OH production rates resulted in approximately equal abundances for both components of $1.2 \pm 0.1\%$ (relative to H_2O), with a best-fitting parent scale length of $L_p = 36 \pm 7 \text{ km}$ for the CH_3OH daughter (in the jet component, Ω_1). The presence of an additional CH_3OH daughter component implies the production of significant amounts of CH_3OH in the near-nucleus coma, likely from the sublimation of icy grains (see Section 5.2). We also attempted to fit the CH_3OH data using a pure daughter model, but this resulted in a statistically worse fit at the 11σ level, so our modeling strongly implies the presence of both nucleus (parent) and coma (daughter) sources for CH_3OH in comet 46P (see also Section 4.4).

A comparison between the best-fitting modeled and observed visibility spectra is shown in Appendix B Figure 9, which includes the total power (autocorrelation) spectrum at the top, followed below by the real part of the interferometric visibilities, binned within successive 20 m baseline ranges. The angular scales probed decrease with




Figure 4. Variation of the 46P coma kinetic temperature with radius in the sunward (jet) and antisunward (ambient) coma regions, derived from modeling the $\text{CH}_3\text{OH } J = 5 - 4$ data. Shaded regions indicate the $\pm 1\sigma$ error envelopes.

291 increasing baseline length, from the $24''$ scale of the autocorrelation spectrum, down to the $0.6''$ scale probed by the
 292 longest baselines. We show only the real part of the visibilities since the imaginary components are weak and noisy.

293 As shown in Figure 4, the average temperature on the sunward (jet) side of the nucleus is significantly lower than the
 294 anti-sunward (ambient) side. This is particularly true closest to the nucleus (within $r = 19$ km), where $T_1 = 59 \pm 3$ K
 295 and $T_2 = 105 \pm 5$ K, amounting to almost a factor of two difference. The temperature in the sunward jet rises with
 296 increasing cometocentric distance, reaching a peak with $T_1 = 79 \pm 5$ K at $r = 121$ km (compared with $T_2 = 71 \pm 5$ K
 297 in the ambient coma at the same radius), before falling smoothly towards $T_1 = 25 \pm 4$ K ($T_2 = 49 \pm 6$ K) at large radii
 298 ($r > 1500$ km). The different temperature behavior on opposite sides of the nucleus is remarkable, since it implies
 299 significant differences in the balance of heating and cooling mechanisms on the sunward and anti-sunward sides of the
 300 comet. Biver et al. (2021) also identified cooler gas on the day side of the nucleus than the night side (57 K vs. 71 K)
 301 based on single-dish CH_3OH observations probing coma radial distances $\lesssim 600$ km. This is qualitatively similar to our
 302 result, and was explained by Biver et al. (2021) as being due to more efficient adiabatic cooling on the sunward side
 303 (for further discussion see Section 5.2).

304 The observed transitions of CH_3CN from different K levels provide an additional measure of the coma temperature.
 305 However, due to the lower spectral resolution and S/N, it was not possible to reliably constrain the spatial distribution
 306 of temperatures for this species. Assuming a constant temperature throughout the coma, we found $T = 80 \pm 8$ K using
 307 the CH_3CN data.

308 4.4. Visibility modeling to derive parent scale lengths

309 Adopting the best-fitting coma kinetic temperature distribution from our CH_3OH modeling, we proceeded to optimize
 310 the remaining free model parameters for HCN, HNC, CS, H_2CO , and CH_3CN . As a result of lower S/N and spectral
 311 resolution for CS and CH_3CN (simultaneously observed with CH_3OH), we employed the same jet opening angle (70°)
 312 and ratio of H_2O production rates as derived for CH_3OH ($Q_1/Q_2 = 1.21$). The HCN jet opening angle (70°) and
 313 Q_1/Q_2 ratio (1.33) was employed for modeling the HNC and H_2CO data. The best-fitting HCN coma outflow velocities
 314 were $v_1 = 0.741 \pm 0.001$ km s^{-1} and $v_2 = 0.443 \pm 0.001$ km s^{-1} , with a jet phase angle $\phi = 44 \pm 1^\circ$ and position angle
 315 $\psi = 37 \pm 1^\circ$, consistent with preferential outgassing in the sunward direction. The jet axis was therefore fixed along
 316 the comet-sun vector for all species apart from CH_3OH . Model fits to the binned visibility spectra for all molecules are
 317 shown in Appendix B (Figures 9 to 14). Since the visibility data for CH_3CN , H_2CO , CS and HNC are noisier than for
 318 CH_3OH and HCN (particularly on large baselines), some of the longer baseline ranges for which no signal is evident
 319 have been omitted from Figures 11 to 14.

320 To visualize the radial flux distribution for each molecule, in Figures 5 and 6 we plot the real part of the interferometric
 321 visibility as a function of baseline length (uv distance). The observed and modeled fluxes are averaged across the
 322 spectral channels containing line emission, and the interferometric data have been further averaged into 10 m-wide uv




Figure 5. Observed interferometric visibilities for HCN, CH₃OH and CH₃CN, with best-fitting models overlaid. The observations and model data have been (spatially) averaged within 10 m uv bins and (spectrally) averaged over all line emission channels for each species, including 13 lines for CH₃OH and 5 lines for CH₃CN. The zero-spacing data points (at $uv = 0$) were taken from the ALMA autocorrelations (observed simultaneously with the interferometric data). Preferred, best-fitting visibility models are shown with solid lines, while dashed and dotted curves show models that are ruled out based on a poorer quality of fit (or other criteria; see text).




Figure 6. Same as Figure 6, for H₂CO, CS and HNC.

bins. Statistical error bars on the interferometric data are typically very small due to the large number of data points being averaged together, whereas the total power data points have an additional 10 % error added in quadrature with the statistical error, to allow for uncertainty in the aperture efficiency of the ALMA antennas. For HCN, a parent outgassing model provides the best fit to the data. In an attempt to better fit the total power ($uv = 0$) data point, a daughter distribution of HCN was added, and the model parameters were re-optimized (with variable parent and daughter abundances, and variable HCN parent scale length L_p). However, this did not significantly improve the χ^2 value, and the optimized L_p value was found to be 5 ± 16 km, which is consistent with zero, showing that HCN is likely a parent species.

The binned CH₃OH visibilities and corresponding model fits are shown in Figure 5 (middle panel). As explained in Section 4.3, in contrast to HCN, a parent + daughter model provides the best fit for CH₃OH (with $L_p = 36 \pm 7$ km for the daughter component). The difference between the CH₃OH “parent” and “parent + daughter” models is difficult to see in Figure 5; unbinned, zoomed CH₃OH visibilities are therefore shown in Appendix B, Figure 15, where the improved fit for the parent + daughter model is evident. For CH₃CN, a parent model fits the data very well. Addition of a CH₃CN daughter component slightly improved the fit (resulting in $L_p = 101^{+194}_{-101}$ km for the daughter), but the associated drop in χ^2 value corresponded to only 1.6σ , so the improvement in fit quality was not statistically significant.

The binned visibility data for H₂CO, CS and HNC are plotted in Figure 6, and as shown by the overlaid model curves, these three species can only be well fit using daughter models (their respective best-fitting parent scale lengths are given in Table 3). To obtain the best visibility fits for these daughter species, we allowed their outflow velocities to differ from those of the underlying H₂O distribution (determined from our HCN and CH₃OH models), which may be physically justified if they originate from a non-nucleus source. We also attempted to fit these data using ‘parent only’ models, by setting $L_p = 0$ and optimizing the abundance to obtain the best fit to the observations. However, in all three cases, a parent model provides a clearly inadequate fit, particularly at the largest scales probed by the total power ($uv = 0$) data. For CS, we also tried to fit the observations assuming production of this species from CS₂ photolysis (at a photodissociation rate $\Gamma = 2.61 \times 10^{-3}$, which corresponds to $L_p = 280$ km in the jet and $L_p = 152$ km




Figure 7. Parent scale-length χ^2 analysis for our observed species in 46P, showing the change in chi-square value ($\Delta\chi^2$), as a function of parent scale-length (L_p). Dotted horizontal lines show the $\Delta\chi^2 = 1$ threshold (1σ), corresponding to 68 % confidence, and dashed lines show the $\Delta\chi^2 = 6.63$ threshold (2.6σ), corresponding to 99 % confidence.

Table 3. 3D visibility modeling results

Species	Abundance ^a (%)	v_1 (km s ⁻¹)	v_2 (km s ⁻¹)	L_p (km)
HCN	$0.1176^{+0.0003}_{-0.0003}$	0.741 ± 0.001	0.443 ± 0.001	< 3
CH ₃ OH ^b	$2.7^{+0.1}_{-0.1}$	0.729 ± 0.002	0.395 ± 0.002	$0 / 36 \pm 7$
CH ₃ CN	$0.011^{+0.002}_{-0.002}$	0.729^c	0.395^c	< 18
H ₂ CO	$0.153^{+0.031}_{-0.023}$	0.65 ± 0.02	0.47 ± 0.02	876^{+250}_{-175}
CS	$0.022^{+0.004}_{-0.002}$	0.66 ± 0.02	0.37 ± 0.02	675^{+159}_{-124}
HNC	$0.0054^{+0.0232}_{-0.0018}$	0.69 ± 0.05	0.34 ± 0.05	$1412^{+14,876}_{-367}$

^a Total production rate relative to H₂O. ^b CH₃OH best-fitting model has both nucleus (parent) and coma (daughter) production; the reported abundance is a sum over both components.

^c Held fixed at the CH₃OH values (simultaneously observed) due to the low spectral resolution for CH₃CN.

347 in the ambient coma; Huebner & Mukherjee 2015). As shown by the dashed orange curve, the CS₂ photolysis model
 348 produces insufficient CS flux on large scales (small uv values) to reproduce the observations. The final, best fitting
 349 model parameters for all species are given in Table 3.

350 Due to the inverse exponential behavior of cometary molecular densities as a function of radius, uncertainties on
 351 the parent scale lengths can be highly asymmetric, and are therefore not adequately represented using the symmetric
 352 error bars obtained from the `lmfit` covariance matrix. To address this issue, we calculated the χ^2 surface for each
 353 species as a function of L_p (with the abundance as a variable, but other model parameters held fixed), and plotted the
 354 resulting $\Delta\chi^2(L_p)$ curve, with cubic spline interpolation between points (with $\Delta\chi^2(L_p) = \chi^2(L_p) - \chi_m^2$, where χ_m^2 is
 355 the minimum value). The results are shown in Figure 7, with dotted horizontal lines to show the 1σ (68 % confidence)
 356 $\Delta\chi^2$ thresholds, and dashed lines to show the 2.6σ (99 % confidence) thresholds.

357 For HCN, the χ^2 minimum is for $L_p = 0$, confirming that this species is a parent. For CH₃CN, the smallest χ^2
 358 value occurs around $L_p = 2.5$ km, but this minimum is much shallower than 1σ , and is therefore not significant. The
 359 CH₃CN $\Delta\chi^2$ curve crosses the 99% confidence threshold at $L_p = 18$ km, which represents a strict upper limit on the

scale of possible coma production for this species assuming a single (distributed) source. CH₃OH, on the other hand, shows a well defined χ^2 minimum at $L_p = 13$ km. However, as explained in Section 4.3, a composite (parent + daughter) model provides the best fit for this species (with a lower χ^2 value than the pure daughter model plotted here); the L_p value of the daughter component in the CH₃OH composite model is well constrained as 36 ± 7 km, so this is reported in Table 3 instead.

Assuming a scale length of $L_p = 36$ km for a sublimating icy grain source (*i.e.* the same as derived for CH₃OH), the abundance upper limit (at the 99% confidence level) for icy grain production of HCN is < 0.001 % relative to H₂O. This corresponds to < 1 % of the total HCN. The associated upper limit for icy grain production of CH₃CN is < 0.01 %, which is comparable to the measured CH₃CN parent abundance, and shows that a significant source of CH₃CN production from icy grains in the coma, close to the nucleus, cannot be ruled out by our data.

H₂CO, CS and HNC all exhibit well-defined χ^2 minima at substantially larger L_p values than the other observed species, so these molecules are confirmed as originating primarily in the coma of comet 46P, rather than from the nucleus. While the H₂CO and CS L_p values are well constrained (at 876^{+250}_{-175} km and 675^{+159}_{-124} km, respectively), the HNC uncertainty interval is much larger and highly asymmetric, with an L_p value lying between 1045 km and 14,880 km (at 1σ confidence). At the stricter, 99 % confidence level, the L_p (HNC) value is only constrained by a lower limit of > 218 km.

5. DISCUSSION

5.1. HCN

The high spatial resolution of ALMA combined with the close geocentric distance of 46P places unusually tight constraints on the production scale length of HCN. With an upper limit of $L_p < 3$ km, it is likely that HCN originates primarily from outgassing by the nucleus of the comet. At the outflow velocity of 0.74 km s^{-1} (in the sunward jet), any HCN daughter production would need to occur on a timescale of < 4 s, which is significantly shorter than the lifetimes of any known coma molecules (typically in the range 10^3 – 10^6 s; Huebner & Mukherjee 2015). Our scale length upper limit is consistent with (although significantly better constrained than) the previous upper limits of $L_p < 50$ km in C/2012 F6 (Lemmon) and C/2012 S1 (ISON) (Cordiner et al. 2014) and $L_p < 100$ km in C/2015 ER61 (Roth et al. 2021a), which were obtained using ALMA and Atacama Compact Array (ACA) data. Including the IRAM Plateau de Bure interferometric study of comet Hale-Bopp (Wink et al. 1997) and the Keck NIRSPEC study of C/2014 Q2 (Lovejoy), the body of evidence demonstrating HCN release from direct sublimation of nucleus ices, with no clear evidence for production in the coma, is now substantial. Coma HCN abundance measurements in fully-activated comets thus provide a valid proxy for the abundance of this molecule in the nucleus ices.

Our derived HCN abundance is 0.1176 ± 0.0003 % with respect to H₂O, with the error bar accounting only for statistical uncertainties. Adopting a conservative estimate for the ALMA flux scale uncertainty of 10 %, and an extra uncertainty of 10 % in the H₂O production rate (added in quadrature), the abundance error increases to ± 0.016 %. Our measured HCN abundance in comet 46P is therefore consistent with the value of 0.11 ± 0.01 % obtained by Biver et al. (2021) within the same month, and matches the average value observed previously in OCCs and JFCs using radio spectroscopy (Bockelée-Morvan & Biver 2017).

5.2. CH₃OH

Our observations reveal the presence of two coexisting outgassing modes for CH₃OH, consisting of (1) production directly from the nucleus and (2) production from a near-nucleus coma source. Our best-fitting SUBLIME model indicates that both sources are responsible for similar amounts of CH₃OH production, with best-fitting abundances (with respect to H₂O) of 1.19 ± 0.13 % for the nucleus (parent) source of CH₃OH gas and 1.18 ± 0.13 % for the distributed (coma) source.

The presence of icy grain sublimation has been known to be an important source of gases in cometary comae since the EPOXI mission to comet 103P/Hartley 2 (A'Hearn et al. 2011). Using infrared spectroscopy, Protopapa et al. (2014) observed $1 \mu\text{m}$ sized water ice grains in the coma of 103P, while Kelley et al. (2013) observed a halo of larger, longer-lived ice chunks (with sizes ≥ 1 cm) surrounding the nucleus at distances up to ~ 40 km. Using a detailed DSMC coma model, Fougere et al. (2013) determined that 77 % of the H₂O outgassing in 103P originates from sublimation of icy grains in the coma. Such an excess of H₂O production compared with its small nucleus size, led to 103P being designated as a ‘hyperactive’ comet.

46P/Wirtanen is also hyperactive (Lis et al. 2019; Moulane et al. 2023), and the large H₂O production rate for its nucleus size can, at least in part, be attributed to the presence of icy grains in the coma (Bonev et al. 2021). Knight et al. (2021) explained the different OH and CN morphologies in 46P as being partly due to OH production from an icy-grain H₂O source. However, Protopapa et al. (2021) exclude pure H₂O ice grains in the coma of this comet. Based on the non-detection of H₂O ice absorption features, these authors suggested two possible explanations for the hyperactivity of 46P: icy grains ($\sim 1 \mu\text{m}$ in size) containing a small amount of low-albedo dust, or larger chunks ejected from the nucleus, containing significant amounts of water ice. Previous ALMA observations by Cordiner et al. (2017b) identified nucleus outgassing as the primary source of CH₃OH in the coma of C/2012 K1 (PanSTARRS). On the other hand, a distributed CH₃OH source (with a scale length ~ 550 km) was observed using the James Clerk Maxwell Telescope (JCMT) in comet 252P/LINEAR, and was attributed to CH₃OH production from icy grains in the coma (Coulson et al. 2017). It is therefore reasonable to attribute the source of our observed CH₃OH daughter distribution in 46P (around 40 km from the nucleus) to the presence of sublimating icy grains in the coma, composed of a mixture of CH₃OH, H₂O and dust. The sublimation rate of coma icy grains depends on their size and composition. Based on the calculations of Beer et al. (2006), and assuming that the gas and dust outflow velocities are coupled, our derived lifetime of 496 ± 97 s for the CH₃OH-producing grains in 46P implies a dirty ice grain size of $\sim 10 \mu\text{m}$. If the dust is moving outward slower than the gas, then the grains could be larger.

The presence of sublimating icy grains can significantly impact the coma temperature, as shown by the direct simulation Monte Carlo (DSMC) models of Fougere et al. (2012) and Fougere (2014), due to the transfer of excess kinetic energy from sublimated molecules to the surrounding gas. In the absence of such a mechanism, the gas kinetic temperature in coma physical models (e.g. Rodgers et al. 2004; Tenishev et al. 2008) falls rapidly with distance from the nucleus, from $\gtrsim 100$ K at $r = 0$ to ~ 10 K at $r = 100$ km, due to quasi-adiabatic expansion. Such steeply declining temperatures are at odds with coma observations. For example, the observations of comets 73P-B/Schwassmann–Wachmann, 103P/Hartley 2, C/2012 S1 and 46P/Wirtanen (Bonev et al. 2008, 2013, 2014, 2021) and C/2012 K1 (Cordiner et al. 2017b) revealed a shallower temperature decay — and in some cases, increasing temperatures — as a function of radius, attributable to sublimative heating. Spatially resolved temperature observations therefore provide an indirect probe for the presence of icy grains in cometary comae.

Our derived 46P coma kinetic temperature profile (Figure 4) shows an initial increase with radius on the sunward side of the nucleus, followed by a relatively slow decrease, consistent with strong coma heating due to the sublimation of icy grains within a few hundred kilometers of the nucleus. On the antisunward side, the temperature drop is also less steep than predicted by hydrodynamical and DSMC models for other comets, and flattens out to a higher temperature than predicted without the presence of icy grain heating (Rodgers et al. 2004; Tenishev et al. 2008; Fougere et al. 2012). This temperature behavior therefore suggests the presence of significant icy grain sublimation in the 46P coma, consistent with conclusions based on spatially resolved infrared observations of H₂O in this comet (Bonev et al. 2021). Dedicated theoretical modeling will be required to further investigate this hypothesis.

Our total CH₃OH abundance of 2.7 ± 0.1 % including the nucleus and coma sources is consistent (at the 2σ level) with the value of 3.03 ± 0.23 % observed by Bonev et al. (2021) on 2018-12-18, but somewhat smaller than the value of 3.38 ± 0.03 % observed by Biver et al. (2021) during December 11–18, and larger than the value of 1.6 ± 0.1 % observed by Bergman et al. (2022) on Dec. 22–28. These differences could be explained by temporal variability of the CH₃OH and H₂O outgassing rates, since significant short-timescale variations in $Q(\text{CH}_3\text{OH})$ (over a period of several hours to several days) were observed in this comet by Roth et al. (2021b). Indeed, our CH₃OH production rate of $(1.9 \pm 0.1) \times 10^{26} \text{ s}^{-1}$ on 2018-12-07 is very close to the value of $(2.2 \pm 0.1) \times 10^{26} \text{ s}^{-1}$ observed on 2018-12-06 by Khan et al. (2023), and $(2.1 \pm 0.1) \times 10^{26} \text{ s}^{-1}$ on 2018-12-11 (Biver et al. 2021), demonstrating good consistency between the ALMA, IRTF and IRAM measurements in this case.

The presence of an icy grain source for CH₃OH, but not HCN implies that the icy grains are chemically distinct from the bulk (sublimating) ice within the nucleus. Drahus et al. (2012) also inferred the presence of icy grain sources of CH₃OH and HCN in comet 103P, and concluded that CH₃OH is more intimately mixed with H₂O ice than HCN, with a larger abundance of CH₃OH in the icy grains. The apparent dichotomy between the abundances of CH₃OH and HCN in the two cometary ice storage reservoirs (icy grains *vs.* bulk nucleus) was interpreted by Drahus et al. (2012) as due to thermal evolution of the nucleus. A primordial origin in the protoplanetary disk or interstellar cloud, prior to accretion of the comet, is also possible. Spatial differentiation of O and CN-rich molecules between gas and ice phases has been observed in disks (Bergner et al. 2018; Booth et al. 2021; Öberg & Bergin 2021), indicating an active carbon chemistry occurs in the gas phase, while O-bearing species such as CH₃OH and H₂O remain largely frozen on

461 grains. Non-uniform mixing of these distinct chemical reservoirs during comet accretion could lead to the observed
 462 spatial heterogeneity of the CH₃OH/HCN ratio within the nucleus.

463 5.3. CH₃CN

464 A production scale length upper limit of $L_p < 18$ km was derived for CH₃CN, based on our visibility modeling.
 465 The ALMA data show no significant evidence for production of this molecule as a coma daughter species (either
 466 from photolysis of a parent molecule, or from sublimation of icy grains), so we conclude that the primary source of
 467 CH₃CN is from direct sublimation of molecular ices in the nucleus. The abundance of this molecule (0.011 ± 0.002
 468 %) is consistent with, but toward the lower end of the range of values (0.008–0.054 %) previously observed in comets
 469 (Bockelée-Morvan & Biver 2017).

470 This is the most complex nitrile confirmed to be present in cometary ice to-date. It has long been detected in the
 471 gas phase in the interstellar medium and protostellar envelopes (Herbst & van Dishoeck 2009), while recent studies
 472 using ALMA have found CH₃CN to be widespread in protoplanetary disks where it is believed to be formed primarily
 473 *via* chemistry on grain surfaces (Öberg et al. 2015; Bergner et al. 2018; Ilee et al. 2021), before being thermally-
 474 desorbed into the gas phase where it can be observed. CH₃CN has not been detected so-far in interstellar ices, but our
 475 measured abundance in 46P is consistent with the upper limit of $\lesssim 2$ % (with respect to H₂O) found recently along
 476 two interstellar sightlines using the James Webb Space Telescope (McClure et al. 2023).

477 The CH₃CN/HCN ratios we measure in 46P are consistent with, or somewhat higher than the values observed
 478 in nearby protoplanetary disks by Bergner et al. (2018) and Ilee et al. (2021), implying a likely genetic relationship
 479 between protoplanetary disk and cometary nitriles. Our observed CH₃CN/H₂O ratio also matches that found in the
 480 (gas plus ice) phase of the GM Aur planet-forming disk by Ilee et al. (2021), but is ~ 8 times lower than that found
 481 in the AS 209 disk. The utility of such comparisons is limited, however, due to the action of gas-phase chemical
 482 processes, which can modify the CH₃CN/HCN/H₂O ratios found in the disk gas compared with those in the ice.
 483 Cometary observations therefore remain as a perhaps more useful probe of the abundances of complex nitriles in the
 484 ice reservoir of our solar system’s planet-forming disk.

485 5.4. H₂CO

486 Our measured H₂CO production scale length of $L_p = 876^{+250}_{-175}$ km at $r_H = 1.07$ is compatible with the values
 487 measured previously using ALMA of 2200^{+1100}_{-800} km at $r_H = 1.17$ au in C/2015 ER61 (PanSTARRS) (Roth et al.
 488 2021b), 1200^{+1200}_{-400} at $r_H = 1.47$ au in C/2012 F6 (Lemmon) and 280 ± 50 at $r_H = 0.54$ au in C/2012 S1 (ISON)
 489 (Cordiner et al. 2014). When scaled by r_H^{-2} , these L_p values are in reasonably close agreement, consistent with H₂CO
 490 production by photodissociation of a (molecular) parent species at a rate $\sim 7 \times 10^{-4} \text{ s}^{-1}$ (at $r_H = 1$ au), in an
 491 optically thin coma. However, Biver et al. (1999) measured a much larger L_p value of $7000r_H^{-1.5}$ km in comet C/1996
 492 B2 (Hyakutake) using a single-dish radio telescope, and Meier et al. (1993) derived $L_p \sim 3600$ km in 1P/Halley at
 493 $r_H = 0.89$ au *via* in-situ mass spectrometry. CH₃OH photolysis is expected to produce H₂CO in the outer coma, but
 494 cannot be responsible for the observed H₂CO in these comets, since it occurs at a rate of $1.0 \times 10^{-5} \text{ s}^{-1}$ at $r_H = 1.06$ au
 495 (Huebner & Mukherjee 2015; see also Heays et al. 2017), which is almost two orders of magnitude smaller than the
 496 H₂CO parent photodissociation rate required to fit our ALMA observations. Indeed, the corresponding CH₃OH
 497 dissociation scale in the 46P sunward jet is 73,000 km, which is much larger than the maximum angular scale of
 498 1400 km spanned by the ALMA primary beam FWHM. Alternative H₂CO parents must therefore be sought.

499 The absence of other known C,H,O-bearing molecules with sufficient abundances has led to the idea that H₂CO
 500 may be released in the coma from the thermal breakdown of organic-rich dust particles. Formaldehyde polymer (or
 501 polyoxymethylene; POM) was found to provide a plausible explanation for the observed H₂CO parent scale length
 502 in comet 1P/Halley (Cottin et al. 2004), and for the heliocentric dependence of H₂CO production rates observed in
 503 O1/Hale-Bopp (Fray et al. 2006). The dissociation scale length of solid-phase POM depends strongly on the particle
 504 size and temperature (Fray et al. 2006), so the observed variations in $L_p(\text{H}_2\text{CO})$ can be explained as a result of
 505 differing size distributions for the POM-rich dust grains in the different comets observed to-date. Conversely, Milam
 506 et al. (2006) noted a lack of evidence for POM in Giotto mass spectra of comet Halley, and argued that it constitutes
 507 an unlikely source of H₂CO in the coma. Furthermore, mass spectrometry by the Rosetta mission found no evidence
 508 for POM in comet 67P (Hänni et al. 2022), so alternative explanations may be required to explain the distributed
 509 H₂CO source. Moreover, given the limited sample size and large disparity between the observed $L_p(\text{H}_2\text{CO})$ values, it
 510 will be important to conduct more observations of H₂CO distributions in different comets, over a range of coma size

scales, at differing heliocentric distances, to better characterize the behavior of the H₂CO source(s) and help constrain the properties of its still-elusive parent.

By simultaneously fitting the H₂CO parent scale length and production rate ratio relative to water, we derive an H₂CO abundance of $x = 0.153_{-0.023}^{+0.031}$ %, which is larger than the value of $x = 0.38 \pm 0.02$ % obtained from IRAM 30-m observations by Biver et al. (2021) using $L_p(\text{H}_2\text{CO}) = 5000$ km. This discrepancy is not surprising considering the retrieved H₂CO production rate from single-dish observations scales with the adopted L_p value. Using $L_p = 880$ km, the IRAM data are consistent with $x = 0.13 \pm 0.01$ %. The H₂CO production rate of $(9.1 \pm 0.9) \times 10^{24}$ s⁻¹ obtained by Coulson et al. (2020) using the JCMT assuming $L_p = 860$ km, also matches our derived H₂CO production rate. Our H₂CO abundance is the second-lowest value reported in a comet to-date, at radio wavelengths (Bockelée-Morvan & Biver 2017), but it should be remembered that the majority of those values were obtained using assumed (rather than directly measured) $L_p(\text{H}_2\text{CO})$ values. We therefore emphasize the importance of using an accurate parent scale length when deriving H₂CO production rates.

An upper limit of $x < 0.064$ % was obtained by Bonev et al. (2021) using Keck infrared observations, but this was calculated assuming H₂CO release directly from the nucleus (*i.e.* $L_p = 0$) — a common assumption among infrared spectroscopists (*e.g.* DiSanti et al. 2006; Dello Russo et al. 2016b) due to the difficulty of deriving L_p values from infrared spatial profiles — which inevitably underestimates the total coma H₂CO abundance in the presence of distributed sources. Our ALMA observations show no evidence for H₂CO production by the nucleus, with a (3σ) upper limit of 0.08 % on the abundance of any parent H₂CO, consistent with the Bonev et al. (2021) result. The relatively short H₂CO parent lifetimes derived using ALMA imply significant daughter production of H₂CO in the near-nucleus coma that could, in some situations, appear similar to a parent source. Further investigations of H₂CO spatial emission profiles at the angular scales of a few arcseconds probed by infrared spectroscopy are therefore warranted,

Despite the evidence for a lack of H₂CO in cometary ices inferred using ALMA observations, it remains one of the most widespread gas-phase molecules in the Galaxy. Abundances of H₂CO relative to water can reach several percent in warm protostellar gas (Ceccarelli et al. 2000; Ehrenfreund & Charnley 2000), while mid-infrared spectroscopy indicates possible abundances ~ 6 % in ices around low-mass protostars (Boogert et al. 2015). The lack of detectable H₂CO in cometary nuclei using ALMA therefore provides evidence for chemical processing to destroy H₂CO ice in (or during its passage to) the protosolar disk, which is expected to occur as part of the pathway to forming more complex organic molecules (including biologically relevant species), starting with hydrogenation of H₂CO to make CH₃OH ice (Chuang et al. 2016). Although formaldehyde is commonly detected in the gas-phase in protoplanetary disks (*e.g.* Pegues et al. 2020), total H₂CO/H₂O masses in the range $\sim 10^{-5}$ – 10^{-3} were recently measured in a sample of five disks by Guzmán et al. (2021), so relatively low H₂CO abundances in cometary nuclei ($\lesssim 6 \times 10^{-4}$, as found by our study and Bonev et al. 2021), may not be surprising.

Of relevance to the nature of the H₂CO parent, it is noteworthy that the H₂CO outflow velocity of 0.65 ± 0.02 km s⁻¹ in the sunward jet is significantly less than the HCN (parent molecule) outflow velocity of 0.74 km s⁻¹ measured only 2 hours earlier. Such a rapid drop in the jet velocity over this period seems unlikely, so this result may constitute evidence that the H₂CO parent is flowing radially outward at a slower rate than the gases sublimating directly from the nucleus. A similar effect is also observed for the daughter species CS (observed simultaneously with CH₃OH) as well as HNC (albeit, at lower confidence). The outflow velocities of coma dust grains are significantly lower than that of the gas particles due to the mass disparity between these fluids, which causes the dust to lag behind the gas (Crifo et al. 2004). A relatively low outflow velocity for H₂CO, CS and HNC may therefore indicate production of these molecules from a slower-moving dust precursor.

5.5. CS

The CS radical in cometary comae is commonly believed to originate from photodissociation of the CS₂ parent molecule (Jackson et al. 1982; Rodgers & Charnley 2006; Feldman et al. 2010). Our derived CS parent scale length of $L_p = 675_{-124}^{+159}$ km, at an outflow velocity of 0.73 km s⁻¹ would therefore correspond to a photodissociation rate $\Gamma(\text{CS}_2) = (1.08 \pm 0.20) \times 10^{-3}$ s⁻¹ at $r_H = 1.06$ au (or $(1.21 \pm 0.22) \times 10^{-3}$ s⁻¹ at $r_H = 1$ au). This value is consistent (within errors) with the value derived by Feldman et al. (1999) using spatially resolved HST STIS imaging of CS in comet C/1999 H1 (Lee). However, our inferred $\Gamma(\text{CS}_2)$ value is significantly smaller than the published rates: 2.9×10^{-3} s⁻¹ (Huebner & Mukherjee 2015), 1.9×10^{-3} s⁻¹ (Heays et al. 2017) and 1.7×10^{-3} s⁻¹ (Jackson et al. 1986), obtained using experimentally-derived CS₂ photodissociation cross sections. Even if only one of these rates is correct, the smaller value for the CS parent photodissociation rate obtained by our study implies that CS₂ photolysis

cannot be the main source of CS in the coma of 46P. This is similar to the conclusion of Roth et al. (2021a), who derived a CS parent photolysis rate (at $r_H = 1$ au) of $(3.6_{-2.2}^{+5.4}) \times 10^{-4} \text{ s}^{-1}$ in comet C/2015 ER61, which is even smaller than our value derived for 46P, although the rates remain consistent at the 2σ level. Biver et al. (2022) also derived a small CS parent photolysis rate of $(4-8) \times 10^{-4} \text{ s}^{-1}$ in C/2020 F3 (NEOWISE).

After considering the possible CS sources in comet C/2015 ER61, Roth et al. (2021a) determined that H_2CS was a possible parent for CS, but noted that the abundance of H_2CS in many comets is actually less than the CS abundance, so it cannot be the parent of all cometary CS. Indeed, the upper limit on the H_2CS abundance in comet 46P of < 0.016 % (Biver et al. 2021) is at odds with this hypothesis based on our derived CS (parent) abundance of $0.022_{-0.002}^{+0.004}$ %, as well as with respect to Biver et al. (2021)'s CS abundance of 0.028 ± 0.003 %. Based on its abundance upper limit of 0.07 % (Biver et al. 2021), OCS is worth considering as another possible CS parent, but this species can be ruled out due to its photodissociation rate of $9.6 \times 10^{-5} \text{ s}^{-1}$ (Huebner & Mukherjee 2015), which is too small to be compatible with the observed CS distribution. Other, larger carbon and sulfur-containing molecules should be considered as plausible candidates for the CS parent, such as CH_3SH , $\text{C}_2\text{H}_6\text{S}$, CH_4S_2 and $\text{C}_4\text{H}_6\text{S}$, which were detected using the ROSINA instrument at comet 67P (Calmonte et al. 2016; Hänni et al. 2022). However, since the photodissociation rates for these molecules are unknown, it remains to be seen if they could be consistent with the ALMA data. The difficulty of forming CS from any of these species in the inner coma should also be emphasized, due to the need to break multiple covalent bonds before CS can be liberated into the gas phase.

Further evidence against CS_2 as the parent of cometary CS was provided by the IRAM 30 m observations of comet 67P by Biver et al. (2023), who obtained a CS abundance of 0.05 ± 0.01 %. This is significantly larger than the CS_2 abundance measured by Rosetta during the previous apparition of 67P (0.02 %; Läuter et al. 2020), so assuming the coma chemistry was the same on the two apparitions, an additional source of CS is required in that comet.

The identity of the main CS parent in comets therefore remains elusive. Considering our present knowledge, the most plausible sources include dust grains rich in carbon and sulfur, or possibly large C and S-bearing molecules (such as $\text{H}_n\text{C}_m\text{S}_k$). Thermal degradation of large molecules or dust could also explain the variation in CS/HCN ratio as a function of heliocentric distance observed by Biver et al. (2006, 2011), assuming the CS production rate depends on the dust temperature. Future studies of the CS parent may benefit from the use of more physically realistic vectorial (or Monte Carlo) coma models, to account for the excess CS kinetic energy introduced during CS_2 photolysis, that might broaden its spatial distribution, leading to larger L_p values. The possibility that the CS_2 photolysis lifetime is larger than the currently-accepted value should also be considered.

The CS production rate in the 46P coma ($0.022_{-0.002}^{+0.004}$ % with respect to water), is at the lower end of the range of values previously found in comets ($0.02-0.20$ %; Bockelée-Morvan & Biver 2017).

5.6. HNC

Our HNC parent scale length of $1412_{-367}^{+14,876}$ km in 46P is comparable with the values of $700_{-400}^{+1,100}$ km in S1/ISON (Cordiner et al. 2014) and $3300_{-2800}^{+19,700}$ km in ER61 (Roth et al. 2021b). Unfortunately, the large error bars hinder the identification of any trends in $L_p(\text{HNC})$ with heliocentric distance or any other cometary parameters, so more observations at higher S/N will be needed to further elucidate the behavior of the HNC parent.

A strong variation in the HNC/HCN abundance ratio as a function of r_H was observed in a sample of 14 moderately active comets by Lis et al. (2008). This trend was interpreted as arising from variations in the HNC production rate as a function of coma dust grain temperature. Given the difficulty in producing HNC from known gas-phase chemical processes in the coma (Rodgers & Charnley 2001), it was therefore concluded that HNC is released from the thermal breakdown of macromolecules or polymeric material originating from inside the nucleus (see also Cordiner et al. 2017a). Our 46P HNC/HCN ratio of 5_{-2}^{+19} % is in line with the observed trend with r_H , so similar breakdown of nitrogen-rich particles could be responsible for the observed HNC in comet 46P. The precise nature of those particles, however, remains to be elucidated.

6. CONCLUSIONS

ALMA observations of comet 46P provided an unusually close-up view of the coma, allowing quantitative measurements of molecular production scale lengths for the first time in a Jupiter-family comet. The interferometric data were analysed using a 3D radiative transfer model, consisting of a radially-expanding ambient coma and a broad, (near-)sunward jet of half-opening angle $\theta_{jet} = 70^\circ$. HCN is identified as a parent species, with a production scale length $L_p < 3$ km (at 99% confidence) and no evidence for any production of this molecule in the coma. CH_3OH is

found to have comparable contributions from a nucleus source and a near-nucleus coma source (with $L_p = 36 \pm 7$ km in the jet; 20 ± 4 km in the ambient coma). The CH_3OH coma source is explained as most likely originating from the sublimation of dirty ice grains or larger ice chunks in the coma, consistent with the hyperactive nature of this comet. The CH_3CN data are also consistent with a parent model, with an upper limit of $L_p < 18$ km, demonstrating for the first time, that this species is most likely a parent molecule and originates from ices stored within the nucleus. We therefore conclude that measurements of HCN, CH_3OH and CH_3CN abundances in fully-activated comets provide a valid proxy for the abundances of these molecules within cometary nuclei.

The H_2CO , CS and HNC observations, on the other hand, cannot be reproduced using a parent model. Our models show that these species originate in the coma, either as photochemical daughters, or from the breakdown of macromolecules or dust grains, with L_p values in the range 550–16,000 km. The detection of a distributed H_2CO source with $L_p = 876_{-175}^{+250}$ km, combined with a lack of any detectable nucleus (parent) source for this molecule, is consistent with the non-detection of H_2CO from the 46P nucleus at infrared wavelengths.

Additional, spatial mapping of the distributed/daughter molecules in different comets (at various heliocentric distances), combined with laboratory studies of the dissociation rates of their putative parent species will be required to conclusively identify the parent materials. Given the consistency of the measured production scale lengths for H_2CO and HNC with previous ALMA observations of Oort cloud comets, our observations suggest that similar parent materials are present in JFCs and OCCs. Such parent materials must therefore be resistant to the thermal processing experienced by JFCs during their repeated perihelion passages.

The observed spectral and spatial data are consistent with an asymmetric coma, and we find enhanced outflow velocities and production rates for all species on the sunward (day side) of the comet. Conversely, the coma kinetic temperature is found to be significantly lower on the day side than the night side, which may be a result of enhanced adiabatic cooling rates on the day side.

The abundances of HCN and CH_3OH in 46P are consistent with average values found in OCCs and JFCs. On the other hand, CH_3CN , H_2CO and CS are at the lower end of the range of previously observed abundances, consistent with a depletion of these species (or their parents) in the 46P nucleus. We find that the abundances measured for daughter species (H_2CO , HNC and CS) are strongly dependent on the assumed parent scale length (L_p). We therefore emphasize the importance of using the correct L_p value when deriving the abundances of cometary daughter/product species, particularly for comets at relatively small geocentric distances, thus demonstrating a crucial need for further spatially resolved studies of cometary molecular emission.

This work was supported by the National Science Foundation under Grant No. AST-2009253 (MAC) and AST-2009398 (BPB). The work of MAC, SNM, NXR and SBC was also supported by NASA’s Planetary Science Division Internal Scientist Funding Program through the Fundamental Laboratory Research work package (FLaRe). This work makes use of ALMA dataset ADS/JAO.ALMA#2018.1.01114.S. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under NASA contract 80NM0018D0004. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

APPENDIX

REFERENCES

- A’Hearn, M. F., Belton, M. J. S., Delamere, W. A., et al. 2011, *Science*, 332, 1396, doi: [10.1126/science.1204054](https://doi.org/10.1126/science.1204054)
- Altwegg, K., Balsiger, H., Hänni, N., et al. 2020, *Nature Astronomy*, 4, 533, doi: [10.1038/s41550-019-0991-9](https://doi.org/10.1038/s41550-019-0991-9)
- Altwegg, K., Balsiger, H., & Fuselier, S. A. 2019, *ARA&A*, 57, 113, doi: [10.1146/annurev-astro-091918-104409](https://doi.org/10.1146/annurev-astro-091918-104409)
- Bardyn, A., Baklouti, D., Cottin, H., et al. 2017, *MNRAS*, 469, S712, doi: [10.1093/mnras/stx2640](https://doi.org/10.1093/mnras/stx2640)

- 659 Beer, E. H., Podolak, M., & Prialnik, D. 2006, *Icarus*, 180,
660 473, doi: [10.1016/j.icarus.2005.10.018](https://doi.org/10.1016/j.icarus.2005.10.018)
- 661 Bergman, P., Lerner, M. S., Olofsson, A. O. H., et al. 2022,
662 *A&A*, 660, A118, doi: [10.1051/0004-6361/202142583](https://doi.org/10.1051/0004-6361/202142583)
- 663 Bergner, J. B., Guzmán, V. G., Öberg, K. I., Loomis,
664 R. A., & Pegues, J. 2018, *ApJ*, 857, 69,
665 doi: [10.3847/1538-4357/aab664](https://doi.org/10.3847/1538-4357/aab664)
- 666 Biver, N., Bockelée-Morvan, D., Colom, P., et al. 2011,
667 *A&A*, 528, A142, doi: [10.1051/0004-6361/201016250](https://doi.org/10.1051/0004-6361/201016250)
- 668 Biver, N., Boissier, J., Bockelée-Morvan, D., et al. 2022,
669 *A&A*, 668, A171, doi: [10.1051/0004-6361/202244970](https://doi.org/10.1051/0004-6361/202244970)
- 670 Biver, N., Bockelée-Morvan, D., Crovisier, J., et al. 1999,
671 *AJ*, 118, 1850, doi: [10.1086/301033](https://doi.org/10.1086/301033)
- 672 —. 2006, *A&A*, 449, 1255, doi: [10.1051/0004-6361:20053849](https://doi.org/10.1051/0004-6361:20053849)
- 673 Biver, N., Bockelée-Morvan, D., Hofstadter, M., et al. 2019,
674 *A&A*, 630, A19, doi: [10.1051/0004-6361/201834960](https://doi.org/10.1051/0004-6361/201834960)
- 675 Biver, N., Bockelée-Morvan, D., Boissier, J., et al. 2021,
676 *A&A*, 648, A49, doi: [10.1051/0004-6361/202040125](https://doi.org/10.1051/0004-6361/202040125)
- 677 Biver, N., Bockelée-Morvan, D., Crovisier, J., et al. 2023,
678 *A&A*, 672, A170, doi: [10.1051/0004-6361/202245672](https://doi.org/10.1051/0004-6361/202245672)
- 679 Bockelée-Morvan, D., & Biver, N. 2017, *Philosophical*
680 *Transactions of the Royal Society of London Series A*,
681 375, 20160252, doi: [10.1098/rsta.2016.0252](https://doi.org/10.1098/rsta.2016.0252)
- 682 Bockelée-Morvan, D., Boissier, J., Biver, N., & Crovisier, J.
683 2010, *Icarus*, 210, 898, doi: [10.1016/j.icarus.2010.07.005](https://doi.org/10.1016/j.icarus.2010.07.005)
- 684 Boissier, J., Bockelée-Morvan, D., Biver, N., et al. 2007,
685 *A&A*, 475, 1131, doi: [10.1051/0004-6361:20078380](https://doi.org/10.1051/0004-6361:20078380)
- 686 Bonev, B. P., Disanti, M. A., Villanueva, G., et al. 2014,
687 *ApJL*, 796, L6
- 688 Bonev, B. P., Mumma, M. J., Kawakita, H., Kobayashi, H.,
689 & Villanueva, G. L. 2008, *Icarus*, 196, 241,
690 doi: [10.1016/j.icarus.2008.02.023](https://doi.org/10.1016/j.icarus.2008.02.023)
- 691 Bonev, B. P., Villanueva, G. L., Paganini, L., et al. 2013,
692 *Icarus*, 222, 740, doi: [10.1016/j.icarus.2012.07.034](https://doi.org/10.1016/j.icarus.2012.07.034)
- 693 Bonev, B. P., Dello Russo, N., DiSanti, M. A., et al. 2021,
694 *PSJ*, 2, 45, doi: [10.3847/PSJ/abd03c](https://doi.org/10.3847/PSJ/abd03c)
- 695 Boogert, A. C. A., Gerakines, P. A., & Whittet, D. C. B.
696 2015, *ARA&A*, 53, 541,
697 doi: [10.1146/annurev-astro-082214-122348](https://doi.org/10.1146/annurev-astro-082214-122348)
- 698 Booth, A. S., Walsh, C., Terwisscha van Scheltinga, J.,
699 et al. 2021, *Nature Astronomy*, 5, 684,
700 doi: [10.1038/s41550-021-01352-w](https://doi.org/10.1038/s41550-021-01352-w)
- 701 Brinch, C., & Hogerheijde, M. R. 2010, *A&A*, 523, A25,
702 doi: [10.1051/0004-6361/201015333](https://doi.org/10.1051/0004-6361/201015333)
- 703 Calmonte, U., Altwegg, K., Balsiger, H., et al. 2016,
704 *MNRAS*, 462, S253, doi: [10.1093/mnras/stw2601](https://doi.org/10.1093/mnras/stw2601)
- 705 Capaccioni, F., Coradini, A., Filacchione, G., et al. 2015,
706 *Science*, 347, aaa0628, doi: [10.1126/science.aaa0628](https://doi.org/10.1126/science.aaa0628)
- 707 CASA Team, Bean, B., Bhatnagar, S., et al. 2022, *PASP*,
708 134, 114501, doi: [10.1088/1538-3873/ac9642](https://doi.org/10.1088/1538-3873/ac9642)
- 709 Ceccarelli, C., Loinard, L., Castets, A., Tielens,
710 A. G. G. M., & Caux, E. 2000, *A&A*, 357, L9
- 711 Chuang, K. J., Fedoseev, G., Ioppolo, S., van Dishoeck,
712 E. F., & Linnartz, H. 2016, *MNRAS*, 455, 1702,
713 doi: [10.1093/mnras/stv2288](https://doi.org/10.1093/mnras/stv2288)
- 714 Cochran, A. L., Lévassieur-Regourd, A.-C., Cordiner, M.,
715 et al. 2015, *Space Sci. Rev.*, 197, 9,
716 doi: [10.1007/s11214-015-0183-6](https://doi.org/10.1007/s11214-015-0183-6)
- 717 Combi, M. R., Mäkinen, T., Bertaux, J. L., et al. 2020,
718 *PSJ*, 1, 72, doi: [10.3847/PSJ/abb026](https://doi.org/10.3847/PSJ/abb026)
- 719 Cordiner, M. A., & Charnley, S. B. 2021, *MNRAS*, 504,
720 5401, doi: [10.1093/mnras/stab1123](https://doi.org/10.1093/mnras/stab1123)
- 721 Cordiner, M. A., Remijan, A. J., Boissier, J., et al. 2014,
722 *ApJL*, 792, L2, doi: [10.1088/2041-8205/792/1/L2](https://doi.org/10.1088/2041-8205/792/1/L2)
- 723 Cordiner, M. A., Boissier, J., Charnley, S. B., et al. 2017a,
724 *ApJ*, 838, 147, doi: [10.3847/1538-4357/aa6590](https://doi.org/10.3847/1538-4357/aa6590)
- 725 Cordiner, M. A., Biver, N., Crovisier, J., et al. 2017b, *ApJ*,
726 837, 177, doi: [10.3847/1538-4357/aa6211](https://doi.org/10.3847/1538-4357/aa6211)
- 727 Cordiner, M. A., Palmer, M. Y., de Val-Borro, M., et al.
728 2019, *ApJL*, 870, L26, doi: [10.3847/2041-8213/aafb05](https://doi.org/10.3847/2041-8213/aafb05)
- 729 Cordiner, M. A., Milam, S. N., Biver, N., et al. 2020, *Nature*
730 *Astronomy*, 4, 861, doi: [10.1038/s41550-020-1087-2](https://doi.org/10.1038/s41550-020-1087-2)
- 731 Cordiner, M. A., Coulson, I. M., Garcia-Berrios, E., et al.
732 2022, *ApJ*, 929, 38, doi: [10.3847/1538-4357/ac5893](https://doi.org/10.3847/1538-4357/ac5893)
- 733 Cortes, P., Remijan, A., Hales, A., et al. 2022, *ALMA*
734 *Cycle 9 Technical Handbook*, 2022, *ALMA Technical*
735 *Handbook*, ALMA Doc. 9.3, ver. 1.0 ISBN
736 978-3-923524-66-2, doi: [10.5281/zenodo.7764458](https://doi.org/10.5281/zenodo.7764458)
- 737 Cottin, H., Bénilan, Y., Gazeau, M.-C., & Raulin, F. 2004,
738 *Icarus*, 167, 397, doi: [10.1016/j.icarus.2003.10.009](https://doi.org/10.1016/j.icarus.2003.10.009)
- 739 Cottin, H., & Fray, N. 2008, *Space Sci. Rev.*, 138, 179,
740 doi: [10.1007/s11214-008-9399-z](https://doi.org/10.1007/s11214-008-9399-z)
- 741 Coulson, I. M., Cordiner, M. A., Kuan, Y.-J., et al. 2017,
742 *AJ*, 153, 169, doi: [10.3847/1538-3881/aa6440](https://doi.org/10.3847/1538-3881/aa6440)
- 743 Coulson, I. M., Liu, F.-C., Cordiner, M. A., et al. 2020, *AJ*,
744 160, 182, doi: [10.3847/1538-3881/abafe0](https://doi.org/10.3847/1538-3881/abafe0)
- 745 Crifo, J. F., Fulle, M., Kömle, N. I., & Szego, K. 2004, in
746 *Comets II*, ed. M. C. Festou, H. U. Keller, & H. A.
747 Weaver, 471
- 748 Crovisier, J., & Encrenaz, T. 1983, *A&A*, 126, 170
- 749 Delahaye, T., Armante, R., Scott, N. A., et al. 2021,
750 *Journal of Molecular Spectroscopy*, 380, 111510,
751 doi: [10.1016/j.jms.2021.111510](https://doi.org/10.1016/j.jms.2021.111510)
- 752 Dello Russo, N., Kawakita, H., Vervack, R. J., & Weaver,
753 H. A. 2016a, *Icarus*, 278, 301,
754 doi: [10.1016/j.icarus.2016.05.039](https://doi.org/10.1016/j.icarus.2016.05.039)
- 755 Dello Russo, N., Vervack, R. J., Kawakita, H., et al. 2016b,
756 *Icarus*, 266, 152, doi: [10.1016/j.icarus.2015.11.030](https://doi.org/10.1016/j.icarus.2015.11.030)
- 757 DiSanti, M. A., Bonev, B. P., Magee-Sauer, K., et al. 2006,
758 *ApJ*, 650, 470, doi: [10.1086/507118](https://doi.org/10.1086/507118)

- 759 Disanti, M. A., Mumma, M. J., Dello Russo, N., et al. 1999,
760 Nature, 399, 662, doi: [10.1038/21378](https://doi.org/10.1038/21378)
- 761 Drahus, M., Jewitt, D., Guilbert-Lepoutre, A., Waniak, W.,
762 & Sievers, A. 2012, ApJ, 756, 80,
763 doi: [10.1088/0004-637X/756/1/80](https://doi.org/10.1088/0004-637X/756/1/80)
- 764 Dubernet, M. L., & Quintas-Sánchez, E. 2019, Molecular
765 Astrophysics, 16, 100046,
766 doi: [10.1016/j.molap.2019.100046](https://doi.org/10.1016/j.molap.2019.100046)
- 767 Ehrenfreund, P., & Charnley, S. B. 2000, ARA&A, 38, 427,
768 doi: [10.1146/annurev.astro.38.1.427](https://doi.org/10.1146/annurev.astro.38.1.427)
- 769 Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., &
770 Müller, H. S. P. 2016, Journal of Molecular Spectroscopy,
771 327, 95, doi: [10.1016/j.jms.2016.03.005](https://doi.org/10.1016/j.jms.2016.03.005)
- 772 Feldman, P. D., McCandliss, S. R., Morgenthaler, J. P.,
773 et al. 2010, ApJ, 711, 1051,
774 doi: [10.1088/0004-637X/711/2/1051](https://doi.org/10.1088/0004-637X/711/2/1051)
- 775 Feldman, P. D., Weaver, H. A., A’Hearn, M. F., et al. 1999,
776 in AAS/Division for Planetary Sciences Meeting
777 Abstracts, Vol. 31, AAS/Division for Planetary Sciences
778 Meeting Abstracts #31, 35.02
- 779 Fougere, N. 2014, PhD thesis, University of Michigan
- 780 Fougere, N., Combi, M. R., Rubin, M., & Tennishev, V.
781 2013, Icarus, 225, 688, doi: [10.1016/j.icarus.2013.04.031](https://doi.org/10.1016/j.icarus.2013.04.031)
- 782 Fougere, N., Combi, M. R., Tennishev, V., et al. 2012,
783 Icarus, 221, 174, doi: [10.1016/j.icarus.2012.07.019](https://doi.org/10.1016/j.icarus.2012.07.019)
- 784 Fray, N., Bénilan, Y., Biver, N., et al. 2006, Icarus, 184,
785 239, doi: [10.1016/j.icarus.2006.04.014](https://doi.org/10.1016/j.icarus.2006.04.014)
- 786 Fray, N., Bardin, A., Cottin, H., et al. 2016, Nature, 538,
787 72, doi: [10.1038/nature19320](https://doi.org/10.1038/nature19320)
- 788 Gkotsinas, A., Guilbert-Lepoutre, A., Raymond, S. N., &
789 Nesvorný, D. 2022, The Astrophysical Journal, 928, 43,
790 doi: [10.3847/1538-4357/ac54ac](https://doi.org/10.3847/1538-4357/ac54ac)
- 791 Gomes, R., Levison, H. F., Tsiganis, K., & Morbidelli, A.
792 2005, Nature, 435, 466, doi: [10.1038/nature03676](https://doi.org/10.1038/nature03676)
- 793 Gordon, I., Rothman, L., Hargreaves, R., et al. 2022,
794 Journal of Quantitative Spectroscopy and Radiative
795 Transfer, 277, 107949,
796 doi: <https://doi.org/10.1016/j.jqsrt.2021.107949>
- 797 Guzmán, V. V., Bergner, J. B., Law, C. J., et al. 2021,
798 ApJS, 257, 6, doi: [10.3847/1538-4365/ac1440](https://doi.org/10.3847/1538-4365/ac1440)
- 799 Hänni, N., Altwegg, K., Combi, M., et al. 2022, Nature
800 Communications, 13, 3639,
801 doi: [10.1038/s41467-022-31346-9](https://doi.org/10.1038/s41467-022-31346-9)
- 802 Hartogh, P., Crovisier, J., de Val-Borro, M., et al. 2010,
803 A&A, 518, L150, doi: [10.1051/0004-6361/201014665](https://doi.org/10.1051/0004-6361/201014665)
- 804 Haser, L. 1957, Bulletin de la Societe Royale des Sciences
805 de Liege, 43, 740
- 806 Heays, A. N., Bosman, A. D., & van Dishoeck, E. F. 2017,
807 A&A, 602, A105, doi: [10.1051/0004-6361/201628742](https://doi.org/10.1051/0004-6361/201628742)
- 808 Herbst, E., & van Dishoeck, E. F. 2009, ARA&A, 47, 427,
809 doi: [10.1146/annurev-astro-082708-101654](https://doi.org/10.1146/annurev-astro-082708-101654)
- 810 Huebner, W. F., & Mukherjee, J. 2015, Planet. Space Sci.,
811 106, 11, doi: [10.1016/j.pss.2014.11.022](https://doi.org/10.1016/j.pss.2014.11.022)
- 812 Ilee, J. D., Walsh, C., Booth, A. S., et al. 2021, ApJS, 257,
813 9, doi: [10.3847/1538-4365/ac1441](https://doi.org/10.3847/1538-4365/ac1441)
- 814 Jackson, W. M., Butterworth, P. S., & Ballard, D. 1986,
815 ApJ, 304, 515, doi: [10.1086/164185](https://doi.org/10.1086/164185)
- 816 Jackson, W. M., Halpern, J. B., Feldman, P. D., & Rahe, J.
817 1982, A&A, 107, 385
- 818 Kelley, M. S., Lindler, D. J., Bodewits, D., et al. 2013,
819 Icarus, 222, 634, doi: [10.1016/j.icarus.2012.09.037](https://doi.org/10.1016/j.icarus.2012.09.037)
- 820 Khan, Y., Gibb, E. L., Roth, N. X., et al. 2023, AJ, 165,
821 231, doi: [10.3847/1538-3881/acc074](https://doi.org/10.3847/1538-3881/acc074)
- 822 Knight, M. M., Schleicher, D. G., & Farnham, T. L. 2021,
823 PSJ, 2, 104, doi: [10.3847/PSJ/abef6c](https://doi.org/10.3847/PSJ/abef6c)
- 824 Läter, M., Kramer, T., Rubin, M., & Altwegg, K. 2020,
825 MNRAS, 498, 3995, doi: [10.1093/mnras/staa2643](https://doi.org/10.1093/mnras/staa2643)
- 826 Lis, D. C., Bockelée-Morvan, D., Boissier, J., et al. 2008,
827 ApJ, 675, 931, doi: [10.1086/527345](https://doi.org/10.1086/527345)
- 828 Lis, D. C., Bockelée-Morvan, D., Güsten, R., et al. 2019,
829 A&A, 625, L5, doi: [10.1051/0004-6361/201935554](https://doi.org/10.1051/0004-6361/201935554)
- 830 Loomis, R. A., Öberg, K. I., Andrews, S. M., et al. 2018,
831 AJ, 155, 182, doi: [10.3847/1538-3881/aab604](https://doi.org/10.3847/1538-3881/aab604)
- 832 McClure, M. K., Rocha, W. R. M., Pontoppidan, K. M.,
833 et al. 2023, Nature Astronomy,
834 doi: [10.1038/s41550-022-01875-w](https://doi.org/10.1038/s41550-022-01875-w)
- 835 Meech, K. J., & Svoren, J. 2004, in Comets II, ed. M. C.
836 Festou, H. U. Keller, & H. A. Weaver, 317
- 837 Meier, R., Eberhardt, P., Krankowsky, D., & Hodges, R. R.
838 1993, A&A, 277, 677
- 839 Mekhtiev, M. A., Godfrey, P. D., & Hougen, J. T. 1999,
840 Journal of Molecular Spectroscopy, 194, 171,
841 doi: [10.1006/jmsp.1998.7782](https://doi.org/10.1006/jmsp.1998.7782)
- 842 Milam, S. N., Remijan, A. J., Womack, M., et al. 2006,
843 ApJ, 649, 1169, doi: [10.1086/506501](https://doi.org/10.1086/506501)
- 844 Moulane, Y., Jehin, E., Manfroid, J., et al. 2023, A&A, 670,
845 A159, doi: [10.1051/0004-6361/202244779](https://doi.org/10.1051/0004-6361/202244779)
- 846 Mumma, M. J., & Charnley, S. B. 2011, ARA&A, 49, 471,
847 doi: [10.1146/annurev-astro-081309-130811](https://doi.org/10.1146/annurev-astro-081309-130811)
- 848 Mumma, M. J., Charnley, S. B., Cordiner, M., Paganini, L.,
849 & Villanueva, G. L. 2017, in AAS/Division for Planetary
850 Sciences Meeting Abstracts, Vol. 49, AAS/Division for
851 Planetary Sciences Meeting Abstracts #49, 414.19
- 852 Newville, M., Stensitzki, T., Allen, D. B., et al. 2016, Lmfit:
853 Non-Linear Least-Square Minimization and
854 Curve-Fitting for Python, Astrophysics Source Code
855 Library, record ascl:1606.014. <http://ascl.net/1606.014>
- 856 Öberg, K. I., & Bergin, E. A. 2021, PhR, 893, 1,
857 doi: [10.1016/j.physrep.2020.09.004](https://doi.org/10.1016/j.physrep.2020.09.004)

- 858 Öberg, K. I., Guzmán, V. V., Furuya, K., et al. 2015,
859 Nature, 520, 198, doi: [10.1038/nature14276](https://doi.org/10.1038/nature14276)
- 860 Pegues, J., Öberg, K. I., Bergner, J. B., et al. 2020, ApJ,
861 890, 142, doi: [10.3847/1538-4357/ab64d9](https://doi.org/10.3847/1538-4357/ab64d9)
- 862 Poch, O., Istiqomah, I., Quirico, E., et al. 2020, Science,
863 367, aaw7462, doi: [10.1126/science.aaw7462](https://doi.org/10.1126/science.aaw7462)
- 864 Protopapa, S., Kelley, M. S. P., Woodward, C. E., & Yang,
865 B. 2021, PSJ, 2, 176, doi: [10.3847/PSJ/ac135a](https://doi.org/10.3847/PSJ/ac135a)
- 866 Protopapa, S., Sunshine, J. M., Feaga, L. M., et al. 2014,
867 Icarus, 238, 191, doi: [10.1016/j.icarus.2014.04.008](https://doi.org/10.1016/j.icarus.2014.04.008)
- 868 Rodgers, S. D., & Charnley, S. B. 2001, MNRAS, 323, 84,
869 doi: [10.1046/j.1365-8711.2001.04099.x](https://doi.org/10.1046/j.1365-8711.2001.04099.x)
- 870 —. 2006, Advances in Space Research, 38, 1928,
871 doi: [10.1016/j.asr.2005.10.006](https://doi.org/10.1016/j.asr.2005.10.006)
- 872 Rodgers, S. D., Charnley, S. B., Huebner, W. F., & Boice,
873 D. C. 2004, Comets II, ed. G. W. Kronk, 505
- 874 Roth, N. X., Milam, S. N., Cordiner, M. A., et al. 2021a,
875 ApJ, 921, 14, doi: [10.3847/1538-4357/ac0441](https://doi.org/10.3847/1538-4357/ac0441)
- 876 —. 2021b, PSJ, 2, 55, doi: [10.3847/PSJ/abdd3a](https://doi.org/10.3847/PSJ/abdd3a)
- 877 Tenishev, V., Combi, M., & Davidsson, B. 2008, ApJ, 685,
878 659, doi: [10.1086/590376](https://doi.org/10.1086/590376)
- 879 Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H. F.
880 2005, Nature, 435, 459, doi: [10.1038/nature03539](https://doi.org/10.1038/nature03539)
- 881 van der Tak, F. F. S., Lique, F., Faure, A., Black, J. H., &
882 van Dishoeck, E. F. 2020, Atoms, 8, 15,
883 doi: [10.3390/atoms8020015](https://doi.org/10.3390/atoms8020015)
- 884 Villanueva, G. L., DiSanti, M. A., Mumma, M. J., & Xu,
885 L.-H. 2012, ApJ, 747, 37,
886 doi: [10.1088/0004-637X/747/1/37](https://doi.org/10.1088/0004-637X/747/1/37)
- 887 Wink, J., Bockelée-Morvan, D., Despois, D., et al. 1997,
888 Earth Moon and Planets, 78, 63,
889 doi: [10.1023/A:1006263026604](https://doi.org/10.1023/A:1006263026604)




Figure 8. SUBLIME model geometry used in the present study, showing the conical jet (Ω_1) and ambient coma (Ω_2), in cartesian coordinates. θ_{jet} is the half-opening angle of the conical jet, ϕ is the (phase) angle of the jet axis with respect to the observer (z), and ψ is the jet position angle in the plane of the sky (x is north, y is east).

A. SUBLIME MODEL GEOMETRY

Figure 8 shows a schematic diagram of the two-component coma radiative transfer model geometry used in the present study.

B. INTERFEROMETRIC VISIBILITY SPECTRAL MODELING

Figures 9 to 14 show the real part of the observed and modeled ALMA visibilities for comet 46P, as a function of spectral channel. The top panel of each figure shows the autocorrelation (total power) spectrum, while each successive sub-panel shows the data averaged over an range of increasing antenna baseline lengths (corresponding to decreasing angular scales). Figure 15 shows a zoomed, un-binned version of Figure 5 for CH_3OH , to highlight the difference between the two best-fitting models and the data.




Figure 9. $\text{CH}_3\text{OH } J_K = 5_K - 4_K$ binned visibility spectra (real J component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 10. HCN $J = 4 - 3$ binned visibility spectra (real component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 11. $\text{CH}_3\text{CN } J_K = 14_K - 13_K$ binned visibility spectra (real component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 12. $\text{H}_2\text{CO } J_{K_a, K_c} = 5_{1,5}-4_{1,4}$ binned visibility spectra (real component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 13. $\text{CS } J = 5-4$ binned visibility spectra (real component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 14. HNC $J = 4-3$ binned visibility spectra (real component), with best-fitting model overlaid in red. Each sub-panel shows the average spectrum for a range of uv distances (antenna baselines), given upper-left, which correspond to a range of angular scales (θ) on the sky.




Figure 15. Observed, spectrally averaged real interferometric visibilities for CH_3OH , zoomed in on the region of shortest baselines to show the improved fit of the parent + daughter model (blue circles) compared with the pure parent model (orange circles).