

SPACEFLIGHT-INDUCED CHANGES IN MICROBIAL VIRULENCE AND THE IMPACT TO THE HOST IMMUNE RESPONSE

C.M. Ott¹, J. Barrila², S. Koroli², S. Gangaraju², R.R. Davis², S.G. Thornhill³, J. Yang², K.A. Hoffman⁴, R.A. Rice⁴, A.A. Medina-Colorado⁴, C. Vu², B.Y. Kang², L.L. Banken^{2,5}, P. Stafford⁶, C. Oubre¹, B.E. Crucian¹, and C.A. Nickerson^{2,5}

¹NASA Johnson Space Center, Houston, TX, ²Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University, Tempe, AZ, ³JES Tech, Houston, TX, ⁴KBR, Houston, TX, ⁵School of Life Sciences, Arizona State University, Tempe, AZ; ⁶Biodesign Center for Innovations in Medicine, Arizona State University, Tempe, AZ

Many microbial pathogens have repeatedly exhibited unexpected responses relevant to infectious disease when grown in microgravity and microgravity analogue environments, including changes in final cell concentration, biofilm production, stress resistance, antibiotic sensitivity, gene expression, host-pathogen interactions, and virulence. Notably, the classic foodborne pathogen *Salmonella enterica* serovar Typhimurium displayed increased virulence in animals when cultured in either the spaceflight analogue or true spaceflight environment. Recently, *Serratia marcescens* also was shown to increase virulence when cultured in the spaceflight environment. In parallel, astronaut studies have characterized a persistent spaceflight-induced dysregulation of the human immune system at multiple levels, which suggests an increased risk of infectious diseases. Moreover, astronauts have some degree of clinical infectious disease incidence. However, the contribution of the microgravity environment on host-pathogen interactions and potential for clinical disease remains understudied and poorly characterized.

The goal of this study is to gain insight into the breadth of other medically significant microbial pathogens that may exhibit altered virulence and pathogenesis-related responses when cultured in spaceflight analogue conditions. Specifically, we are characterizing the effect of spaceflight analogue culture (Low Shear Modeled Microgravity/LSMMG) on microbial pathogenesis-related stress responses, *in vitro* host-pathogen interactions, gene expression, and virulence potential in animals using five important model bacterial pathogens, *Salmonella enterica* Enteritidis, *Pseudomonas aeruginosa*, *Burkholderia cepacia*, *Streptococcus pneumoniae*, and enterohemorrhagic *Escherichia coli*.

The information to date is providing a better understanding into the potential impact of microgravity on alterations in microbial virulence and associated infectious disease risk to crew health during spaceflight missions.