

Xenon in Benu, Ryugu, and Winchcombe

S. A. Crowther¹, J. D. Gilmour¹, J. S. Cowpe¹, L. Fawcett¹, J. J. Barnes², A. N. Nguyen³, H. C. Connolly Jr^{2,4,5}, D. S. Lauretta², R. Okazaki⁶, Hayabusa2-initial-analysis volatiles team and Hayabusa2-initial-analysis core.

¹Department of Earth and Environmental Sciences, The University of Manchester, Manchester, UK (sarah.crowther@manchester.ac.uk); ²Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ, USA; ³ARES, NASA JSC, Houston, TX, USA; ⁴Department of Geology, Rowan University, Glassboro, NJ, USA; ⁵Department of Earth and Planetary Sciences, American Museum of Natural History, New York, NY, USA; ⁶Department of earth and Planetary Sciences, Kyushu University, Japan.

Carbonaceous chondrites sample some of the most primitive materials from our Solar System and are implicated in the delivery of water, organics and volatile species to Earth [1]. However, exposure to the terrestrial atmosphere can lead to alteration or loss of information about their composition and history [2]. Returned samples, such as those collected from asteroids Ryugu and Benu by JAXA's Hayabusa2 [3] and NASA's OSIRIS-REx [4] missions, respectively, are protected from the terrestrial atmosphere, allowing more accurate interpretation. Careful curation of the Winchcombe (CM2) carbonaceous chondrite [5,6] has provided a near-pristine meteorite to compare with returned samples.

Xenon is particularly useful among the noble gases for tracing Solar System evolution. Nine isotopes are sufficient to unambiguously identify multiple sources contributing to the observed composition: different reservoirs, radioactive decay of ²⁴⁴Pu and ²³⁸U, spallation and secondary neutron capture reactions of samples containing Ba and light REE, and artificial neutron irradiation of samples containing ¹²⁷I, ¹³⁰Ba, ¹³⁰Te and ²³⁵U, all produce characteristic xenon isotopic signatures. The I-Xe chronometer [7,8] provides a high-resolution means to investigate the timing and sequence of events in the first ~70 Myr of Solar System history. Here we report and compare xenon isotopic analyses of Ryugu, Benu and Winchcombe, conducted using the RELAX mass spectrometer [9,10]. The high sensitivity of RELAX enables us to separate different components and observe small isotopic excesses which may not be resolved using a conventional noble gas mass spectrometer.

Funding: STFC ST/V000675/1 and ST/Y002369/1, NASA Contract NNM10AA11C.

[1] Alexander C.M.O'D. et al. (2012) *Science*, 337, 721-723.

[2] Bland P. et al. (2006) *Meteorites and the Early Solar System II*, 853-867.

[3] Watanabe Si. et al. (2017) *Space Sci. Rev.*, 208, 3-16.

[4] Lauretta D.S. & Connolly H.C. Jr. et al. (2024) *MAPS*, 59, 2453-2486.

[5] King A.J. et al. (2022) *Sci. Adv.*, 8, eabq3925.

[6] Russell S.S. et al. (2024) *MAPS*, 59, 973-987.

[7] Gilmour J.D. et al. (2006) *MAPS*, 41, 19-31.

[8] Pravdivtseva O. et al. (2017) *GCA*, 201, 320-330.

[9] Crowther S.A. et al. (2008) *JASS*, 23, 938-947.

[10] Gilmour J.D. et al. (1994) *Rev. Sci. Instru.*, 65, 617-625.