

Trends (2000 to 2022) from TOAR II/HEGIFTOM Global Ground-based Tropospheric Measurements: A Reference Dataset for Satellite Products & Models

Anne M. Thompson^{*1,2}, **Debra E. Kollonige^{1,3}**, **Ryan M. Stauffer¹**

Roeland Van Malderen⁴, **Herman GJ Smit^{5,6}**, **Bryan J. Johnson⁷**, **Kai-Lan Chang^{6,8}**

(1) Atmospheric Chemistry & Dynamics Lab, NASA/GSFC, Greenbelt, MD; (2) GESTAR, Univ. of Maryland, Baltimore County, MD; (3) Science Systems & Applications, Inc, Lanham, MD;



(4) RMI, Uccle, Belgium; (5) FZ-Juelich, Germany; (6) CIRES, U. of Colorado, Boulder, CO;
(7) NOAA/Global Monitoring Lab, Boulder, CO; (8) NOAA Chemical Sciences Lab, Boulder, CO

*** anne.m.thompson@nasa.gov**

SHADOZ=So. Hemisphere Additional Ozonesondes
IAGOS = In-service Aircraft for a Global Observing System
HEGIFTOM = Harmonization and Evaluation of Ground-based Instruments for Free Tropospheric Ozone Measurements

- **WHY Is HEGIFTOM** (Harmonization and Evaluation of Ground-based Instruments for Free Tropospheric Ozone Measurements) **so important in Ozone (TOAR II) & Climate Assessments?**
- **HEGIFTOM: WHAT, HOW, WHERE. Data Status.**
- **Present “Total tropospheric ozone column” (TrOC, surface to 300 hPa) trends**
 - Focus on 55 station/instrument time-series from 2000-2022
 - Five instrument types with TrOC measurements included in calculations
 - Compare trends by 2 standard statistical methods (QR, MLR)
 - Free Tropospheric (FT) ozone column trends computed (not shown)
- **Summary: Trends to date for 2000-2022 show:**
 - *All sites within ± 3 ppbv/dec $\rightarrow$ equivalent to $\pm(1-8)\%/dec$, for TrOC, depending on location, and independent of statistical method*
 - *HEGIFTOM data = *the* independent reference for satellite, model evaluation*

Why Does IGAC/TOAR II Need HEGIFTOM?



- Tropospheric ozone (TCO) satellite products ([Keppens, next Talk](#)) struggle to match aircraft, ozonesondes, each other! New tropical comparisons (Gaudel et al., 2024 for TOAR II) illustrate noise, varying biases, correlation, r^2 0.3-0.6 (**Left**)
- Typical CCM puts positive FT O_3 trend, greatest radiative forcing region, in wrong months. BL O_3 trend too low. (**Right**, update of Stauffer et al., 2019)

HEGIFTOM: IGAC/TOAR II Activity, Co-Leads: R. Van Malderen & H. G. J. Smit

Alternative to still-evolving satellite TrOC (tropospheric ozone column) products:

- Ozone from 5 ground-based instrument types, most from NDACC & related networks: in-service aircraft [IAGOS], ozonesondes, FTIR, Brewer/Dobson Umkehr, Lidar (**Photos, Right**)
 - All instrument types in HEGIFTOM database. *Reprocessed data based on rigorous protocols and absolute standards*, thus ensuring harmonized time-series with minimal artifacts.
- Contributing networks
- Each measurement is delivered with uncertainty and a quality flag




IAGOS



Ozonesondes




Brewer/Dobson Umkehr




FTIR



Lidar




TrOC = surface -300 hPa
 } FT, 700-400 hPa



<http://hegiftom.meteo.be/datasets>

HEGIFTOM Sites/Datasets for 2000-2022 Trends

Global Sites Contributing to HEGIFTOM (55 L3 Data) Trends





TOAR II Protocol: Minimum Sample No. to reduce trends uncertainty, start in 2000-2002, end 2020-2022
 QR (Quantile Regression) analyses use all data (“L1”) from **55** sites; **only 50%-ile results shown**.
 Monthly means (L3) from **55** sites are analyzed with QR and MLR (multiple linear regression)

Trends Questions Addressed with HEGIFTOM Data

1. What do TrOC trends (surface to 300 hPa) for 2000-2022 look like? *Examine “all-site” QR trends & median distribution at 55 sites*
2. How do TrOC trends computed with QR and MLR compare? *Answer with analysis of L3 (monthly mean) TrOC from 55 stations*
3. How do TrOC trends from the various instrument types (sondes, IAGOS, FTIR, Dobson Umkehr, Lidar) compare? *Examine trends at colocated stations*
4. How do TrOC trends vary by region? *Examine trends on map*


Ques 1. TrOC QR “All-Site” Trends, 5 Instrument Types





- **Left:** QR L1 trends for TrOC in TOAR-preferred ppbv/decade (2σ). (1) **Median trends nearly all within +/- 3 ppbv/decade.** (2) Medians for most stations ~ 0 . (3) Positive and negative trends appear at all longitudes
- **Right:** Medians in %/decade suggest possible instrument bias. More negatives than positives

Ques 2. TrOC QR, MLR Trends Similar

HEGIFTOM QR and MLR Trends (2000-2022) in L3 TrOC (ppbv/decade) for surface to 300hPa



Global HEGIFTOM QR/MLR Trends (2000-2022) in L3 TrOC (ppbv/decade) for surface to 300hPa





QR-MLR trends, ppbv/dec



- **Left & Center:** TrOC trends, QR= color-coded for 5 instruments. MLR= green shades for 5 instrument types. QR trends tend to be higher than MLR (**Right**)
- **Left:** Preponderance of negative trends in No. Hemisphere. **Center:** few Asian data

Ques 2, Cont'd. TrOC QR, MLR Trends Similarity



QR, MLR trends,
+/-3 ppbv/decade
range



QR, MLR trends,
%/decade,
-6 to +8%/dec
range




Ques 3. TrOC Trends from 5 Instruments Compared



- **Left:** L1 QR trends, 4 of 6 sites with > 1 instrument show large offsets: MLO/Hilo, Boulder, Izaña, Lauder (Zeng et al., 2024; Björklund et al., 2024)
- **Right:** Pronounced offsets observed at same sites (MLO/Hilo, Boulder, Lauder) with both statistical methods. Causes unclear – different sampling frequency, protocols, diurnal, seasonal effects investigated

Ques 4. TrOC Trends – Global Map View



Summarizes: 55 station Trends, moderately positive to negative trends with more confidence (lower p-value) in no. hemisphere (NH). More negative trends in NH. Large divergences at multi-instrument stations. Lack of SH and Asian data in HEGIFTOM limits “global” assessment

Conclusions – HEGIFTOM TOAR II Trends

SUMMARY

- TrOC trends (2000-2022) determined from 55 Ground-based Instruments display *moderately positive and negative* median trends; many with no detectable change!
- Results are independent of QR, MLR statistical method. **D. Kollonige SHADOZ talk illustrates outstanding MLR application to equatorial SE Asia trends!**

SIGNIFICANCE: HEGIFTOM data are *the* Definitive TOAR Reference. Use them!

- Recommend TOAR Model Comparison project with these values, site by site
- If model (or satellite TOC) disagree, HEGIFTOM can guide improvements

WORK IN PROGRESS:

- Understand instrument trend offsets
- Nearby site trend disagreements- why?
- Investigate COVID-19 impact on trends
- “Merge” stations for robust “regional trends” – examine 5%-ile, 95%-ile trends

Thank you! Acknowledgments. Bibliography

- **Acknowledgments: Dozens of funding organizations. Hundreds of researchers who have operated and collected ozone ground-based data over the past 30 years! Ozonesonde Funding by NASA UACO (K. Jucks), SAGE III (R. Eckman) and NOAA/OAR - GML**

Chang, K-L. et al. (2024) Challenges of detecting free tropospheric ozone ...<https://doi.org/10.5194/egusphere-2023-2739>

Gaudel, A., et al. (2018) Tropospheric Ozone Assessment Report: Present-day distribution and trends...
<https://doi.org/10.1525/elementa.291>

Gaudel, A., et al. (2024) Tropical tropospheric ozone distribution and trends from in situ..., *ACP*,
<https://doi.org/10.5194/egusphere-2023-3095>

Kollonige, D. E., et al. (2024) SHADOZ ozonesonde network – Overview and updates, This Meeting, Talk Today

Stauffer, R. M., et al. (2019) The effects of a changing observing system on MERRA-2-based ozone ...
<https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2018JD030090>

Stauffer, R. M., et al. (2024) Dynamical drivers of free-tropospheric ozone... *ACP* <https://doi.org/10.5194/egusphere-2023-2618>

Smit, H. G. J., A. M. Thompson et al. (2021) WMO/GAW ASOPOS Report 268 <https://library.wmo.int/records/item/57720-ozonesonde-measurement-principles-and-best-operational-practices>


Smit, H. G. J. et al. (2024) New insights from the Juelich OzoneSonde ... <https://doi.org/10.5194/amt-17-73-2024>

Thompson, A. M., et al. (2021) Regional and seasonal trends in tropical ozone from SHADOZ profiles...
<https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021JD034691>




HEGIFTOM Trends. Input & Guidelines

Nairobi Annual Trend in TrOC (DU or ppbv/decade) for surface -> 150hPa



Largest Increase at Low O_3

Nairobi Tropospheric Column MLR Trends (1998–2021)




Large Mar-Apr Increase masked by negligible annual mean trend in O_3

- Recommended TOAR II statistical approach is Quantile Regression (QR) with NOAA-provided test code, e.g., K-L Chang et al., (2023; JGR; 10.1029/2022JD038090)
- Alternative: Multiple-Linear Regression (MLR) as used in Thompson et al., 2021 & Stauffer et al., *ACP*, 2024. MLR is standard of stratospheric ozone Assessment community
- Above example for a typical SHADOZ station shows merits of each approach. QR gives insights into low-mid-ozone- O_3 profiles. Monthly means from MLR give insight into meteorological or chemical signatures responsible for O_3 trends



Tropospheric ozone column: COVID impact



○ ozonesondes

△ Umkehr

◇ FTIR

□ IAGOS

▽ Lidar

- Relative change of mean TrOC for the time period 2000-2022 vs. 2000-2019

Blue: 2000-2022 < 2000-2019

Red: 2000-2022 > 2000-2019

- Decline in 75% of the sites, on average -0.3% prominent in NH (spring + summer), stronger in FT.
- From R. Van Malderen, RMI
- Impact on trends!

Ques 3. All-Site FTOC Median Trend Variability

- **Left:** Sites (sondes, IAGOS, Umkehr, lidar), **50-%ile median** profiles, L1 analyzed with QR, 2000-2022. **Upper Right:** FT OC, 700-300 hPa, mostly smaller than TrOC trends; several exceptions in tropics.
- **Changes mostly ≤ 2.0 ppbv/dec, positive OR negative**

Global Median All Instruments L1 Trend (2000-2022) in TrOC (ppbv/decade) for FT (300hPa<p<700hPa)

